

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 July 2003 (03.07.2003)

PCT

(10) International Publication Number
WO 03/053349 A2

(51) International Patent Classification⁷: **A61K**

(21) International Application Number: PCT/US02/39926

(22) International Filing Date:
13 December 2002 (13.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/344,080 20 December 2001 (20.12.2001) US
60/382,103 20 May 2002 (20.05.2002) US

(71) Applicant: **BRISTOL-MYERS SQUIBB COMPANY**
[US/US]; P.O. Box 4000, Route 206 and Provinceline
Road, Princeton, NJ 08543-4000 (US).

(72) Inventors: **CAMPBELL, Jeffrey, Allen**; 525 Alpine
Drive, Cheshire, CT 06410 (US). **GOOD, Andrew,**
Charles; 52 High Hill Road, Wallingford, CT 06492 (US).

(74) Agents: **DUBOFF, Samuel** et al.; Bristol-Myers Squibb
Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INHIBITORS OF HEPATITIS C VIRUS

(57) Abstract: The present invention relates to macrocyclic compounds, methods for making these compounds, pharmaceutical compositions and the therapeutic or prophylactic use of these compounds by administering said compounds to mammals to prevent or treat hepatitis C virus (HCV) infection.

WO 03/053349 A2

INHIBITORS OF HEPATITIS C VIRUS**REFERENCE TO RELATED APPLICATIONS**

5 This application claims the benefit of U.S.
Provisional Application Serial Numbers 60/382,103 filed
May 20, 2002 and 60/344,080 filed December 20, 2001.

BACKGROUND OF THE INVENTION

10

Hepatitis C virus (HCV) is the major etiological agent of 90% of all cases of non-A, non-B hepatitis (Choo et al., 1989, Kuo et al., 1989). The incidence of HCV infection is becoming an increasingly severe public
15 health concern with 2-15% individuals infected worldwide. While primary infection with HCV is often asymptomatic, most HCV infections progress to a chronic state that can persist for decades. Of those with chronic HCV infections, it is believed that about 20-50% will
20 eventually develop chronic liver disease (e.g. cirrhosis) and 20-30% of these cases will lead to liver failure or liver cancer. As the current HCV-infected population ages, the morbidity and mortality associated with HCV are expected to triple.

25

An approved treatment for HCV infection uses interferon (IFN) which indirectly effects HCV infection by stimulating the host antiviral response. IFN treatment is suboptimal, however, as a sustained antiviral response
30 is produced in less than 30% of treated patients. Further, IFN treatment induces an array of side effects of varying severity in upwards of 90% of patients (eg: acute pancreatitis, depression, retinopathy,

thyroiditis). Therapy with a combination of IFN and ribavirin has provided a slightly higher sustained response rate, but not alleviated the IFN-induced side effects.

5

The use of protease inhibitors, particularly those selectively targeting HCV serine protease, has great potential to be useful in treating HCV infections in patients by inhibiting HCV replication.

10

Amongst the compounds that have demonstrated efficacy in inhibiting HCV replication, as selective HCV serine protease inhibitors, are the macrocyclic peptide compounds disclosed in International Application PCT/CA00/00353 (Publication No. WO 00/59929). The present invention describes selective inhibitors of the HCV NS3/NS4A serine protease complex of improved chemical design relative to the previously described invention, that have the potential to demonstrate suitable whole cell permeability for the treatment of HCV-infected patients.


15

20

SUMMARY OF THE INVENTION

25

The present invention relates to compounds of Formula I, including pharmaceutically acceptable salts thereof,



wherein:

5

(a) R_1 is C_{1-6} alkyl or unsubstituted C_{3-7} cycloalkyl;

10

(b) R_2 is H, halo, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkoxy, C_6 or 10 aryl or heterocycle; wherein heterocycle is a five-, six-, or seven-membered saturated or unsaturated heterocycle containing from one to four heteroatoms selected from nitrogen, oxygen and sulfur; said aryl or heterocycle being substituted with R_{24} ; wherein R_{24} is H, halo, C_{1-6} alkyl, $-N(R_a)(R_b)$, $-N(R_a)C(O)(OR_d)$, $-N(R_a)C(O)(R_b)$, $-NHC(O)(R_a)(R_c)$ or $-N(R_a)SO_2R_b$; wherein R_a and R_b are independently H, C_{1-6} alkyl or C_{3-6} cycloalkyl; R_c is H, C_{1-6} alkyl, C_{3-6} cycloalkyl or C_{1-6} alkoxy; R_d is C_{1-6} alkyl or C_{3-6} cycloalkyl;

20

(c) R_3 is H, halo, CF_3 , C_{1-6} alkoxy or C_{3-6} cycloalkoxy;

- (d) R_4 is NH_2 , or $-NH-R_{31}$; wherein R_{31} is $-C(O)-R_{32}$, $C(O)-NHR_{32}$ or $C(O)-OR_{32}$, wherein R_{32} is C_1-C_6 alkyl optionally substituted with halo, $-(CH_2)_p-C_3-7$ cycloalkyl or a tetrahydrofuran ring linked through the C3 or C4 position of the ring; wherein p is 0-6;
- (e) R_5 is H; and
- (f) Q is a three to nine atom saturated or unsaturated alkylene chain optionally containing one to three heteroatoms independently selected from O or $S(O)_m$; wherein m is 0, 1 or 2.

The present invention also relates to a pharmaceutical composition, useful for inhibiting HCV NS3 protease, or for treating patients infected with the hepatitis C virus, comprising a therapeutically effective amount of a compound of the present invention and a pharmaceutically acceptable carrier.

The present invention further relates to a method of treating mammals infected with hepatitis C virus, comprising administering to said mammal an effective amount of a compound of the present invention.

Additionally, the present invention relates to a method of inhibiting HCV NS3 protease by administering to a patient an effective amount of a compound of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

As used herein, the following definitions apply unless otherwise noted. With reference to the instances
5 where (R) or (S) is used to designate the configuration of a substituent in context to the whole compound and not in context to the substituent alone.

The term "halo" as used herein means a halogen
10 substituent selected from bromo, chloro, fluoro or iodo.

The term "C₁₋₆ alkyl" as used herein means acyclic, straight or branched chain alkyl substituents containing from 1 to six carbon atoms and includes, for example,
15 methyl, ethyl, propyl, butyl, tert-butyl, hexyl, 1-methylethyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl.

The term "C₁₋₆ alkoxy" as used herein means the
20 radical -O(C₁₋₆ alkyl) wherein alkyl is as defined above containing up to six carbon atoms. Alkoxy includes, for example, methoxy, ethoxy, propoxy, 1-methylethoxy, butoxy and 1,1-dimethylethoxy. The latter radical is known commonly as tert-butoxy.

25

The term "C₃₋₇ cycloalkyl" as used herein means a cycloalkyl substituent containing from three to seven carbon atoms and includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

30


The term "C₃₋₇ cycloalkoxy" as used herein means a C₃₋₇ cycloalkyl group linked to an oxygen atom, such as, for example, butyloxy or cyclopropyloxy.

The term "C₆ or 10 aryl" as used herein means either an aromatic monocyclic group containing 6 carbon atoms or an aromatic bicyclic group containing 10 carbon atoms, for example, aryl includes phenyl, 1-naphthyl or 2-naphthyl.

The term "carboxy(C₁₋₆ alkyl)" as used herein means a carboxyl group (COOH) linked through a C₁₋₆ alkyl group as defined above and includes, for example, butyric acid.

The term "heterocycle", as used herein means a monovalent radical derived by removal of a hydrogen from a five-, six-, or seven-membered saturated or unsaturated (including aromatic) heterocycle containing from one to four heteroatoms selected from nitrogen, oxygen and sulfur. Furthermore, the term heterocycle includes heterocycles, as defined above, that are fused to one or more other ring structure. Examples of suitable heterocycles include, but are not limited to, pyrrolidine, tetrahydrofuran, thiazolidine, pyrrole, thiophene, diazepine, 1H-imidazole, isoxazole, thiazole, tetrazole, piperidine, 1,4-dioxane, 4-morpholine, pyridine, pyrimidine, thiazolo[4,5-b]-pyridine, quinoline, or indole, or the following heterocycles:


25



The term "C₁₋₆ alkyl-heterocycle" as used herein, means a heterocyclic radical as defined above linked through a chain or branched alkyl group, wherein alkyl as


defined above containing from 1 to 6 carbon atoms.

Examples of C₁₋₆ alkyl-Het include:



5

In a preferred embodiment, the compounds of the present invention have the structure of Formula II




Formula II

10 wherein R₁, R₂, R₃, R₄, R₅ and Q are as defined for Formula I.

For compounds of Formula II, it is preferred that R₁ is cyclopropyl. In addition, it is preferred that R₂ is
 15 phenyl. It is further preferred that R₃ is methoxy. Furthermore, it is further preferred that R₄ is NH₂ or *tert*-butoxycarbonylNH-. It is also preferred that Q is -(CH₂)_nCH=C- wherein n is 1-5. It is more preferred that n is 3 or 4.

20

In a more preferred embodiment, the compounds of the present invention have the structure of Formula III





III

5

wherein R_4 is NH_2 or *tert*-butoxycarbonylNH-, Q is $(CH_2)_nCH=C-$, and n is 1-5. For the compounds of Formula III, it is more preferred that n is 3 or 4.

10 In another preferred embodiment for compounds of Formula II, wherein:



and R_1 is cyclopropyl.

Compounds of the present invention, by virtue of
20 their basic moiety, can form salts by the addition of a

pharmaceutically acceptable acid. The acid addition salts are formed from a compound of Formula I and a pharmaceutically acceptable inorganic acid, including but not limited to hydrochloric, hydrobromic, hydroiodic, 5 sulfuric, phosphoric, or organic acid such as p-toluenesulfonic, methanesulfonic, acetic, benzoic, citric, malonic, fumaric, maleic, oxalic, succinic, sulfamic, or tartaric. Thus, examples of such pharmaceutically acceptable salts include chloride, 10 bromide, iodide, sulfate, phosphate, methanesulfonate, citrate, acetate, malonate, fumarate, sulfamate, and tartrate.


Salts of an amine group may also comprise quaternary 15 ammonium salts in which the amino nitrogen carries a suitable organic group such as an alkyl, alkenyl, alkynyl or aralkyl moiety.

Compounds of the present invention, which are 20 substituted with a acidic group, may exist as salts formed through base addition. Such base addition salts include those derived from inorganic bases which include, for example, alkali metal salts (e.g. sodium and potassium), alkaline earth metal salts (e.g. calcium and 25 magnesium), aluminum salts and ammonium salts. In addition, suitable base addition salts include salts of physiologically acceptable organic bases such as trimethylamine, triethylamine, morpholine, pyridine, piperidine, picoline, dicyclohexylamine, N,N'- 30 dibenzylethylenediamine, 2-hydroxyethylamine, bis-(2-hydroxyethyl)amine, tri-(2-hydroxyethyl)amine, procaine, dibenzylpiperidine, N-benzyl- β -phenethylamine, dehydroabietylamine, N,N'-bishydroabietylamine,


glucamine, N-methylglucamine, collidine, quinine, quinoline, ethylenediamine, ornithine, choline, N,N'-benzylphenethylamine, chlorprocaine, diethanolamine, diethylamine, piperazine, tris(hydroxymethyl)aminomethane
5 and tetramethylammonium hydroxide and basic amino acids such as lysine, arginine and N-methylglutamine. These salts may be prepared by methods known to those skilled in the art.

10 In addition, compounds of the present invention, or a salt thereof, may exhibit polymorphism. The present invention also encompasses any such polymorphic form.


Compounds of the present invention (Formula I, II or
15 III) also contain two or more chiral centers and exist in different optically active forms. For example, compounds of Formula I may include a cyclopropyl of formula



20 wherein C₁ and C₂ each represent an asymmetric carbon atom at positions 1 and 2 of the cyclopropyl ring. Notwithstanding other possible asymmetric centers at other segments of the compounds of formula I, the presence of these two asymmetric centers means that the compounds of
25 formula I can exist as mixtures of diastereomers, such as the diastereomers of compounds of Formula II wherein Q is configured either syn to the amide or syn to the carbonyl as shown below.




syn to carbonyl




syn to amide

Alternatively, the structures can be viewed as:



syn to carbonyl



syn to amide

The present invention includes both enantiomers and mixtures of enantiomers such as racemic mixtures.

5

As illustrated in the examples, the racemic mixtures can be prepared and thereafter separated into individual optical isomers, or these optical isomers can be prepared by chiral synthesis.

10

The enantiomers may be resolved by methods known to those skilled in the art, for example, by formation of diastereoisomeric salts which may be separated by crystallization, gas-liquid or liquid chromatography, selective reaction of one enantiomer with an enantiomer-specific reagent. It will be appreciated that where the desired enantiomer is converted into another chemical entity by a separation technique, then an additional step

15

is required to form the desired enantiomeric form.
Alternatively, specific enantiomers may be synthesized by
asymmetric synthesis using optically active reagents,
substrates, catalysts or solvents, or by converting one
5 enantiomer into the other by asymmetric transformation.

Certain compounds of the present invention may also
exist in different stable conformational forms which may
be separable. Torsional asymmetry due to restricted
10 rotation about an asymmetric single bond, for example
because of steric hindrance or ring strain, may permit
separation of different conformers. The present
invention includes each conformational isomer of these
compounds and mixtures thereof.

15

Certain compounds of the present invention may exist
in zwitterionic form and the present invention includes
each zwitterionic form of these compounds and mixtures
thereof.

20

The compounds of the present invention are useful in
the inhibition of HCV NS3 protease, as well as, the
prevention or treatment of infection by the hepatitis C
virus and the treatment of consequent pathological
25 conditions. The treatment involves administering to a
patient, in need of such treatment, a compound of the
present invention, or a pharmaceutically acceptable salt
thereof, or a pharmaceutical composition comprising a
pharmaceutical carrier and a therapeutically effective
30 amount of a compound of the present invention, or a
pharmaceutically acceptable salt thereof.

It will be appreciated by those skilled in the art that reference herein to treatment extends to prophylaxis as well as the treatment of established infections or symptoms. This includes initiating treatment pre- and
5 post-exposure to the virus. In addition, the present invention can be administered in conjunction with immunomodulators, such as α -, β -, or γ -interferons; other antiviral agents such as ribavirin, amantadine; other inhibitors of HCV NS3 protease; inhibitors of other
10 targets in the HCV life cycle, which include but not limited to, helicase, polymerase, metalloprotease, or internal ribosome entry site (IRES); or combinations thereof. The additional agents may be combined with the compounds of this invention to create a single dosage
15 form. Alternatively these additional agents may be separately administered to a mammal as part of a multiple dosage form.

These methods are useful in decreasing HCV NS3
20 protease activity in a mammal. These methods are useful for inhibiting viral replication in a mammal. If the pharmaceutical composition comprises only a compound of this invention as the active component, such methods may additionally comprise the step of administering to said
25 mammal an agent selected from an immunomodulatory agent, an antiviral agent, a HCV protease inhibitor, or an inhibitor of other targets in the HCV life cycle such as helicase, polymerase, or metalloprotease or IRES. Such additional agent may be administered to the mammal prior
30 to, concurrently with, or following the administration of the compositions of this invention.

The compounds of the present invention are also useful in the preparation and execution of screening or replication assays for antiviral compounds. Furthermore, the compounds of the present invention are useful in
5 establishing or determining the binding site of other
antiviral compounds to HCV NS3 protease, for example, by
competitive inhibition.

The compounds of the present invention may be
10 administered orally, parenterally (including subcutaneous
injections, intravenous, intramuscular, intrasternal
injection or infusion techniques), by inhalation spray or
rectally, in dosage unit formulations containing
conventional non-toxic pharmaceutically acceptable
15 carriers, adjuvants and vehicles.

This invention also provides a pharmaceutical
composition for use in the above-described therapeutic
method. A pharmaceutical composition of the present
20 invention comprises an effective amount of a compound of
Formula I in association with a pharmaceutically
acceptable carrier, excipient or diluent.

The active ingredient in such formulations comprises
25 from 0.1 percent to 99.9 percent by weight of the
formulation. By "pharmaceutically acceptable" it is
meant that the carrier, diluent or excipient must be
compatible with the other ingredients of the formulation
and not deleterious to the recipient thereof.

30

The present pharmaceutical compositions are prepared
by known procedures using well-known and readily
available ingredients. The compositions of this

invention may be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art. In making the

5 compositions of the present invention, the active ingredient will usually be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may

10 be a solid, semi-solid or liquid material which acts as a vehicle, excipient or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, beadlets, lozenges, sachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols, (as

15 a solid or in a liquid medium), soft and hard gelatin capsules, suppositories, sterile injectable solutions, sterile packaged powders and the like.

The compounds can be administered by a variety of

20 routes including oral, intranasally, rectal, transdermal, subcutaneous, intravenous, intramuscular and intranasal.

When administered orally, these compositions are prepared according to techniques well-known in the art of

25 pharmaceutical formulation. For oral administration, the compound is typically formulated with excipients such as binders, fillers, lubricants, extenders, diluents, disintegration agents and the like as are known in the art.

30

For parenteral administration, the compound is formulated in pharmaceutically acceptable non-toxic, parenterally-acceptable diluents or solvents, such as

mannitol, 1,3-butanediol, water, 5 percent dextrose, Ringer's solution or isotonic sodium chloride solution, or suitable dispersing or wetting and suspending agents, such as sterile, bland, fixed oils, including synthetic
5 mono- or diglycerides, and fatty acids, including oleic acid.


A compound of the present invention, or a salt or solvate thereof, can be formulated in unit dosage
10 formulations comprising a dose between about 0.1 mg and about 1000 mg, or more, according to the particular treatment involved. An example of a unit dosage formulation comprises 5 mg of a compound of the present invention in a 10 mL sterile glass ampoule. Another
15 example of a unit dosage formulation comprises about 10 mg of a compound of the present invention as a pharmaceutically acceptable salt in 20 mL of isotonic saline contained in a sterile ampoule.





20 The compounds of the present invention can also be administered to humans in a dosage range of 1 to 100 mg/kg body weight in divided doses. One preferred dosage range is 1 to 20 mg/kg body weight orally in divided doses. It will be understood, however, that the specific
25 dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the route of administration, the age, body
30 weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

General methods useful for the synthesis of compounds embodied in this invention are shown below. The preparations shown below are disclosed for the purpose of illustration and are not meant to be
5 interpreted as limiting the processes to make the compounds by any other methods.

It will be appreciated by those skilled in the art that a number of methods are available for the
10 preparation of the compounds of the present invention. These compounds may be prepared by processes which include processes known in the chemical art for the production of structurally analogous compounds or by a novel process described herein. A process for the
15 preparation of these compounds (or a pharmaceutically acceptable salt thereof) and novel intermediates for the manufacture of these compounds provide further features of the invention and are illustrated by the following procedures in which the meanings of the generic radicals
20 are as defined above, unless otherwise specified. It will be recognized that it may be preferred or necessary to prepare such a compound in which a functional group is protected using a conventional protecting group then to remove the protecting group to provide a compound of the
25 present invention.

For example, compounds of the present invention, having the structure of Formula I, II or III, such as those compounds of Formula IIIA and IIIB, shown below,
30 were synthesized, as shown in the following scheme, indirectly from compounds of formula IV, and directly from compounds of formula VIIA or VIIB.






The α -carboxylic acid of a compound of Formula VIIA or VIIB is coupled with $R_1SO_2NH_2$, which was prepared by treatment of R_1SO_2Cl in ammonia saturated tetrahydrofuran solution, in the presence of peptide coupling agent, such as CDI or EDAC, and in the presence of a base, such as 4-dimethylaminopyridine (4-DMAP) and/or 1,8-diazabicyclo [5.4.0]undec-7-ene (DBU), to form a compound of Formula I, II or III.

Compounds having the structure of Formula IV, VI, VIIA or VIIB can be prepared as described herein, and in

non-provisional application U.S.S.N. 10/001,850 filed November 20, 2001; International Application Number PCT/CA00/00353, Publication No. WO 00/59929, published October 12, 2000 titled "Macrocyclic Peptides Active
5 Against the Hepatitis C Virus"; and U.S. Patent 6,323,180 granted November 27, 2001 (corresponding to International Application Number PCT/CA99/00736, Publication No. WO 00/09543), titled "Hepatitis C Inhibitor TriPeptides". The disclosure of U.S. Patent 6,323,180 is incorporated,
10 in its entirety, herein by reference.


Compounds having the structure of Formula IV can also be prepared by coupling chemical precursor A with chemical precursor B as shown below.

15



Chemical precursor A, which is shown above, may be synthesized as described in the following.

21




In this scheme, the ethyl ester (A1) is condensed with the appropriate aniline (A2), in the presence of
5 acid, to form an imine. The imine is then cyclized by heating at about 260-280°C to give the corresponding 4-hydroxyquinoline (A3). The 4-hydroxyquinoline (A3) is subsequently refluxed with phosphorous oxychloride to form a 4-chloroquinoline (A4).

10

Commercially available Boc-4(R)-hydroxyproline (A5), in a suitable solvent such as DMSO, is then treated with a base, for example, sodium hydride or potassium tert-butoxide, to form an alkoxide of A5 which is then reacted
15 with the 4-chloroquinoline (A4) to give chemical precursor A.

Alternate means of forming the chemical precursor A, and of forming fragments or precursors of chemical precursor (A), are described in International Application Number PCT/CA00/00353, Publication No. WO 00/59929, 5 titled "Macrocyclic Peptides Active Against the Hepatitis C Virus", and U.S. Patent 6,323,180 granted November 27, 2001 (corresponding to International Application Number PCT/CA99/00736, Publication No. WO 00/09543), titled "Hepatitis C Inhibitor Tri-Peptides". The disclosure of 10 U.S. Patent 6,323,180 is incorporated, in its entirety, herein by reference.

Chemical precursor B, which is also shown above, may be synthesized as described in the following scheme with 15 slight modifications depicted in the experimental herein.



Treatment of commercially available or easily 20 obtainable imine (B1) with 1,4-dihalobutene (B2) in presence of a base produces, provides the resulting imine (B3). Acid hydrolysis of B3 then provides B, which has an allyl substituent syn to the carboxyl group. It is preferred that for compounds B3 and B that the vinyl 25 group is syn to the ester.

Exemplification

The specific examples that follow illustrate the syntheses of the compounds of the instant invention, and are not to be construed as limiting the invention in sphere or scope. The methods may be adapted to variations in order to produce compounds embraced by this invention but not specifically disclosed. Further, variations of the methods to produce the same compounds in somewhat different manner will also be evident to one skilled in the art.

Chemical abbreviations commonly used to identify chemical compounds in the literature include Bn: benzyl; Boc: tert-butyloxycarbonyl {Me₃COC(O)}; BSA: bovine serum albumin; CDI: carbonyldiimidazole; DBU: 1,8-diazabicyclo[5.4.0]-undec-7-ene; CH₂Cl₂=DCM: methylene chloride; DEAD: diethylazodicarboxylate; DIAD: diisopropylazodicarboxylate; DIEA: diisopropylethylamine; DIPEA: diisopropylethylamine; 4-DMAP: 4-dimethylaminopyridine; DCC: 1,3-dicyclohexyl-carbodiimide; DMF: dimethylformamide; DMSO: dimethylsulfoxide; DPPA: diphenylphosphoryl azide; EDAC: ethyldimethylaminopropylcarbodiimide hydrochloride; EDTA: ethylenediaminetetraacetic acid; Et: ethyl; EtOH: ethanol; EtOAc: ethyl acetate; Et₂O: diethyl ether; Grubb's Catalyst: bis(tricyclohexylphosphine)benzylidene ruthenium (IV) dichloride; HATU: [O-7-azabenzotriazol-1-yl)-1, HBTU: [O-(1H-benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate; PYBROP: Bromotris-pyrrolidino-phosphonium hexafluorophosphate; HOAT, 1-hydroxy-7-azabenzotriazole; HPLC: high performance liquid chromatography; MS: mass spectrometry; Me: methyl;

MeOH: methanol; NMM: N-methylmorpholine; NMP: N-methylpyrrolidine; Pr: propyl; Succ: 3-carboxypropanoyl; PPA: polyphosphoric acid; TBAF: tetra-n-butylammonium fluoride; 1,2-DCE or DCE: 1,2-dichloroethane; TBTU: 2-(1H-5 benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate; TFA: trifluoroacetic acid; THF: tetrahydrofuran

Solution percentages express a weight to volume relationship, and solution ratios express a volume to volume relationship, unless stated otherwise. Nuclear magnetic resonance (NMR) spectra were recorded either on a Bruker 300, 400 or 500 MHz spectrometer; the chemical shifts (δ) are reported in parts per million. Flash chromatography was carried out on silica gel (SiO_2) evident to one skilled in the art. All Liquid Chromatography (LC) data were recorded on a Shimadzu LC-10AS liquid chromatograph using a SPD-10AV UV-Vis detector and Mass Spectrometry (MS) data were determined with a Micromass Platform for LC in electrospray mode (ES+).

Unless otherwise noted, each compound was analyzed, by LC/MS, using one of five methodologies, having the following conditions.


- Columns: (Method A) - YMC ODS S7 C18 3.0x50 mm
5 (Method B) - YMC ODS-A S7 C18 3.0x50 mm
(Method C) - YMC S7 C18 3.0x50 mm
(Method D) - YMC Xterra ODS S7 3.0x50 mm
(Method E) - YMC Xterra ODS S7 3.0x50 mm
(Method F) - YMC ODS-A S7 C18 3.0x50 mm
10 (Method G) - YMC C18 S5 4.6x50 mm]
(Method H) - Xterra S7 3.0x50 mm
(Method I) - Xterra S7 C18 3.0x50 mm
- Gradient: 100% Solvent A/0% Solvent B to
15 0% Solvent A/100% Solvent B
Gradient time: 2 min. (A, B, D, F, G, H, I); 8 min. (C, E)
Hold time: 1 min. (A, B, D, F, G, H, I); 2 min. (C, E)
Flow rate: 5 mL/min
20 Detector Wavelength: 220 nm
Solvent A: 10% MeOH / 90% H₂O / 0.1% TFA
Solvent B: 10% H₂O / 90% MeOH / 0.1% TFA.
- (Method J) - YMC Xterra ODS S7 3.0x50 mm
25 Gradient: 100% Solvent A/0% Solvent B to
0% Solvent A/100% Solvent B
Gradient time: 4 min (A)
Hold time: 1 min (A); 2 min
Flow rate: 4 mL/min
30 Detector Wavelength: 220 nm
Solvent A: 10% MeOH / 90% H₂O / 0.1% TFA
Solvent B: 10% H₂O / 90% MeOH / 0.1% TFA.

The compounds and chemical intermediates of the present invention, described in the following examples, were prepared according to the following methods. For the description of the examples, in a synthetic step such as 1a, the numeral refers to the example number and the letter refers to the relative sequential step in the sequence.

10

Example 1Step 1a: Preparation of 4-hydroxy-2-phenyl-7-methoxyquinoline

15



20

25


30

To a solution of *m*-anisidine (300g, 2.44 mole) and ethyl benzoylacetate (234.2g, 1.22mole) in toluene (2.0 L) was added HCl (4.0 N in dioxane, 12.2 mL, 48.8 mmole). The resulting solution was refluxed for 6.5 hr using a Dean-Stark apparatus (about 56 ml of aqueous solution was collected). The mixture was cooled to rt, partitioned multiple times with aqueous HCl (10%, 3x500mL), aqueous NaOH (1.0 N, 2x200mL), water (3x200 mL), and the organic layer dried (MgSO₄) and concentrated in vacuo to supply an oily residue (329.5g). The crude product was heated in an oil bath (280 °C) for 80 min using a Dean-Stark apparatus (about 85 mL liquid was collected). The reaction mixture was cooled down to rt, the solid residue triturated with CH₂Cl₂ (400 mL), the resulting suspension

filtered, and the filter cake washed with more CH_2Cl_2 (2x150 mL). The resulting solid was dried in vacuo (50 °C; 1 torr; 1 day) affording analytically pure 4-hydroxy-7-methoxy-2-phenylquinoline as a light brown solid

5 (60.7g, 20% overall): ^1H NMR δ (DMSO) 3.86 (s, 3H), 6.26 (s, 1H), 6.94 (dd, $J=9.0$, 2.4 Hz, 1H), 7.21 (d, $J=2.4$ Hz, 1H), 7.56~7.61 (m, 3H), 7.80~7.84 (m, 2H), 8.00 (d, $J=9.0$ Hz, 1H), 11.54 (s, 1H); ^{13}C NMR (DMSO- d_6) δ 55.38, 99.69, 107.07, 113.18, 119.22, 126.52, 127.17, 128.97, 130.34,
10 134.17, 142.27, 149.53, 161.92, 176.48.


Step 1b: Preparation of 4-Chloro-7-methoxy-2-phenylquinoline



The 4-hydroxy-2-phenyl-7-methoxyquinoline (21.7g, 86.4mmole) was suspended in POCl_3 (240 mL). The suspension was refluxed for 2 hours. After removal of the POCl_3 in vacuo, the residue was partitioned between EtOAc (1L), and cold aqueous NaOH (generated from 1.0N 200 mL NaOH and 20 mL 10.0 N NaOH) and stirred for 15 min. The organic layer was washed with water (2x200mL), brine (200mL), dried (MgSO_4), and concentrated in vacuo to
25 supply 4-chloro-2-phenyl-7-methoxyquinoline (21.0g, 90%) as a light brown solid. ^1H NMR (DMSO- d_6) δ 3.97 (s, 3H), 7.36 (dd, $J=9.2$, 2.6 Hz, 1H), 7.49~7.59 (m, 4H), 8.08 (d, $J=9.2$ Hz, 1H), 8.19 (s, 1H), 8.26~8.30 (m, 2H); ^{13}C NMR (DMSO- d_6) δ 55.72, 108.00, 116.51, 119.52, 120.48,
30 124.74, 127.26, 128.81, 130.00, 137.58, 141.98, 150.20,


156.65, 161.30. LC-MS (Method D: retention time: 1.547),
MS m/z 270 ($M^+ + 1$).

5 Step 1c: Preparation of *N*-Boc (2*S*,4*R*)-(2-phenyl-7-methoxyquinoline-4-oxo)proline



To a suspension of *N*-Boc (2*S*,4*R*)-hydroxyproline (16.44g, 71.1mmol) (Fluka) in DMSO (250mL) was added *t*-
10 BuOK (19.93g, 177.6mmol) at 0 °C. The generated mixture was stirred for 1.5 hour and then 4-chloro-7-methoxy-2-phenylquinoline (21.02g, 77.9 mmol) added in three portions over 1 h. The reaction was stirred for one day, the reaction mixture was poured into cold water
15 (1.5L) and washed with Et₂O (4x200mL). The aqueous solution was acidified to pH 4.6, filtered to obtain a white solid, and dried in vacuo to supply the product, *N*-Boc (2*S*,4*R*)-(2-phenyl-7-methoxyquinoline-4-oxo)proline (32.5g, 98%): ¹H NMR (DMSO) δ 1.30, 1.32 (2s, 9H), 2.27-
20 2.31 (m, 1H), 2.58-2.69 (m, 1H), 3.73 (s, 2H), 3.88 (s, 3H), 4.30-4.35 (m, 1H), 5.72 (s, 1H), 7.11-7.17 (m, 1H), 7.35-7.57 (m, 5H), 7.92-7.97 (m, 1H), 8.24 (d, *J*=9.0 Hz, 2H), 12.76 (bs, 1H); LC-MS (Method D: retention time: 2.01; MS m/z 465 ($M^+ + H$)).


Step 1d: Preparation of racemic (1R,2S)/(1S,2R)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride (Method A and Method B) and chiral resolution of this racemate for the preparation of N-(1R, 2S)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride (Method C).



The named compound was made (Step d) racemic by each of the following methods A and B. This racemate could also be resolved to afford chiral Boc-(1R, 2S)-1-amino-2-vinylcyclopropyl carboxylic acid ester which was deprotected under acid conditions to afford (1R, 2S)-1-amino-2-vinylcyclopropane carboxylic acid ester hydrochloride (Method C).

Method A

A.1 Preparation of N-Benzyl Imine of Glycine Ethyl Ester



Glycine ethyl ester hydrochloride (303.8 g, 2.16 mole) was suspended in *tert*-butylmethyl ether (1.6 L). Benzaldehyde (231 g, 2.16 mole) and anhydrous sodium sulfate (154.6 g, 1.09 mole) were added and the mixture cooled to 0 °C using an ice-water bath. Triethylamine (455 mL, 3.26 mole) was added dropwise over 30 min and the mixture stirred for 48 h at rt. The reaction was


then quenched by addition of ice-cold water (1 L) and the organic layer was separated. The aqueous phase was extracted with *tert*-butylmethyl ether (0.5 L) and the combined organic phases washed with a mixture of

5 saturated aqueous NaHCO₃ (1 L) and brine (1 L). The solution was dried over MgSO₄, concentrated in vacuo to afford 392.4 g of the *N*-benzyl imine product as a thick yellow oil that was used directly in the next step. ¹H

10 NMR (CDCl₃, 300 MHz) δ 1.32 (t, *J*=7.1 Hz, 3H), 4.24 (q, *J*=7.1 Hz, 2H), 4.41 (d, *J*=1.1 Hz, 2H), 7.39–7.47 (m, 3H), 7.78–7.81 (m, 2H), 8.31 (s, 1H).

A.2 Preparation of racemic *N*-Boc-(1*R*,2*S*)/(1*S*,2*R*)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester

15




To a suspension of lithium *tert*-butoxide (84.06 g, 1.05 mol) in dry toluene (1.2 L), was added dropwise a

20 mixture of the *N*-benzyl imine of glycine ethyl ester (100.4 g, 0.526 mol) and *trans*-1,4-dibromo-2-butene (107.0 g, 0.500 mol) in dry toluene (0.6 L) over 60 min. After completion of the addition, the deep red mixture was quenched by addition of water (1 L) and *tert*-

25 butylmethyl ether (TBME, 1 L). The aqueous phase was separated and extracted a second time with TBME (1 L). The organic phases were combined, 1 N HCl (1 L) was added and the mixture stirred at room temperature for 2 h. The

organic phase was separated and extracted with water (0.8 L). The aqueous phases were then combined, saturated with salt (700 g), TBME (1 L) was added and the mixture cooled to 0 °C. The stirred mixture was then basified to pH 14 by the dropwise addition of 10 N NaOH, the organic layer separated, and the aqueous phase extracted with TBME (2 x 500 mL). The combined organic extracts were dried (MgSO₄) and concentrated to a volume of 1L. To this solution of free amine, was added BOC₂O or di-tert-butylidicarbonate (131.0 g, 0.6 mol) and the mixture stirred 4 days at rt. Additional di-tert-butylidicarbonate (50 g, 0.23 mol) was added to the reaction, the mixture refluxed for 3 h, and was then allowed cool to room temperature overnight. The reaction mixture was dried over MgSO₄ and concentrated in vacuo to afford 80 g of crude material. This residue was purified by flash chromatography (2.5 Kg of SiO₂, eluted with 1% to 2% MeOH/CH₂Cl₂) to afford 57 g (53%) of racemic *N*-Boc-(1*R*,2*S*)/(1*S*,2*R*)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester as a yellow oil which solidified while sitting in the refrigerator: ¹H NMR (CDCl₃, 300 MHz) δ 1.26 (t, *J*=7.1 Hz, 3H), 1.46 (s, 9H), 1.43-1.49 (m, 1H), 1.76-1.82 (br m, 1H), 2.14 (q, *J*=8.6 Hz, 1H), 4.18 (q, *J*=7.2 Hz, 2H), 5.12 (dd *J*=10.3, 1.7 Hz, 1H), 5.25 (br s, 1H), 5.29 (dd, *J*=17.6, 1.7 Hz, 1H), 5.77 (ddd, *J*=17.6, 10.3, 8.9 Hz, 1H); MS *m/z* 254.16 (M-1)

A.3 Preparation of Racemic (1*R*,2*S*)/(1*S*,2*R*) 1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride




N-Boc-(1*R*,2*S*)/(1*S*,2*R*)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester (9.39 g, 36.8 mmol) was dissolved in 4 N HCl/dioxane (90ml, 360 mmol) and was stirred for 2 h at rt. The reaction mixture was concentrated to supply (1*R*,2*S*)/(1*S*,2*R*)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride in quantitative yield (7 g, 100%). ¹H NMR (methanol-*d*₄) δ 1.32 (t, *J*=7.1, 3H), 1.72 (dd, *J*=10.2, 6.6 Hz, 1H), 1.81 (dd, *J*=8.3, 6.6 Hz, 1H), 2.38 (q, *J*=8.3 Hz, 1H), 4.26-4.34 (m, 2H), 5.24 (dd, 10.3, 1.3 Hz, 1H) 5.40 (d, *J*=17.2, 1H), 5.69-5.81 (m, 1H).

10

15


Method B



To a solution of potassium *tert*-butoxide (11.55 g, 102.9 mmol) in THF (450 mL) at -78 °C was added the commercially available *N,N*-dibenzyl imine of glycine ethyl ester (25.0 g, 93.53 mmol) in THF (112 mL). The reaction mixture was warmed to 0 °C, stirred for 40 min,

25

and was then cooled back to -78°C . To this solution was added *trans*-1,4-dibromo-2-butene (20.0 g, 93.50 mmol), the mixture stirred for 1 h at 0°C and was cooled back to -78°C . Potassium *tert*-butoxide (11.55 g, 102.9 mmol) was
5 added, the mixture immediately warmed to 0°C , and was stirred one more hour before concentrating in vacuo. The crude product was taken up in Et_2O (530 mL), 1N aq. HCl solution (106 mL, 106 mmol) added and the resulting biphasic mixture stirred for 3.5 h at rt. The layers
10 were separated and the aqueous layer was washed with Et_2O (2x) and basified with a saturated aq. NaHCO_3 solution. The desired amine was extracted with Et_2O (3x) and the combined organic extract was washed with brine, dried (MgSO_4), and concentrated in vacuo to obtain the free
15 amine. This material was treated with a 4N HCl solution in dioxane (100 mL, 400 mmol) and concentrated to afford (1*R*,2*S*)/(1*S*,2*R*)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride as a brown semisolid (5.3 g, 34% yield) identical to the material obtained from
20 procedure A, except for the presence of a small unidentified aromatic impurity (8%).

Method CC.1 Resolution of *N*-Boc-(1*R*,2*S*)/(1*S*,2*R*)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride

5

A solution of phosphate buffer (pH 8.0, 6.12 g, KH_2PO_4 , 121.41 g, Na_2PO_4 , in 9 L of water) was warmed to 40°C in a jacketed reactor with a mechanical stirrer. If needed, the pH was adjusted to 8.0 with the addition of phosphoric acid. A solution (1.8 kg) of protease enzyme from *Bacillus globigii* (Sigma) was added and the pH of the resulting mixture was adjusted to 8.0 by the addition of 50% NaOH. To this mixture was then added the racemic ester (180 g, 0.75 mmol) as a DMSO solution (1.8 L) over a period of 2 hours. A few ml of 50% NaOH was added in the first 4 hours to keep the pH at 8.0. Aliquots (1 ml) were taken from the reaction mixture at various times and analyzed by the following procedure: 1) Acidification to pH 4 by addition of 6(N) sulfuric acid. 2) The aliquot (1 ml) was extracted with MTBE (2 ml) and the MTBE extract divided into two portions and evaporated. 3) One portion was redissolved in acetonitrile and analyzed by HPLC to determine the conversion: HPLC Column: YMC Propack C-18, 150 X 4.6 mm, S-3 μm . 40°C Solvent:


Gradient 0.2% H₃PO₄-CH₃CN 70:30 to 10:90 in 15 minutes,
Re-equilibration back to original at 16 minutes and
continue till 20 min before next injection. Flow Rate: 1
ml/min, Detection: UV 210 nm. Elution Time: Acid 4.7

5 min, Ester 8 min

4) The second portion portion was redissolved in the
mobile phase and analyzed by HPLC to determine the
enantiomeric excess: Column: Chiralpak AD, Room
Temperature, Solvent: Heptane:Ethanol (Absolute, 200
10 proof):Trifluoroacetic acid (97.4:2.5:0.1)
Flow rate: 1 ml/min, Detection: UV 210 nm
Elution: Desired (1R,2S) enantiomer 9.2 min, Undesired
(1S,2R) enantiomer 9.7 min

15 Note: The reaction was terminated when the analysis
showed that the enantiomeric excess of the ester has
reached the desired level (usually ee>95% in 16-24 hrs).
After 22 hours, the reaction was cooled to room
temperature. The pH was 7.75. Analyses of an aliquot (as
20 above) showed acid to ester ratio of 55:45. Sodium
hydroxide (10%) was added to the reaction mixture to
bring the pH to 8.5. The aqueous layer was extracted
with MTBE (2 X 6 Liter) until no ester remained in the
aqueous layer. The MTBE extracts were washed with 5%
25 NaHCO₃ (4 Liter) to remove any acid remaining and then
with water (2 X 4 L). The solvent is evaporated to give
the ester 87.9g (97%, based on theoretical maximum yield
90g) as a yellow oil.

C.2 Preparation of chiral (1*R*,2*S*)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride




5

N-BOC (1*R*,2*S*)-1-amino-2-vinylcyclopropanecarboxylic acid ethyl ester (8.5 g, 33.3 mmol) was stirred under an N₂ atmosphere with 200 mL of 4N HCl/dioxane (Aldrich) at
 10 rt for 3 h. The solvent was removed under reduced pressure keeping the temperature below 40 C. This gave 6.57 g (~100%) of (1*R*,2*S*)-1-amino-2-vinylcyclopropanecarboxylic acid ethyl ester hydrochloride as a light tan solid. ¹H NMR (300 MHz,
 15 CD₃OD) δ 1.31 (t, *J*=7.0 Hz, 3 H), 1.69-1.82 (m, 2 H), 2.38 (q, *J*=8.8 Hz, 1 H), 4.29 (q, *J*=7.0 Hz, 2 H), 5.22 (d, *J*=10.3 Hz, 1 H), 5.40 (d, *J*=17.2 Hz, 1 H), 5.69-5.81 (m, 1 H). LC-MS (Method J, retention time: 0.58 min), MS *m/z* 156 (*M*⁺+1).

20

Step 1e: Preparation of 2(*S*)-(1(*R*)-ethoxycarbonyl-2(*S*)-vinylcyclopropylcarbamoyl)-4(*R*)-(7-methoxy-2-phenylquinolin-4-yloxy)pyrrolidine-1-carboxylic acid *tert*-butyl ester (High R_f Diastereomer) and 2(*S*)-(1(*S*)-ethoxycarbonyl-2(*R*)-vinylcyclopropylcarbamoyl)-4(*R*)-(7-methoxy-2-phenylquinolin-4-yloxy)-pyrrolidine-1-
 25 carboxylic acid *tert*-butyl ester (Low R_f Diastereomer)



Method A: To a solution of Boc-(2S,4R)-(2-phenyl-7-methoxyquinoline-4-oxo)proline (11.0 g, 23.7 mmole),
 5 (1R,2S)/(1S,2R)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride where carboxy group is syn to vinyl moiety (5.40 g, 28.2 mmole), NMM (20.8 mL; 18.9 mmole) in 500 mL of 50% CH₂Cl₂/THF was added the coupling reagent PyBrop or Bromotrispyrrolidinophosphonium
 10 hexafluorophosphate (16.0 g, 34.3 mmole) in three portions in 10 min at 0 °C. The solution was stirred at rt for one day and then was washed with pH 4.0 buffer (4x50 mL). The organic layer was washed with saturated aqueous NaHCO₃ (100 mL), the aqueous washing extracted
 15 with ethyl acetate (150 mL), and the organic layer

backwashed with pH 4.0 buffer (50 mL), and saturated aqueous NaHCO₃ (50 mL). The organic solution was dried (MgSO₄), concentrated and purified using a Biotage 65M column (eluted with 50% EtOAc/Hexanes) to provide over
5 7.5 g of a 1:1 mixture of (1R,2S) and (1S,2R) cyclopropane diastereomers (50% overall) or alternatively elution using a slow to 15% to 60% EtOAc in hexanes gradient to supply 3.54 g, 25% overall of the high R_f eluted (1R,2S) cyclopropane diastereomer and 3.54 g, 25%
10 overall of the low R_f eluted (1S,2R) cyclopropane diastereomer.

Method B: To a solution of (1R,2S)/(1S,2R)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester
15 hydrochloride (7.5 g, 39.1 mmol) was combined with diisopropylethylamine (32.5 mL, 186 mmol) in dichloromethane (150 mL). To the resulting mixture was added HOBt hydrate (6.85 g, 44.7 mmol) and *N*-Boc (2S,4R)-(2-phenyl-7-methoxyquinoline-4-oxo)proline (17.3 g, 37.3
20 mmol) followed by addition of HBTU (16.96 g, 44.7 mmol). A slight exotherm occurred immediately, and the mixture was stirred at room temperature overnight. The mixture was then concentrated *in vacuo* and redissolved in ethyl acetate (600 mL). The solution was washed with water (2
25 x 200 mL), then with 10% aqueous sodium bicarbonate (2 x 200 mL), then with water (150 mL) and finally with brine (150 mL). The organic was dried over anhydrous magnesium sulfate and filtered, and the filtrate was concentrated
30 *in vacuo* to a beige glassy solid. Purification was performed in multiple batches (7 g each) by flash chromatography on a Biotage Flash 75M cartridge (66% hexanes/ethyl acetate) to provide the high R_f eluted (1R,2S) cyclopropane diastereomer (9.86 g total, 44.0%

yield), followed by the low Rf eluted cyclopropane diastereomer (10.43 g total, 46.5% yield). A total of 1.97 g of mixed fractions were recovered to give an overall conversion of 99.3% to the two diastereomers.

5

Data for 2(S)-(1(R)-ethoxycarbonyl-2(S)-vinyl-cyclopropylcarbamoyl)-4(R)-(7-methoxy-2-phenylquinolin-4-yloxy)pyrrolidine-1-carboxylic acid tert-butyl ester


(High Rf Cyclopropane Diastereomer): ^1H NMR (CDCl_3) δ 1.22
10 (t, $J=7.1$ Hz, 3H), 1.44 (s, 9H), 1.48-1.58 (m, 1H), 1.89 (bs, 1H), 2.05-2.20 (m, 1H), 2.4 (bs, 1H), 2.90 (bs, 1H), 3.80, 3.87 (2bs, 2H), 3.94 (s, 3H), 4.07-4.22 (m, 2H), 4.56 (bs, 1H), 5.13 (d, $J=10.2$ Hz, 1H), 5.30 (d, $J=17.1$ Hz, 1H), 5.70-5.82 (m, 1H), 7.03 (bs, 1H), 7.09 (dd,
15 $J=9.1$, 2 Hz, 1H), 7.42-7.53 (m, 4H), 7.61 (bs, 1H), 7.96 (d, $J=9.1$ Hz, 1H), 8.05 (d, $J=7.2$ Hz, 2H); ^1H NMR (methanol- d_4) δ 1.23 (t, $J = 7.2$ Hz, 3H), 1.4 (s, 4H), 1.45 (s, 6H), 1.73 (dd, $J = 7.9$, 1.5 Hz, 0.4H), 1.79 (dd, $J = 7.8$, 2.4 Hz, 0.6H), 2.21 (q, $J = 8.2$ Hz, 1H), 2.44-
20 2.49 (m, 1H), 2.66-2.72 (m, 0.4H), 2.73-2.78 (m, 0.6H), 3.93-3.95 (m, 2H), 3.96 (s, 3H), 4.10-4.17 (m, 2H), 4.44 (q, $J = 7.8$ Hz, 1H), 5.13 (d, $J = 10.7$ Hz, 1H), 5.31 (d, $J = 17.7$ Hz, 0.4H), 5.32 (d, $J = 17.4$ Hz, 0.6H), 5.49 (bs, 1H), 5.66-5.82 (m, 1H), 7.16 (dd, $J = 9.2$, 2.5 Hz, 1H),
25 7.26 (s, 1H), 7.42 (d, $J = 2.4$ Hz, 1H), 7.48-7.55 (m, 3H), 8.02-8.05 (m, 3H); ^{13}C NMR (CDCl_3) δ 14.22; 22.83, 28.25, 33.14, 33.58, 39.92, 51.84, 55.47, 58.32, 61.30, 75.86, 81.27, 98.14, 107.42, 115.00, 117.84, 118.27, 122.63, 123.03, 127.50, 128.72, 129.26, 133.39, 140.06,
30 151.23, 159.16, 160.34, 161.35, 169.78, 171.68. LC-MS (Method D: retention time: 1.62), MS m/z 602 ($M^+ + 1$).

Data for 2(*S*)-(1(*S*)-ethoxycarbonyl-2(*R*)-vinylcyclopropylcarbamoyl)-4(*R*)-(7-methoxy-2-phenylquinolin-4-yloxy)pyrrolidine-1-carboxylic acid *tert*-butyl ester (Low Rf Cyclopropane Diastereomer): ^1H NMR (CDCl_3)

5 δ 1.25 (t, $J=7.2$ Hz, 3 H), 1.45 (s, 9 H), 1.35-1.52 (m, 1 H), 1.85 (bs, 1 H), 2.13-2.22 (m, 1 H), 2.4 (bs, 1 H), 2.95 (bs, 1 H), 3.83 (bs, 2 H), 3.99 (s, 3 H), 4.13 (q, $J=7.2$ Hz, 2 H), 4.59 (s, 1 H), 5.16 (d, $J=10.8$ Hz, 1 H), 5.33 (d, $J=17.2$ Hz, 1 H), 5.40 (bs, 1 H), 5.74-5.86 (m, 1 H), 7.06 (s, 1 H), 7.14 (dd, $J=9.2, 2.3$ Hz, 1 H), 7.46-7.6 (m, 3 H), 8.00 (d, $J=9.2$, 1 H), 8.08, 8.10 (2s, 2H).

LC-MS (Method D: retention time: 1.66), MS m/z 602 (M^++1).

15 Step 1f: Preparation of 1-([4-(7-Methoxy-2-phenylquinolin-4(*R*)-yloxy)pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino}-2(*S*)-vinylcyclopropanecarboxylic acid ethyl ester, dihydrochloride




20

5.88g (9.77mmol) of 2(*S*)-(1(*R*)-ethoxycarbonyl-2(*S*)-vinylcyclopropylcarbamoyl)-4(*R*)-(7-methoxy-2-phenylquinolin-4-yloxy)pyrrolidine-1-carboxylic acid *tert*-butyl ester (High Rf Cyclopropane Diastereomer) was dissolved in HCl/dioxane (4.0M; 200ml) and was stirred for 2.5 h at rt. The reaction mixture was concentrated to supply the titled compound which was directly used in the

next step. ^1H NMR (methanol- d_4) δ 1.25 (t, $J=7.1$ Hz, 3H),
1.51 (dd, $J=9.6, 5.4$ Hz, 1H), 1.78 (dd, $J=8.1, 5.5$ Hz,
1H), 2.30 (dd, $J=17.7, 8.7$ Hz, 1H), 2.27-2.67 (m, 1H),
3.06 (dd, $J=14.6, 7.4$ Hz, 1H), 3.56-3.76 (m, 3H), 3.99
5 (s, 2H), 4.07 (s, 3H), 4.16 (q, 2H), 4.77 (dd, $J=10.4,$
7.4 Hz, 1H), 5.13 (dd, $J=10.3, 1.7$ Hz), 5.33 (dd, $J=17.1,$
1.7 Hz), 5.71-5.81 (m, 1H), 6.06 (bs, 1H), 7.49 (dd,
 $J=9.4, 2.3$ Hz, 1H), 7.66-7.77 (m, 4H), 8.12-8.16 (m, 2H),
8.55 (d, $J=9.4$ Hz, 1H). ^{13}C NMR (methanol- d_4) δ 14.77,
10 23.23, 34.86, 37.25, 41.19, 43.90, 52.66, 60.35, 62.32,
62.83, 68.27, 72.58, 73.70, 81.21, 100.70, 102.44,
116.13, 118.67, 122.25, 126.93, 130.27, 130.94, 133.19,
134.14, 134.89, 143.79, 158.39, 166.84, 167.44, 169.57,
171.33. LC-MS (Method D: retention time: 1.55), MS m/z
15 502 (M^++1)

Examples 1,2


Preparation of Compounds 1 and 2



Compound 1 - R is Boc
Compound 2 - R is H

Step 1g: Preparation of 1-[[1-(2(S)-tert-butoxycarbonyl-amino-hex-5-enoyl)-4-(7-methoxy-2-phenyl-quinolin-4(R)-yloxy)pyrrolidine-2(S)-carbonyl]-1(R)-amino]-2(S)-vinyl-cyclopropanecarboxylic acid ethyl ester

5




To suspension of (0.600g; 1.04mmol) of 1-[[4-(7-Methoxy-2-phenylquinolin-4(R)-yloxy)pyrrolidine-2(S)-carbonyl]-1(R)-amino]-2(S)-vinylcyclopropanecarboxylic acid ethyl ester, dihydrochloride, 2(S)-tert-butoxycarbonyl-amino-5-hexenoic acid purchased from RSP Amino Acids (0.34g, 1.46mmol), NMM (0.57ml, 5.22mmol) in DMF (5mL) was added HATU (0.56g, 1.46 mmol) at 0 °C.

After being stirred for 2 days, the reaction mixture was diluted with EtOAc (200 mL), washed with pH 4.0 buffer (2x30 mL), saturated aqueous NaHCO₃ (30 mL), brine (30mL), dried (MgSO₄), purified by a Biotage 40 M column (eluted with 30% to 60% EtOAc in Hexanes) to supply the titled compound, 1-[[1-(2(S)-tert-butoxycarbonylamino-hex-5-enoyl)-4-(7-methoxy-2-phenyl-quinolin-4(R)-yloxy)-pyrrolidine-2(S)-carbonyl]-1(R)-amino]-2(S)-vinyl-cyclopropanecarboxylic acid ethyl ester, as a yellow oil (0.56g, 76%): ¹H NMR (300 MHz, CDCl₃) δ 1.20 (t, J=7 Hz, 3H), 1.39 (s, 9H), 1.43-1.48 (m, 1H), 1.64-1.92 (m, 4H), 2.10-2.20 (m, 2H), 2.36-2.44 (m, 1H), 2.91-2.99 (m, 1H),

3.94 (s, 3H), 4.04-4.22 (m, 2H), 4.29 (d, $J=11$ Hz, 1H), 4.46-4.51 (m, 1H), 4.81 (dd, $J=8.1, 5.5$ Hz, 1H), 4.98-5.15 (m, 4H), 5.29 (dd, $J=17.2, 1.5$ Hz, 1H), 5.41 (brs, 1H), 5.67-5.86 (m, 2H), 7.04 (s, 1H), 7.08 (dd, $J=9.2, 2.6$ Hz, 1H), 7.42-7.64 (m, 4H), 7.97-8.06 (m, 3H). LC-MS (Method E-retention time: 1.59), MS m/z 713 (M^++1).

Step 1h: Preparation of (1*S*,4*R*,6*S*,11*S*)-7-*trans*-11-*tert*-butoxycarbonylamino-15-(7-methoxy-2-phenylquinolin-4-yloxy)-2,12-dioxo-3,13-diazatricyclo[11.3.0.0^{4,6}]hexadec-7-ene-4-carboxylic acid ethyl ester




To a solution of 1-[[1-(2(*S*)-*tert*-butoxycarbonylamino-hex-5-enoyl)-4-(7-methoxy-2-phenylquinolin-4(*R*)-yloxy)-pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino]-2(*S*)-vinyl-cyclopropanecarboxylic acid ethyl ester (360 mg, 0.505 mmol) in 0.25L of Argon degassed 1,2-DCE, was added 70 mg of Grubbs catalyst, the mixture degassed under Ar and heated to reflux for 3 h. The reaction mixture was cooled to rt, the mixture degassed once more, a final 70 mg portion of Grubbs catalyst added, and the mixture heated to reflux for 12 h. The dark brown solution was concentrated in vacuo, purified by a Biotage 40 M column (eluted with 30% to 65% EtOAc in Hexanes) to supply the intermediate (1*S*,4*R*,6*S*,11*S*, 15*R*)-7-*trans*-11-

tert-butoxycarbonylamino-15-(7-methoxy-2-phenylquinolin-4-yloxy)-2,12-dioxo-3,13-diazatricyclo-[11.3.0.0^{4,6}]hexadec-7-ene-4-carboxylic acid ethyl ester (165 mg, 46%) as a pale yellow oil: ¹H NMR (500 MHz, CD₃OD) δ 1.12 (s, 9H), 1.23 (t, J=7Hz, 3H), 1.41 (dd, J=7.5, 5.8 Hz, 1H), 1.64-1.71 (m, 2H), 2.04-2.18 (m, 2H), 2.32 (m, 2H), 2.60-2.70 (m, 2H), 3.95 (s, 3H), 4.11-4.23 (m, 4H), 4.34 (m, 1H), 4.64 (m, 1H), 5.25 (m, 1H), 5.54 (d, J=15.3 Hz, 1H), 5.65 (s, 1H), 7.18 (dd, J=9.2, 2.4 Hz, 1H), 7.29 (s, 1H), 7.40 (d, J=2.4 Hz, 1H), 7.56-7.59 (m, 3H), 8.02-8.04 (m, 2H), 8.09 (d, J=9.2 Hz, 1H). LC-MS (Method E-retention time: 1.66), MS m/z 685 (M⁺+1).

Step 1i: Preparation of (1S,4R,6S,11S,15R)-7-trans-11-tert-butoxycarbonylamino-15-(7-methoxy-2-phenyl-quinolin-4-yloxy)-2,12-dioxo-3,13-diazatricyclo[11.3.0.0^{4,6}]hexadec-7-ene-4-carboxylic acid


To a solution of (1S,4R,6S,11S,15R)-7-trans-11-tert-butoxycarbonylamino-15-(7-methoxy-2-phenylquinolin-4-yloxy)-2,12-dioxo-3,13-diazatricyclo[11.3.0.0^{4,6}]hexadec-7-ene-4-carboxylic acid ethyl ester (80 mg, 0.12 mmol) in 7 mL of 4:2:1 THF/H₂O/MeOH was added 28 mg of LiOH (0.70 mmol), the mixture stirred 24 h, the solution adjusted to pH 7, and the mixture concentrated until only the water layer remained. The solution was adjusted to pH 4, the acidic aqueous solution extracted with EtOAc (3 x 100 mL portions) and the organic extracts combined, dried (MgSO₄) and concentrated to afford (1S,4R,6S,11S,15R)-7-trans-11-tert-butoxycarbonyl-amino-15-(7-methoxy-2-phenylquinolin-4-yloxy)-2,12-dioxo-3,13-diazatricyclo[11.3.0.0^{4,6}]hexadec-7-ene-4-carboxylic acid (62.2 mg, 80%): ¹H NMR (300 MHz, CD₃OD) δ 1.12 (s, 9H),

1.18-2.48 (m, 8H), 2.65-2.69 (m, 1H), 3.94 (s, 3H), 4.18-4.34 (m, 3H), 4.83 (t, $J=8$ Hz, 1H), 5.30-5.38 (m, 2H), 5.53-5.61 (m, 2H), 7.14 (dd, $J=9.2, 2.4$ Hz, 1H), 7.24 (s, 1H), 7.39 (d, $J=2.4$ Hz, 1H), 7.48-7.64 (m, 3H), 8.02-8.07 (m, 3H). LC-MS (Method E-retention time: 1.45), MS m/z 657 (M^++1). HRMS m/z ($M+H$)⁺ calcd for $C_{36}H_{41}N_4O_8$: 657.2924, found 657.2924.



Preparation of Cyclopropylsulfonamide from Cyclopropyl sulfonyl chloride.

To a solution of 100 mL of THF cooled to 0 °C was bubbled in gaseous ammonia until saturation was reached. To this solution was added a solution of 5 g (28.45 mmol) of cyclopropylsulfonyl chloride (purchased from Array Biopharma) in 50 mL of THF, the solution warmed to rt overnight and stirred one additional day. The mixture was concentrated until 1-2 mL of solvent remained, applied on to 30 g plug of SiO_2 (eluted with 30% to 60% EtOAc/Hexanes) to afford 3.45g (100%) of cyclopropyl sulfonamide as a white solid. 1H NMR (Methanol- d_4) δ 0.94-1.07 (m, 4H), 2.52-2.60 (m, 1H); ^{13}C NMR (methanol- d_4) δ 5.92, 33.01.




Preparation of cyclobutylsulfonamide from
cyclobutylbromide

To a solution of 5.0 g (37.0 mmol) of cyclobutyl
5 bromide in 30 mL of anhydrous diethyl ether (Et₂O) cooled
to -78 °C was added 44 mL (74.8 mmol) of 1.7M *tert*-butyl
lithium in pentanes and the solution slowly warmed to -35
°C over 1.5 h. This mixture was cannulated slowly into a
solution of 5.0 g (37.0 mmol) freshly distilled sulfonyl
10 chloride in 100 mL of hexanes cooled to -40 °C, warmed to
0 °C over 1 h and carefully concentrated in vacuo. This
mixture was redissolved in Et₂O, washed once with some
ice-cold water, dried (MgSO₄) and concentrated carefully.
This mixture was redissolved in 20 mL of THF, added
15 dropwise to 500 mL of saturated NH₃ in THF and was
allowed to stir overnight. The mixture was concentrated
in vacuo to a crude yellow solid and was recrystallized
from the minimum amount of CH₂Cl₂ in hexanes with 1-2
drops of MeOH to afford 1.90 g (38%) of
20 cyclobutylsulfonamide as a white solid. ¹H NMR (CDCl₃) δ
1.95-2.06 (m, 2H), 2.30-2.54 (m, 4H), 3.86 (p, J=8 Hz,
1H), 4.75 (brs, 2H); ¹³C NMR (CDCl₃) δ 16.43, 23.93,
56.29. HRMS m/z (M-H)⁻ calcd for C₄H₈NSO₂: 134.0276, found
134.0282.

Step 1j: Preparation of Compound 1, Example 1

(1S,4R,6S,11S,15R)-7-trans-[4-Cyclopropanesulfonyl-amino-
carbonyl-15-(7-methoxy-2-phenylquinolin-4-yloxy)-2,12-
dioxo-3,13-diazatricyclo[11.3.0.0^{4,6}]hexa-dec-7-en-11-
yl]carbamic acid tert-butyl ester




To a solution of (1S,4R,6S,11S,15R)-7-trans-11-tert-
 10 Butoxycarbonylamino-15-(7-methoxy-2-phenylquin-olin-4-
 yloxy)-2,12-dioxo-3,13-diazatricyclo[11.3.0.0^{4,6}]hexadec-
 7-ene-4-carboxylic acid (47.7 mg, 0.0726 mmol) in THF (2
 mL) was added was added CDI (16.5mg, 0.102 mmol) in one
 portion under argon. The resulting solution was stirred
 15 for 30 min, refluxed for 30 min and allowed to cool down
 to rt. Cyclopropylsulfonamide (12.3 mg, 0.102 mmol) was
 added in one portion before the addition of a neat
 solution of DBU (0.016 mL, 0.102 mmol). The reaction was
 stirred for 24 h, diluted with EtOAc (100mL) and washed
 20 pH 4.0 buffer (3x20mL), brine (20mL), dried (MgSO₄) and
 purified using one preparative TLC plate (Analtech 20x40
 cm 1000 uM thickness) to supply (1S,4R,6S,11S,15R)-7-
 trans-[4-cyclopropanesulfonyl-aminocarbonyl-15-(7-
 methoxy-2-phenylquinolin-4-yloxy)-2,12-dioxo-3,13-
 25 diazatricyclo[11.3.0.0^{4,6}]hexadec-7-en-11-yl]carbamic acid

tert-butyl ester, Compound 1, Example 1, as a foam (31.8 mg, 58%): ^1H NMR (300 MHz, CD_3OD) δ 0.86-1.43 (m, 5H), 1.15 (s, 9H), 1.63-1.73 (m, 2H), 2.03-2.20 (m, 2H), 2.27-2.41 (m, 2H), 2.56-2.80 (m, 3H), 3.92 (s, 3H), 4.16-4.37 (m, 3H), 4.83 (t, $J=8$ Hz, 1H), 5.28-5.36 (m, 1H), 5.52-5.56 (m, 2H), 7.09 (dd, $J=9.2$, 2.2 Hz, 1H), 7.22 (s, 1H), 7.37 (d, $J=2.2$ Hz, 1H), 7.44-7.54 (m, 3H), 8.00-8.04 (m, 3H). LC-MS (Method D-retention time: 1.43), MS m/z 760 (M^++1). HRMS m/z ($M+H$) $^+$ calcd for $\text{C}_{39}\text{H}_{46}\text{SN}_5\text{O}_9$: 760.3016, found 760.3002.

Step 2k: Preparation of Compound 2, Example 2


(1S,4R,6S,11S,15R)-7-trans-Cyclopropanesulfonic acid [11-amino-15-(7-methoxy-2-phenylquinolin-4-yloxy)-2,12-dioxo-3,13-diazatricyclo-[11.3.0.0^{4,6}]hexadec-7-ene-4-carbonyl]amide dihydrochloride



A solution of 2 mL (8 mmol) of 4N HCl/dioxane was added to 11 mg (0.014 mmol) of (1S,4R,6S,11S,15R)-7-trans-[4-cyclopropanesulfonylaminocarbonyl-15-(7-methoxy-2-phenylquinolin-4-yloxy)-2,12-dioxo-3,13-diazatricyclo-[11.3.0.0^{4,6}]hexadec-7-en-11-yl]carbamic acid tert-butyl

ester, the mixture stirred 2h, concentrated and dried (50 °C; 20 torr; 12h) to afford 10.5 mg (~100%) of (1*S*,4*R*,6*S*,11*S*,15*R*)-7-*trans*-cyclopropanesulfonic acid [11-amino-15-(7-methoxy-2-phenylquinolin-4-yloxy)-2,12-dioxo-3,13-diazatricyclo-[11.3.0.0^{4,6}]hexadec-7-ene-4-carbonyl]amide dihydrochloride as a white solid: ¹H NMR (300 MHz, CD₃OD) δ 0.87-1.41 (m, 8H), 1.70 (m, 1H), 2.16 (s, 1H), 2.30 (m, 1H), 2.45 (m, 1H), 2.65 (m, 1H), 2.96 (m, 1H), 4.05 (s, 3H), 4.33 (s, 3H), 4.48 (m, 1H), 5.46-5.67 (m, 2H), 5.99 (s, 1H), 7.45 (m, 1H), 7.58-7.85 (m, 5H), 8.12 (brs, 2H), 8.35 (m, 1H). LC-MS (Method E- retention time: 1.09), MS *m/z* 660 (*M*⁺+1). HRMS *m/z* (*M*+H)⁺ calcd for C₃₄H₃₈SN₅O₇: 660.2492, found 660.2492.


15

Examples 3,4Preparation of Compounds 3 and 4

Compound 3 - R is Boc
Compound 4 - R is H
(Bis HCl Salt)

20

Step 3g: Preparation of 1-([1-(2(*S*)-tert-butoxycarbonyl-amino-hept-6-enoyl)-4-(7-methoxy-2-phenylquinolin-4(*R*)-yloxy)pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino]-2(*S*)-vinylcyclopropanecarboxylic acid ethyl ester




5

Following the procedure of Step 1g, except that (0.356g, 1.46 mmol) of 2(*S*)-tert-butoxycarbonylamino-6-heptenoic acid purchased from RSP Amino Acids, was coupled in place of 2(*S*)-tert-butoxycarbonylamino-5-hexenoic acid, to give the titled compound, 1-([1-(2(*S*)-tert-butoxycarbonylamino-hept-6-enoyl)-4-(7-methoxy-2-phenylquinolin-4(*R*)-yloxy)pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino]-2(*S*)-vinylcyclopropanecarboxylic acid ethyl ester as a yellow oil (0.600g, 79%): ¹H NMR (CDCl₃) δ (300 MHz, CDCl₃); 1.21 (t, *J*=7 Hz, 3H), 1.40 (s, 9H), 1.42-2.18 (m, 9H), 2.36-2.45 (m, 1H), 2.91-2.99 (m, 1H), 3.94 (s, 3H), 4.05-4.22 (m, 2H), 4.26-4.30 (m, 1H), 4.39-4.49 (m, 1H), 4.83 (dd, *J*=8.2, 5.6 Hz, 1H), 4.92-5.14 (m, 4H), 5.29 (dd, *J*=17.2, 1.4 Hz, 1H), 5.41 (brs, 1H), 5.63-5.80 (m, 2H), 7.04 (s, 1H), 7.09 (dd, *J*=9, 2 Hz, 1H), 7.42-7.54 (m, 4H), 8.00-8.05 (m, 3H). LC-MS (Method E-retention time: 1.53), MS *m/z* 727 (*M*⁺+1).

10
15
20

51




Step 3h: Preparation of (1*S*,4*R*,6*S*,12*S*,16*R*)-7-*trans*-12-
 tert-butoxycarbonylamino-16-(7-methoxy-2-phenylquinolin-4-
 5 4-yloxy)-2,13-dioxo-3,14-diazatricyclo[12.3.0.0^{4,6}]-
 heptadec-7-ene-4-carboxylic acid ethyl ester

Following the experimental and purification
 procedure of Step 1h, 1-[[1-(2(*S*)-tert-
 10 butoxycarbonylamino-hept-6-enoyl)-4-(7-methoxy-2-
 phenylquinolin-4(*R*)-yloxy)-pyrrolidine-2(*S*)-carbonyl]-
 1(*R*)-amino]-2(*S*)-vinyl-cyclopropanecarboxylic acid ethyl
 ester (404 mg, 0.555 mmol) was reacted with two 75 mg
 portions of Grubbs catalyst to afford (1*S*,4*R*,6*S*,12*S*,16*R*)-
 15 7-*trans*-12-tert-butoxycarbonylamino-16-(7-methoxy-2-
 phenylquinolin-4-yloxy)-2,13-dioxo-3,14-
 diazatricyclo[12.3.0.0^{4,6}]hepta-dec-7-ene-4-carboxylic
 acid ethyl ester (190 mg, 49%): ¹H NMR (500 MHz, CD₃OD) δ ,
 1.22 (t, *J*=7 Hz, 3H), 1.26 (s, 9H), 1.32-1.43 (m, 3H),
 20 1.59-1.64 (m, 1H), 1.77-1.90 (m, 3H), 2.11 (brs, 1H),
 2.29-2.34 (m, 1H), 2.60-2.65 (m, 1H), 2.70-2.74 (m, 1H),
 3.93 (s, 3H), 4.07-4.18 (m, 3H), 4.30 (d, *J*=7.6 Hz, 1H),
 4.52 (d, *J*=11.3 Hz, 1H), 4.67 (t, *J*=8.2 Hz, 1H), 5.35
 (dd, *J*=15.7, 6.0 Hz, 1H), 5.50-5.56 (m, 1H), 5.60 (brs,

1H), 7.10 (dd, $J=9.2$, 2 Hz, 1H), 7.26 (s, 1H), 7.37 (d, $J=2$ Hz, 1H), 7.53-7.57 (m, 3H), 8.00, 8.01 (2s, 2H), 8.17 (d, $J=9.2$ Hz, 1H). LC-MS (Method E-retention time: 1.58), MS m/z 699 (M^++1).

5

Step 3i: Preparation of (1*S*,4*R*,6*S*,12*S*,16*R*)-7-*trans*-12-*tert*-butoxycarbonylamino-16-(7-methoxy-2-phenylquinolin-4-yloxy)-2,13-dioxo-3,14-diazatricyclo-[12.3.0.0^{4,6}]heptadec-7-ene-4-carboxylic acid




10

Following the experimental and purification procedure of Step 1i, (1*S*,4*R*,6*S*,12*S*,16*R*)-7-*trans*-12-*tert*-butoxycarbonyl-amino-16-(7-methoxy-2-phenylquinolin-4-yloxy)-2,13-dioxo-3,14-diazatricyclo-[12.3.0.0^{4,6}]heptadec-7-ene-4-carboxylic acid ethyl ester (185 mg, 0.265 mmol) was reacted with 64 mg (1.6 mmol) of LiOH in 15.5 mL of 9:5:1.5 of THF/H₂O/MeOH to afford (1*S*,4*R*,6*S*,12*S*,16*R*)-7-*trans*-12-*tert*-butoxycarbonylamino-16-(7-methoxy-2-phenylquinolin-4-yloxy)-2,13-dioxo-3,14-diazatricyclo[12.3.0.0^{4,6}]heptadec-7-ene-4-carboxylic acid (180mg, 100%): ¹H NMR (300 MHz, CD₃OD) δ 1.24 (s, 9H), 1.33-1.43 (m, 1H), 1.61-1.97 (m, 6H), 2.10 (brs, 1H), 2.28-2.36 (m, 1H), 2.70 (m, 2H), 3.94 (s, 3H), 4.17 (d,

$J=10.6$ Hz, 1H), 4.28 (d, $J=4.8$ Hz, 1H), 4.65-4.70 (m, 1H), 5.34 (dd, $J=15.7$, 5.9 Hz, 2H), 5.53-5.62 (m, 2H), 7.14 (d, $J=9$ Hz, 1H), 7.32 (s, 1H), 7.39 (s, 1H), 7.56 (m, 3H), 8.01-8.02 (m, 2H), 8.22 (d, $J=9.2$ Hz, 1H). LC-MS
 5 (Method E-retention time: 1.49), MS m/z 671 (M^++1). HRMS m/z ($M+H$)⁺ calcd for $C_{37}H_{43}N_4O_8$: 671.3081, found 671.3080.

Step 3j: Preparation of Compound 3, Example 3

(1*S*,4*R*,6*S*,12*S*,16*R*)-7-*trans*-[4-Cyclopropanesulfonyl-
 10 aminocarbonyl-16-(7-methoxy-2-phenylquinolin-4-yloxy)-
2,13-dioxo-3,14-diazatricyclo[12.3.0.0^{4,6}]heptadec-7-en-
12-yl]carbamic acid *tert*-butyl ester



Compound 3, Example 3


15

Applying the procedural and purification practices of Step 1j, (144.9 mg, 0.216 mmol) of (1*S*,4*R*,6*S*,12*S*,16*R*)-7-*trans*-12-*tert*-butoxycarbonyl-amino-16-(7-methoxy-2-phenylquinolin-4-yloxy)-2,13-dioxo-3,14-
 20 diazatricyclo[12.3.0.0^{4,6}]heptadec-7-ene-4-carboxylic acid was converted to 92 mg (56%) of the product (1*S*,4*R*,6*S*,12*S*,16*R*)-7-*trans*-[4-cyclopropane-sulfonylaminocarbonyl-16-(7-methoxy-2-phenylquinolin-4-

yloxy)-2,13-dioxo-3,14-diazatricyclo[12.3.0.0^{4,6}]-
 heptadec-7-en-12-yl]carbamic acid *tert*-butyl ester: ¹H
 NMR (300 MHz, CD₃OD) δ 0.81-0.88 (m, 2H), 1.00-1.05 (m,
 2H), 1.15-1.19 (m, 1H), 1.32 (s, 9H), 1.42 (m, 1H), 1.59-
 5 1.68 (m, 1H), 1.77-1.93 (m, 4H), 2.08-2.26 (m, 2H), 2.60-
 2.73 (m, 3H), 3.94 (s, 3H), 4.21-4.34 (m, 2H), 4.43 (d,
 $J=11$ Hz, 1H), 4.56-4.62 (m, 1H), 5.28-5.36 (m, 1H), 5.46-
 5.55 (m, 2H), 7.07 (dd, $J=9.5$, 2.4 Hz, 1H), 7.27 (s, 1H),
 7.37 (d, $J=2.4$ Hz, 1H), 7.42-7.49 (m, 3H), 8.01, 8.03
 10 (2s, 2H), 8.18 (d, $J=9.5$ Hz, 1H). LC-MS (Method D-
 retention time: 1.48), MS m/z 774 ($M^+ + 1$). HRMS m/z ($M + H$)⁺
 calcd for C₄₀H₄₈SN₅O₉: 774.3173, found 774.3173.

Step 4k: Preparation of Compound 4, Example 4

15 (1*S*,4*R*,6*S*,12*S*,16*R*)-7-*trans*-Cyclopropanesulfonic acid [12-
 amino-16-(7-methoxy-2-phenylquinolin-4-yloxy)-2,13-dioxo-
 3,14-diazatricyclo [12.3.0.0^{4,6}]heptadec-7-ene-4-
 carbonyl]amide dihydrochloride



Compound 4, Example 4 - (Bis HCl Salt)


20 A solution of 5 mL (20 mmol) of 4N HCl/dioxane was
 added to 10.8 mg (0.014 mmol) of (1*S*,4*R*,6*S*,12*S*,16*R*)-7-
trans-[4-cyclopropanesulfonyl-aminocarbonyl-16-(7-

methoxy-2-phenyl-quinolin-4-yloxy)-2,13-dioxo-3,14-diazatricyclo[12.3.0.0^{4,6}]hexa-dec-7-en-12-yl]carbamic acid *tert*-butyl ester, the mixture stirred 2h, concentrated and dried (50 °C; 20 torr; 12h) to afford 10.6 mg (~100%) of (1*S*,4*R*,6*S*,12*S*,16*R*)-7-*trans*-cyclopropanesulfonic acid [12-amino-16-(7-methoxy-2-phenylquinolin-4-yloxy)-2,13-dioxo-3,14-diazatricyclo[12.3.0.0^{4,6}]heptadec-7-ene-4-carbonyl]amide dihydrochloride as a white solid: ¹H NMR (300 MHz, CD₃OD) δ 0.89 (m, 1H), 1.05-2.12 (m, 11H), 2.18-2.26 (m, 1H), 2.41-2.49 (m, 1H), 2.62-2.99 (m, 2H), 4.05 (s, 3H), 4.30 (m, 2H), 4.44-4.46 (m, 2H), 5.15-5.23 (m, 1H), 5.68-5.75 (m, 1H), 5.96 brs, 1H), 7.45 (d, *J*=9 Hz, 1H), 7.57 (s, 1H), 7.63-7.73 (m, 4H), 8.08-8.13 (m, 2H), 8.38 (d, *J*=9 Hz, 1H). LC-MS (Method E-retention time: 1.10), MS *m/z* 674 (*M*⁺+1). HRMS *m/z* (*M*+H)⁺ calcd for C₃₅H₄₀SN₅O₇: 674.2649, found 674.2655.

Examples 5,6


20

Preparation of Compounds 5 and 6



Compound 5, Example 5 - R is Boc
Compound 6, Example 6 - R is H (Bis HCl Salt)


Step 5g: Preparation of 1-[[1-(2(S)-tert-Butoxycarbonylamino-oct-7-enoyl)-4-(7-methoxy-2-phenyl-quinolin-4(R)-yloxy)pyrrolidine-2(S)-carbonyl]-1(R)-amino]-2(S)-vinylcyclopropanecarboxylic acid ethyl ester



5

Following the procedure of Step 1g, except that (0.35g, 1.4mmol) of 2(S)-tert-butoxycarbonylamino-7-octenoic acid purchased from RSP Amino Acids, was coupled in place of 2(S)-tert-butoxycarbonylamino-5-hexenoic acid, to give 1-[[1-(2(S)-tert-Butoxycarbonylamino-oct-7-enoyl)-4-(7-methoxy-2-phenyl-quinolin-4(R)-yloxy)pyrrolidine-2(S)-carbonyl]-1(R)-amino]-2(S)-vinylcyclopropanecarboxylic acid ethyl ester as a yellow oil (0.58g, 75%): ¹H NMR (300 MHz, CDCl₃) δ 1.20 (t, J=7 Hz, 3H), 1.32-1.47 (m, 5H), 1.39 (s, 9H), 1.54-1.68 (m, 2H), 1.68-2.05 (m, 3H), 2.09-2.18 (m, 1H), 2.35-2.48 (m, 1H), 2.89-2.99 (m, 1H), 3.93 (s, 3H), 4.02-4.22 (m, 2H), 4.27 (d, J=11.3 Hz, 1H), 4.38-4.47 (m, 1H), 4.80-4.98 (m, 3H), 5.06-5.14 (m, 2H), 5.28 (dd, J=17.2, 1.5 Hz, 1H), 5.41 (brs, 1H), 5.63-5.81 (m, 2H), 7.03 (s, 1H), 7.07 (dd, J=9.2, 2.6 Hz, 1H), 7.41-7.57 (m, 4H), 7.98-8.04 (m, 3H). LC-MS (Method A-retention time: 1.81), MS m/z 741 (M⁺+1).

25




Step 5h: Preparation of (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*trans*-13-
tert-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-
 5 4-yloxy)-2,14-dioxo-3,15-diazatricyclo-
[13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid ethyl ester
 and (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-13-*tert*-butoxycarbonylamino-
 17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-
 diazatricyclo[13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid
 10 ethyl ester

Following the experimental and purification
 procedure of step 1h, 1-{[1-(2(*S*)-*tert*-
 butoxycarbonylamino-oct-7-enoyl)-4-(7-methoxy-2-
 15 phenylquinolin-4(*R*)-yloxy)-pyrrolidine-2(*S*)-carbonyl]-
 1(*R*)-amino]-2(*S*)-vinyl-cyclopropanecarboxylic acid ethyl
 ester (1.16 g, 1.565 mmol) was reacted with 275 mg and
 225 mg portions of Grubbs catalyst to afford
 (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*trans*-13-*tert*-butoxycarbonyl-amino-
 20 17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-
 diazatricyclo-[13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid
 ethyl ester (90 mg, 8%) and (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-13-
tert-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-

4-yloxy)-2,14-dioxo-3,15-diaza-
tricyclo[13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid ethyl
ester (222 mg, 20%). Data for *trans*- or *E*-olefin isomer:
¹H NMR (500 MHz, CD₃OD) δ 1.14-1.48 (m, 6H), 1.21 (t,
5 J=7.0 Hz, 3H), 1.34 (s, 9H), 1.51-1.63 (m, 2H), 1.76 (s,
1H), 1.81-1.91 (m, 2H), 2.00-2.08 (m, 1H), 2.50 (q, J=8
Hz, 1H), 2.61-2.66 (m, 1H), 2.75-2.80 (m, 1H), 3.95 (s,
3H), 4.04-4.18 (m, 2H), 4.40-4.44 (m, 1H), 4.54 (d,
J=11.6 Hz, 1H), 4.79 (t, J=7.3 Hz, 1H), 5.04 (dd, J=15.6,
10 7.6 Hz, 1H), 5.16 (d, J=8.2 Hz, 1H), 5.46 (brs, 1H),
5.53-5.59 (m, 1H), 7.04 (s, 1H), 7.07 (dd, J=9.2, 2.4 Hz,
1H), 7.42-7.52 (m, 4H), 8.03-8.10 (m, 3H). LC-MS (Method
E-retention time: 1.89), MS *m/z* 713 (M⁺+1). Data for *cis*-
or *Z*-olefin isomer: ¹H NMR (300 MHz, CDCl₃) δ 1.15-1.62
15 (m, 5H), 1.21 (t, J=7 Hz, 3H), 1.38 (s, 9H), 1.91 (dd,
J=8.1, 5.9 Hz, 1H), 1.97-2.02 (m, 5H), 2.33-2.41 (m, 1H),
3.03-3.11 (m, 1H), 3.95 (s, 3H), 4.06-4.18 (m, 3H), 4.27
(d, J=11 Hz, 1H), 4.54 (m, 1H), 4.93 (dd, J=8.2, 4.2 Hz,
1H), 5.27-5.40 (m, 2H), 5.71-5.80 (m, 1H), 7.05 (s, 1H),
20 7.08 (dd, J=9.2, 2.6 Hz, 1H), 7.43-7.53 (m, 4H), 8.02 (d,
J=9.2 Hz, 1H), 8.05-8.07 (2s, 2H). LC-MS (Method E-
retention time: 1.80), MS *m/z* 713 (M⁺+1).

59



Trans or E-isomer-
was used in this sequence


Step 5i: Preparation of (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*trans*-13-
tert-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-
4-yloxy)-2,14-dioxo-3,15-diazatricyclo-
[13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid

Following the experimental and purification
procedure of Step 2i, (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*trans*-13-tert-
butoxycarbonyl-amino-17-(7-methoxy-2-phenylquinolin-4-
yloxy)-2,14-dioxo-3,15-diazatricyclo-[13.3.0.0^{4,6}]octadec-
7-ene-4-carboxylic acid ethyl ester (185 mg, 0.265 mmol)
was reacted with 64 mg (1.6 mmol) of LiOH in 15.5 mL of
9:5:1.5 of THF/H₂O/MeOH to afford (1*S*,4*R*,6*S*,13*S*,17*R*)-7-
trans-13-tert-butoxycarbonylamino-17-(7-methoxy-2-
phenylquinolin-4-yloxy)-2,14-dioxo-3,15-
diazatricyclo[13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid
(180mg, 100%): ¹H NMR (300 MHz, CD₃OD) δ 1.14 (s, 9H),
1.20-1.51(m, 5H), 1.53-1.58 (m, 1H), 1.70 (dd, *J*=9.9, 5.1
Hz, 1H), 1.86 (brs, 1H), 2.06 (brs, 1H), 2.28-2.59 (m,
3H), 2.65-2.73 (m, 1H), 3.94 (s, 3H), 4.07 (dd, *J*=11.3,

2.6 Hz, 1H), 4.23 (d, $J=8.4$ Hz, 1H), 4.64–4.70 (m, 2H), 5.54–5.68 (m, 3H), 7.10 (dd, $J=9$, 2 Hz, 1H), 7.26 (s, 1H), 7.39 (d, $J=2.2$ Hz, 1H), 8.02, 8.04 (2s, 2H), 8.12 (d, $J=9.1$ Hz, 1H). LC-MS (Method E-retention time: 1.57),
 5 MS m/z 685 (M^++1). HRMS m/z ($M+H$)⁺ calcd for $C_{38}H_{45}N_4O_8$: 685.3238, found 685.3241.

Step 5j: Preparation of Compound 5, Example 5

(1*S*,4*R*,6*S*,13*S*,17*R*)-7-*trans*-[4-Cyclopropanesulfonylamino-
 10 carbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-
dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octa-dec-7-en-13-
yl]carbamic acid *tert*-butyl ester



15


Using the procedural and purification conditions of Step 2j, 35.6 mg (0.052 mmol) of (1*S*,4*R*,6*S*,13*S*, 17*R*)-7-*trans*-13-*tert*-butoxycarbonyl-amino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo-
 20 [13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid was converted to 30 mg (73%) (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*trans*-[4-cyclopropanesulfonylamino-carbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo-
 [13.3.0.0^{4,6}]octadec-7-en-13-yl]carbamic acid *tert*-butyl

ester: ^1H NMR (300 MHz, CD_3OD) δ 0.85-1.51 (m, 8H), 1.16 (s, 9H), 1.64-1.76 (m, 3H), 1.87 (m, 1H), 2.10 (m, 1H), 2.32 (m, 1H), 2.44 (m, 1H), 2.52-2.72 (m, 2H), 2.83 (brs, 1H), 3.94 (s, 3H), 4.10-4.15 (m, 1H), 4.23-4.32 (m, 1H),
 5 4.61-4.69 (m, 2H), 5.49-5.64 (m, 3H), 7.08 (dd, $J=9.2$, 2.2 Hz, 1H), 7.23 (s, 1H), 7.37 (d, $J=2.2$ Hz, 1H), 7.46-7.56 (m, 3H), 8.03-8.09 (m, 3H). LC-MS (Method E- retention time: 1.57), MS m/z 788 (M^++1). HRMS m/z ($\text{M}+\text{H}$) $^+$ calcd for $\text{C}_{41}\text{H}_{50}\text{SN}_5\text{O}_9$: 788.3329, found 788.3337.

10

Step 6k: Preparation of Compound 6, Example 6

(1S,4R,6S,13S,17R)-7-trans-Cyclopropanesulfonic acid [13-amino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo [13.3.0.0^{4,6}]octadec-7-ene-4-
 15 carbonyl]amide dihydrochloride




Compound 6, Example 6

A solution of 2 mL (8 mmol) of 4N HCl/dioxane was
 20 added to 10.5 mg (0.014 mmol) of (1S,4R,6S,13S,17R)-7-trans-[4-cyclopropanesulfonylaminocarbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo-[13.3.0.0^{4,6}]octadec-7-en-13-yl]carbamic acid *tert*-butyl


ester, the mixture stirred 2h, concentrated and dried (50 °C; 20 torr; 12h) to afford 10.0 mg (~100%) of (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*trans*-Cyclopropanesulfonic acid [13-amino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-7-ene-4-carbonyl]amide dihydrochloride as a white solid: ¹H NMR (300 MHz, CD₃OD) δ 0.86-2.03 (m, 14H), 2.17 (brs, 1H), 2.52-3.00 (m, 3H), 4.06 (s, 3H), 4.16-4.53 (m, 4H), 5.36 (t, *J*=10.6 Hz, 1H), 5.65-5.72 (m, 1H), 5.97 (brs, 1H), 7.44-7.47 (m, 1H), 7.59-7.74 (m, 5H), 8.12 (m, 2H), 8.38-8.41(m, 1H). LC-MS (Method E-retention time: 1.16), MS *m/z* 688 (*M*⁺+1). HRMS *m/z* (*M*+H)⁺ calcd for C₃₆H₄₂SN₅O₇: 688.2805, found 688.2793.

15

Examples 7,8Preparation of Compounds 7 and 8

Compound 7, Example 7 - R is Boc
Compound 8, Example 8 - R is H

Step 7h: Preparation of (1S,4R,6S,13S,17R)-7-cis-13-tert-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo-[13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid ethyl ester




5

This step of this particular sequence uses the product of step 5g as the starting point. To a solution of 1-[[1-(2(*S*)-*tert*-butoxycarbonylamino-oct-7-enoyl)-4-(7-methoxy-2-phenylquinolin-4(*R*)-yloxy)pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino]-2(*S*)-vinylcyclopropanecarboxylic acid ethyl ester (1.44 g, 1.93 mmol) in 2.5L of Argon degassed 1,2-DCE, was added 150 mg of Grubbs catalyst, the mixture degassed under Ar and heated to reflux for 3 h. The reaction mixture was cooled to rt, the mixture degassed again, 140 mg portion of Grubbs catalyst added, and the mixture heated to reflux for 14 h. The reaction mixture was cooled again to rt, the mixture degassed again, 140 mg portion of Grubbs catalyst added, and the mixture heated to reflux for a final 4 h. The dark brown solution was concentrated in vacuo, purified by a Biotage 40 M column (eluted with 15% to 65% EtOAc in hexanes) to supply (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-13-*tert*-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-4-

yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-
 7-ene-4-carboxylic acid ethyl ester (720 mg, 52%). Data
 for *cis*- or *Z*-olefin isomer, (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-13-
 tert-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-
 4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octa-
 dec-7-ene-4-carboxylic acid ethyl ester: ¹H NMR (300
 MHz, CDCl₃) δ 1.15-1.62 (m, 5H), 1.21 (t, *J*=7 Hz, 3H),
 1.38 (s, 9H), 1.91 (dd, *J*=8.1, 5.9 Hz, 1H), 1.97-2.02 (m,
 5H), 2.33-2.41 (m, 1H), 3.03-3.11 (m, 1H), 3.95 (s, 3H),
 4.06-4.18 (m, 3H), 4.27 (d, *J*=11 Hz, 1H), 4.54 (m, 1H),
 4.93 (dd, *J*=8.2, 4.2 Hz, 1H), 5.27-5.40 (m, 2H), 5.71-
 5.80 (m, 1H), 7.05 (s, 1H), 7.08 (dd, *J*=9.2, 2.6 Hz, 1H),
 7.43-7.53 (m, 4H), 8.02 (d, *J*=9.2 Hz, 1H), 8.05-8.07 (2s,
 2H). LC-MS (Method E-retention time: 1.80), MS *m/z* 713
 (M⁺+1).

Step 7i: Preparation of (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-13-*tert*-
 butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-4-
 yloxy)-2,14-dioxo-3,15-diazatricyclo-[13.3.0.0^{4,6}]octa-
 dec-7-ene-4-carboxylic acid




Following the experimental and purification
 procedure of Step 1i, (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-13-*tert*-
 butoxycarbonylamino-17-(7-methoxy-2-phenyl-quinolin-4-

ylxy)-2,14-dioxo-3,15-diazatricyclo [13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid ethyl ester (290 mg, 0.406 mmol) was reacted with 100 mg (2.5 mmol) of LiOH in 23.9 mL of 14.4:7.7:1.8 of THF/H₂O/MeOH to afford

- 5 (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-13-*tert*-butoxy-carbonylamino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo-[13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid (279 mg, 100%): ¹H NMR (300 MHz, CD₃OD) δ 0.97-1.05 (m, 2H), 1.14 (s, 9H), 1.20-1.55 (m, 5H), 1.60-1.73 (m, 2H),
10 1.85 (brs, 1H), 2.06-2.22 (m, 1H), 2.29 (brs, 1H), 2.52-2.73 (m, 2H), 3.93 (s, 3H), 4.14-4.19 (m, 1H), 4.26 (d, *J*=9.5 Hz, 1H), 4.56-4.67 (m, 2H), 5.47-5.53 (m, 2H), 5.71-5.78 (m, 1H), 7.08 (dd, *J*=9.1, 2.2 Hz, 1H), 7.22 (s, 1H), 7.37 (d, *J*=2.2 Hz, 1H), 7.45-7.56 (m, 3H), 8.02,
15 8.04 (2s, 2H), 8.09 (d, *J*=9.1 Hz, 1H). LC-MS (Method E-retention time: 1.60), MS *m/z* 685 (*M*⁺+1).

Step 7j: Preparation of Compound 7, Example 7

- 20 (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-[4-Cyclopropanesulfonylamino-carbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-7-en-13-yl]carbamic acid *tert*-butyl ester



Following the procedural and purification conditions of Step 1j, 219 mg (0.31 mmol) of (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-13-*tert*-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo-


5 [13.3.0.0^{4,6}]octadec-7-ene-4-carboxylic acid was converted to 94 mg (40%) (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-[4-cyclopropanesulfonylamino-carbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-

10 *tert*-butyl ester: ¹H NMR (300 MHz, CD₃OD) δ 0.90-1.50 (m, 9H), 1.16 (s, 9H), 1.64-1.74 (m, 2H), 1.85 (brs, 1H), 2.08 (brs, 1H), 2.30 (brs, 1H), 2.44 (brs, 1H), 2.53-2.71 (m, 2H), 2.84 (brs, 1H), 3.92 (s, 3H), 4.05-4.12 (m, 1H), 4.24 (d, *J*=9.5 Hz, 1H), 4.60-4.69 (m, 2H), 5.44-5.63 (m,

15 3H), 7.06 (dd, *J*=9.2, 2.2 Hz, 1H), 7.20 (s, 1H), 7.35 (d, *J*=2.2 Hz, 1H), 7.45-7.56 (m, 3H), 8.02-8.08 (m, 3H). LC-MS (Method D-retention time: 1.55), MS *m/z* 788 (*M*⁺+1).

Step 8k: Preparation of Compound 8, Example 8

(1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-Cyclopropanesulfonic acid [13-amino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-7-ene-4-carbonyl]amide dihydrochloride



Compound 8, Example 8


A solution of 5 mL (20 mmol) of 4N HCl/dioxane was added to 21 mg (0.027 mmol) of (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*Cis*-[4-Cyclopropanesulfonylamino-carbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-7-en-13-yl]carbamic acid *tert*-butyl ester, the mixture stirred 2h, concentrated and dried (50 °C; 20 torr; 12h) to afford 20 mg (~100%) of (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-Cyclopropanesulfonic acid [13-amino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-7-ene-4-carbonyl]amide dihydrochloride as a white solid: ¹H NMR (500 MHz, CD₃OD) δ 1.03-1.16 (m, 3H), 1.25-1.33 (m, 2H), 1.35-1.57 (m, 3H), 1.65-1.71 (m, 2H), 1.76-1.84 (m, 1H), 1.94-2.01 (m, 1H), 2.10-2.21 (m, 2H), 2.38-2.43 (m, 1H), 2.67-2.72 (m, 1H), 2.85 (dd, *J*=14.6, 7.9 Hz, 1H), 2.93-

2.98 (m, 1H), 4.06 (s, 3H), 4.24 (dd, $J=12.2$, 3.7 Hz, 1H), 4.40, 4.42 (2s, 2H), 4.80-4.81 (m, 1H), 5.53-5.57 (m, 1H), 5.67-5.74 (m, 1H), 5.95 (s, 1H), 7.45 (dd, $J=9$, 2 Hz, 1H), 7.57 (d, $J=2$ Hz, 1H), 7.61 (s, 1H), 7.70-7.77 (m, 3H), 8.09-8.10 (m, 2H), 8.37 (d, $J=9.2$ Hz, 1H). LC-MS (Method E-retention time: 1.21), MS m/z 688 ($M^+ + 1$).

Examples 9, 10

Preparation of Compounds 9 and 10


10



Compound 9, Example 9 - R is Boc
Compound 10, Example 10 - R is H (Bis HCl Salt)


Step 9g: Preparation of 1-{[1-(2(*S*)-tert-butoxycarbonyl-amino-non-8-enoyl)-4-(7-methoxy-2-phenylquinolin-4(*R*)-yloxy)pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino]-2(*S*)-vinylcyclopropanecarboxylic acid ethyl ester

5



Following the procedure of Step 1g, except that
 (0.78g, 3.0mmol) of 2(*S*)-tert-butoxycarbonylamino-8-
 10 nonenoic acid purchased from RSP Amino Acids, was coupled
 in place of 2(*S*)-tert-butoxycarbonylamino-5-hexenoic
 acid, gave 1-{[1-(2(*S*)-tert-butoxycarbonylamino-non-8-
 enoyl)-4-(7-methoxy-2-phenylquinolin-4(*R*)-
 yloxy)pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino}-2(*S*)-
 15 vinylcyclopropanecarboxylic acid ethyl ester as a yellow
 oil (1.55g, 82%): ^1H NMR (300 MHz, CDCl_3) δ 1.15-1.78 (m,
 9H), 1.20 (t, $J=7$ Hz, 3H), 1.39 (s, 9H), 1.84 (dd, $J=8.1$,
 5.5 Hz, 1H), 1.91-2.04 (m, 2H), 2.14 (q, $J=8.8$ Hz, 1H),
 2.35-2.48 (m, 1H), 2.89-3.00 (m, 1H), 3.93 (s, 3H), 4.02-
 20 4.20 (m, 2H), 4.28 (d, $J=11.7$ Hz, 1H), 4.39-4.47 (m, 1H),
 4.80-4.99 (m, 1H), 5.06-5.14 (m, 2H), 5.27 (dd, $J=17$, 1.3
 Hz, 1H), 5.40 (brs, 1H), 5.65-5.81 (m, 2H), 7.03 (s,
 1H), 7.07 (dd, $J=9.1$, 2.6 Hz, 1H), 7.42-7.57 (m, 4H),
 7.99-8.05 (m, 3H). LC-MS (Method E-retention time: 1.79),
 25 MS m/z 755 (M^++1).


70



Step 9h: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-
 butoxycarbonylamino-18-(7-methoxy-2-phenylquinolin-4-
 5 yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-
 7-ene-4-carboxylic acid ethyl ester

To a solution of 1-[[1-(2(*S*)-*tert*-
 butoxycarbonylamino-non-8-enoyl)-4-(7-methoxy-2-
 10 phenylquinolin-4(*R*)-yloxy)-pyrrolidine-2(*S*)-carbonyl]-
 1(*R*)-amino]-2(*S*)-vinylcyclopropanecarboxylic acid ethyl
 ester (1.50 g, 1.99 mmol) in 2.4L of Argon degassed 1,2-
 DCE, was added 150 mg of Grubbs catalyst, the mixture
 degassed under Ar and heated to reflux for 3 h. The
 15 reaction mixture was cooled to rt, the mixture degassed
 again, 150 mg portion of Grubbs catalyst added, and the
 mixture heated to reflux for 3 h. The reaction mixture
 was cooled again to rt, the mixture degassed again, 150
 mg portion of Grubbs catalyst added, and the mixture
 20 heated to reflux for a final 9 h. The dark brown
 solution was concentrated in vacuo, purified by a Biotage
 40 M column (eluted with 15% to 65% EtOAc in hexanes) to
 supply (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-
 butoxycarbonylamino-18-(7-methoxy-2-phenylquinolin-4-

yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-
 7-ene-4-carboxylic acid ethyl ester (1.08 g, 75%): ¹H NMR
 (300 MHz, CDCl₃) δ 1.10-1.42 (m, 6H), 1.17 (t, J=7 Hz,
 3H), 1.33 (s, 9H), 1.54 (dd, J=9.5, 5.5 Hz, 1H), 1.62 (m,
 5 1H), 1.73 (dd, J=8.5, 5.5 Hz, 1H), 1.75-1.86 (m, 1H),
 2.01-2.24 (m, 3H), 2.39-2.47 (m, 1H), 2.81-2.89 (m, 1H),
 3.91 (s, 3H), 3.98-4.17 (m, 3H), 4.34 (d, J=11.0 Hz, 1H),
 4.46 (m, 1H), 4.77 (dd, J=8.1, 5.5 Hz, 1H), 5.21-5.27 (m,
 1H), 5.39 (m, 1H), 5.48-5.57 (m, 1H), 6.99 (s, 1H), 7.04
 10 (dd, J=9.1, 2.2 Hz, 1H), 7.38 (d, J=2.2 Hz, 1H), 7.42-
 7.50 (m, 3H), 7.95-8.01 (m, 3H). LC-MS (Method E-
 retention time: 1.71), MS m/z 727 (M⁺+1).



15 Step 9i: Preparation of (1S,4R,6S,14S,18R)-7-cis-14-tert-
butoxycarbonylamino-18-(7-methoxy-2-phenylquinolin-4-
yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-
7-ene-4-carboxylic acid


20 Following an analogous experimental and purification
 procedure to Step 1i, (1S,4R,6S,14S,18R)-7-cis-14-tert-
 butoxycarbonylamino-18-(7-methoxy-2-phenylquinolin-4-
 yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-
 7-ene-4-carboxylic acid ethyl ester (225 mg, 0.309 mmol)

was reacted with 78 mg (2.0 mmol) of LiOH in 20.8 mL of 12.4:6.6:1.8 of THF/H₂O/MeOH to afford (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-butoxycarbonylamino-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid (178 mg, 82%): ¹H NMR (500 MHz, CD₃OD) δ 1.12-1.54 (m, 8H), 1.19 (s, 9H), 1.59 (dd, *J*=8.4, 5.0 Hz, 1H), 1.64 (dd, *J*=9.8, 4.9 Hz, 1H), 1.75-1.82 (m, 1H), 1.87-1.94 (m, 1H), 2.31 (q, *J*=9 Hz, 1H), 2.52-2.57 (m, 1H), 2.69 (dd, *J*=14, 7.5 Hz, 1H), 3.92 (s, 3H), 4.01 (dd, *J*=11.6, 3.2 Hz, 1H), 4.19 (dd, *J*=10.4, 2.4 Hz, 1H), 4.65 (t, *J*=8 Hz, 1H), 4.75 (d, *J*=11.6 Hz, 1H), 5.33 (t, *J*=9.6 Hz, 1H), 5.55-5.60 (m, 2H), 7.08 (dd, *J*=9.2, 2.1 Hz, 1H), 7.25 (s, 1H), 7.37 (d, *J*=2 Hz, 1H), 7.51-7.57 (m, 3H), 8.02-8.05 (m, 2H), 8.15 (d, *J*=9.2 Hz, 1H). LC-MS (Method E-retention time: 1.61), MS *m/z* 699 (M⁺+1). HRMS *m/z* (M+H)⁺ calcd for C₃₉H₄₇N₄O₈: 699.3394, found 699.3395.

Step 9j: Preparation of Compound 9, Example 9

20 (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-[4-Cyclopropanesulfonylamino-carbonyl-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid *tert*-butyl ester

73




Compound 9, Example 9

Following the procedure of Step 1j, 136.5 mg (0.195 mmol) of (1*S*,4*R*,6*S*,13*S*,18*R*)-7-*cis*-14-*tert*-
 5 butoxycarbonylamino-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid was converted to 115 mg (76%) (1*S*,4*R*,6*S*,13*S*,18*R*)-7-*cis*-[4-Cyclopropanesulfonylamino-
 10 carbonyl-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid *tert*-butyl ester: ¹H NMR (500 MHz, CD₃OD) 0.97-1.54 (m, 11H), 1.20 (s, 9H), 1.59 (m, 1H), 1.71 (m, 1H), 1.75-1.89 (m, 2H), 2.35-2.43 (m, 1H), 2.47-2.75 (m, 3H), 2.87-2.91 (m, 1H),
 15 3.92 (s, 3H), 3.97-4.05 (m, 1H), 4.17 (d, *J*=10.4 Hz, 1H), 4.64 (t, *J*=7.6 Hz, 1H), 5.07 (m, 1H), 5.54 (brs, 1H), 5.63 (brs, 1H), 7.04 (d, *J*=8.6 Hz, 1H), 7.21 (brs, 1H), 7.36 (s, 1H), 7.48-7.55 (m, 3H), 8.02, 8.03 (2s, 2H), 8.15 (d, *J*=8.5 Hz, 1H). LC-MS (Method E-retention time:
 20 1.64), MS *m/z* 802 (*M*⁺+1). HRMS *m/z* (*M*+H)⁺ calcd for C₄₂H₅₂SN₅O₉: 802.3486, found 802.3481.

Step 10k: Preparation of Compound 10, Example 10

Preparation of (1S,4R,6S,14S,18R)-7-cis-Cyclopropane-
sulfonic acid [14-amino-18-(7-methoxy-2-phenylquinolin-4-
yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-


5 nonadec-7-ene-4-carbonyl]amide dihydrochloride




Compound 10, Example 10

A solution of 5 mL (20 mmol) of 4N HCl/dioxane was
 10 added to 26 mg (0.032 mmol) of (1S,4R,6S,13S,18R)-7-cis-
 [4-cyclopropanesulfonylaminocarbonyl-18-(7-methoxy-2-
 phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo-
 [14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid tert-butyl
 ester, the mixture stirred 2h, concentrated and dried (50
 15 °C; 20 torr; 12h) to afford 24.0 mg (~100%) of
 (1S,4R,6S,14S,18R)-7-cis-cyclopropanesulfonic acid [14-
 amino-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-
 3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-
 carbonyl]amide dihydrochloride as a white solid: ¹H NMR
 20 (300 MHz, CD₃OD) δ 0.96-2.05 (m, 16H), 2.24-2.34 (m, 1H),
 2.52 (ms, 1H), 2.73 (m, 1H), 2.91 (m, 2H), 3.57-3.74 (m,
 1H), 4.06 (s, 3H), 4.17 (m, 1H), 4.37 (m, 1H), 4.48 (m

2H), 5.07-5.14 (m, 1H), 5.66-5.75 (m, 1H), 5.95 (m, 1H), 7.47 (m, 1H), 7.60-7.75 (m, 5H), 8.12 (m, 2H), 8.39-8.41 (m, 1H) LC-MS (Method E-retention time: 1.24), MS m/z 702 ($M^+ + 1$). HRMS m/z ($M + H$)⁺ calcd for C₃₇H₄₄SN₅O₇: 702.2962, found 702.2976.



Compound 11, Example 11



Compound 12, Example 11

Compounds 11/12, Example 11

10

Preparation of Compound 11, Example 11,

(1S,4R,6S,13S,18R)-7-cis-[4-Cyclopropanesulfonyl-
aminocarbonyl-18-(7-methoxy-2-phenylquinolin-4-yloxy)-
2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nona-dec-7-en-


15

14-yl]carbamic acid tert-butyl ester and Compound 12,

Example 11, (1S,4R,6S,13R,18R)-7-cis-[4-
Cyclopropanesulfonylaminocarbonyl-18-(7-methoxy-2-phenyl-
quinolin-4-yloxy)-2,15-dioxo-3,16-diazatri-cyclo-
[14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid tert-butyl

20

ester was accomplished using steps 11A-J.



Steps 11A-11B: Preparation of 7-Methoxyquinolin-4-ol
5 from *m*-anisidine.

Step 11A) To a solution of *m*-anisidine (58 g, 471 mmol) in 800 mL of CH₃CN was added Meldrum's acid (75g, 518 mmol), and trimethylformate (60g, 565 mmol). The
10 heterogeneous mixture was refluxed for 2 h went into solution. The solvent was removed in vacuo, 30 mL of MeOH added, and the resulting precipitate filtered and washed with 10-15 mL of MeOH. The MeOH
addition/filtration procedure was repeated on the
15 concentrated mother liquor. The resulting combined solid was dried (~20 torr, 45 °C overnite) to afford (117.6g, 90%) of the intermediate 5-[(3-Methoxyphenyl-amino)methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione.

Step 11B) To a solution of 500 g of Ph₂O heated to 250 °C was added 108.7g (392 mmol) of 5-[(3-Methoxyphenyl-amino)methylene]-2,2-dimethyl-[1,3]dioxane-4,6-dione in portions over a 30 min period. The mixture

was heated an additional 15 min, cooled to rt, diluted with 800 mL of hexanes and the resulting slurry stirred overnite. The hexanes was decanted off, the solid residue dissolved in 600 mL of MeOH at reflux, cooled to
5 rt and the resulting solid filtered and washed with minimal CH₂Cl₂. The analogous recrystallization procedure was followed to afford a total of 20.73 g (30%) of 7-methoxyquinolin-4-ol as a light brown solid. ¹H NMR (methanol-d₄) δ 3.87 (s, 3H), 6.23 d, *J*=7.3 Hz, 1H), 6.68
10 (d, *J*=2.4 Hz, 1H), 6.96 (dd, *J*=9.0, 2.4 Hz, 1H), 8.11 (d, *J*=9 Hz, 1H); LC-MS (retention time: 0.77, method D), MS *m/z* 176 (M⁺+1).

Step 11C: Preparation of BOC-N-P2[(4R)-(7-methoxyquinoline-4-oxo)proline methyl ester]4-(7-methoxyquinolin-4yloxy)-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester from BOC-L-CIS-HYP-OH and 7-methoxyquinolin-4-ol.

15

Step 11C) To a solution of 12.24 g (49.8 mmol) of BOC-L-CIS-HYP-OH (BOC-CIS-HYP-OME, *N*-α-*tert*-butoxycarbonyl-*cis*-L-4-Hydroxyproline methylester) and 26.14g (99.7 mmol) of PPh₃ in 200 mL of THF cooled to 0 °C, was added a solution of 17.36 g (99.7 mmol) of DEAD
25 and 8.73 g (49.8 mmol) of 7-methoxyquinolin-4-ol in 700 mL of THF over a 45 min period. The mixture was slowly allowed to warm to rt overnite, concentrated in vacuo and the residue purified over a Biotage 65M (eluted with 0% to 10% MeOH-EtOAc) to afford 12.78g (64%) of the desired
30 product BOC-N-P2[(4R)-(7-methoxyquinoline-4-oxo)pro-line methyl ester], 4-(7-methoxyquinolin-4yloxy)-pyrrolidine-1,2-dicarboxylic acid 1-*tert*-butyl ester 2-methyl ester as a colorless glass. ¹H NMR (CDCl₃) δ 1.36 (s, 9H),


2.26-2.35 (m, 1H), 2.57-2.68 (m, 1H), 3.71 (s, 3H), 3.75-3.92 (m, 2H), 3.86, 3.87 (two s (rotamers) 3H), 4.41-4.53 (m, 1H), 5.09 (m, 1H), 6.52 (d, $J=5.5$ Hz, 1H), 7.06-7.09 (m, 1H), 7.24-7.26 (m, 1H), 7.94 (d, $J=9.1$ Hz, 1H),
5 8.50-8.56 (m, 1H); LC-MS (retention time: 1.34, method D), MS m/e 403 ($M^+ + 1$).

Step 11D: Preparation of BOC-N-P2[(4R)-(7-methoxyquinoline-4-oxo)proline]-OH, 4-(7-methoxyquinolin-4-yl-oxy)-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester from BOC-N-P2[(4R)-(7-methoxyquinoline-4-oxo)proline methyl ester], 4-(7-methoxyquinolin-4-yloxy)-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl ester 2-methyl ester.

15

Step 11D) To a solution of 8.54 g (21.2 mmol) of BOC-N-P2[(4R)-(7-methoxyquinoline-4-oxo)proline methyl ester] in 600 mL of 5:1 THF/MeOH, was added a solution of 4.0 g (167 mmol) of LiOH in 150 mL of water. The mixture
20 was stirred overnight, the pH adjusted to pH 7 using 6N aqueous HCl, and the solution concentrated until only the water layer remained. The solution was adjusted to pH 4 using 6N aqueous HCl, NaCl added to saturate the mixture and was partitioned repeatedly with first EtOAc and then
25 THF as the product was aqueous soluble. The combined organic layers were dried ($MgSO_4$) and concentrated to afford 8.18 g (99%) of BOC-N-P2[(4R)-(7-methoxyquinoline-4-oxo)proline]-OH, 4-(7-methoxyquinolin-4-yloxy)-pyrrolidine-1,2-dicarboxylic acid 1-tert-butyl
30 ester as a white solid. 1H NMR ($CDCl_3$ -Methanol- d_4) δ 1.42 (s, 9H), 2.40-2.49 (m, 1H), 2.68-2.77 (m, 1H), 3.88 (m, 2H), 3.94 (s, 3H), 4.41-4.53 (m, 1H), 5.32 (m, 1H), 6.86-6.92 (m, 1H), 7.21 (dd, $J=9, 2$ Hz, 1H), 7.30 (d, $J=2$ Hz,

1H), 8.05-8.10 (m, 1H), 8.62 (d, $J=6$ Hz, 1H); LC-MS (retention time 1.20, method A), MS m/z 389 ($M^+ + 1$).



5

Step 11E: Coupling of (1R,2S/1S,2R)-Vinyl Acca)-CO₂Et HCl Salt P1 Moiety with Boc (4R)-(7-methoxyquinoline-4-oxo)-S-proline P2 Moiety to make BOCN-P2[(4R)-7-methoxyquinoline-4-oxo proline]-P1(1R,2S/1S,2R)-vinylAcca-COOEt and chromatographic separation of BOCN-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S VinylAcca)-CO₂Et (most bioactive isomer) and BOCN-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1S,2R Vinyl Acca)-CO₂Et Diastereomers.


15

Step 11E) To a solution of 4.50 g (11.60 mmol) of Boc-4(R)-(7-methoxyquinoline-4-oxo)proline, 2.66 g (13.9 mmol) of the HCl salt of vinyl Acca (existing as a 1:1 mixture of diastereoisomers (1R,2S/1S,2R where cyclopropyl carboxyethyl group is syn to vinyl moiety), 10 mL (57.4 mmol) of DIPEA, and 2.13 g (13.9 mmol) of HOBT.H₂O in 150 mL of CH₂Cl₂ was added 5.27 g (13.9 mmol) of the coupling reagent HBTU, and the mixture stirred overnight. The solution was diluted with 200 mL of CH₂Cl₂ and was partitioned with pH 4.0 buffer (2x50 mL). The organic layer was washed with saturated aqueous NaHCO₃ (2x50mL), water (2x50mL), and brine (2x50mL). The organic solution was dried (MgSO₄), concentrated and

25

purified using a Biotage 65M column (eluted with 0% to 9% MeOH/EtOAc) to provide of BOC-NH-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S vinyl acca P1 moiety)-COOEt as the initial eluted isomer (2.21 g, 36% overall), followed by 1.13 g (19%) of pure lower Rf isomer BOC-NH-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1S,2R Vinyl Acca P1 moiety)-CO₂Et. Mixed fractions were also obtained. Data for BOCN-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S)-(VinylAcca)-COOEt: ¹H NMR (CDCl₃) δ 1.16 (t, J=7 Hz, 3H), 1.35 (s, 9H), 1.37-1.47 (m, 1H), 1.74-1.88 (m, 1H), 2.04-2.13 (m, 1H), 2.32-2.46 (m, 1H), 2.58-2.69 (m, 1H), 3.76 (m, 1H), 3.87 (s, 3H), 4.02-4.13 (m, 2H), 4.30-4.44 (m, 1H), 5.05-5.19 (m, 2H), 5.24 (d, J=17 Hz, 1H), 5.63-5.71 (m, 1H), 6.61 (m, 1H), 7.07 (dd, J=9, 2 Hz, 1H), 7.22 (d, J=2 Hz, 1H), 7.76-7.83 (m, 1H), 7.92 (d, J=9 Hz, 1H), 8.50 (d, J=5 Hz, 1H). LC-MS (retention time: 1.38, method A), MS m/z 526 (M⁺+1).

Data for BOC-NH-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1S,2R Vinyl Acca P1 moiety)-CO₂Et: ¹H NMR (methanol-d₄) δ 1.23 (t, J=7 Hz, 3H), 1.38 (s, 9H), 1.45 (dd, J=9.5, 5.5 Hz, 1H), 1.77-1.87 (m, 1H), 2.10-2.16 (m, 1H), 2.37 (m, 1H), 2.73 (m, 1H), 3.77 (m, 1H), 3.90 (s, 3H), 4.07-4.15 (m, 2H), 4.46 (m, 1H), 5.10 (d, J=10 Hz, 1H), 5.21 (m, 1H), 5.27 (d, J=17 Hz, 1H), 5.69-5.76 (m, 1H), 6.64 (m, 1H), 7.10 (dd, J=9.2, 2.2 Hz, 1H), 7.27 (m, 1H), 7.95 (d, J=9.2 Hz, 1H), 8.56 (d, J=5 Hz, 1H). LC-MS (retention time: 1.45, method B), MS m/z 526 (M⁺+1).



Step 11F: Preparation of N-BOC-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S VinylAcCa)-CO₂H or (1R,2S) Vinyl Acca P1 Isomer of {1-[2-(1-Cyclopropanesulfonylaminocarbon-yl-2-vinyl-cyclopropylcarbamoyl)-4-(7-methoxyquinolin-4-yloxy)pyrrolidine-1-carbonyl]-2,2-dimethylpropyl}-carbamic acid tert-butyl ester.

10

Step 11F) To a solution of 794 mg (1.51 mmol) of N-BOC-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S VinylAcCa)-CO₂Et (from step 11E) in 68 mL of 12% MeOH/THF was added a solution of 218 mg (9.08 mmol) of lithium hydroxide in 30 mL of water and the mixture stirred 16h. The pH was adjusted to neutral by addition of 6N aqueous HCl, concentrated until only the water remained, the solution adjusted to pH 4 using aqueous 1N HCl and was then extracted with 50% THF-EtOAc (5 X 200-mL portions). The combined organic layers were dried (MgSO₄) and concentrated to afford 808 mg of crude which was dried further to provide 752 mg (100%) of N-BOC-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S VinylAcCa)-CO₂H. ¹H NMR (Methanol-d₄) δ 1.37-1.43 (m, 1H), 1.39 (s, 9H), 1.69-1.78 (m, 1 H), 2.16-2.24 (m, 1H), 2.44-2.54 (m, 1H), 2.64-2.74 (m, 1H), 3.89-3.94 (m, 2H),

25

3.96 (s, 3H), 4.40-4.43 (m, 1H), 5.11 (d, $J=10$ Hz, 1H),
5.31 (d, $J=17$ Hz, 1H), 5.40 (m, 1H), 5.79-5.87 (m, 1H),
6.91 (s, 1H), 7.04 (d, $J=6$ Hz, 1H), 7.25 (dd, $J=9.1$, 2
Hz, 1H), 7.29 (m, 1H), 8.09 (d, $J=9.1$ Hz, 1H), 8.66 (d,
5 $J=6$ Hz, 1H). LC-MS (retention time: 1.05, method H). MS
 m/z 498 (M^++1).

Step 11G: Preparation of N-BOC-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S VinylAcCa)-
10 CONHSO₂Cyclopropane or (1R,2S) Vinyl AcCa P1 Diastereomer
of 2-(1-Cyclopropane-sulfonylamino-carbonyl-2-
vinylcyclopropylcarbamoyl)-4-(7-methoxy-quinolin-4-
xyloxy)pyrrolidine-1-carboxylic acid tert-butyl ester.

15 **Step 11G)** To a solution of 399.5 mg (0.668 mmol)
of N-BOC-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-
P1(1R,2S VinylAcCa)-CO₂H in THF (4 mL), was added CDI
(434 mg, 2.68 mmol), and the resulting solution refluxed
for 60 min and allowed to cool down to rt.
20 Cyclopropylsulfonamide (406 mg, 3.35 mmol) was added in
one portion before the addition of a neat solution of
DBU (0.50 mL, 3.35 mmol). The reaction was stirred for 16
h, diluted with 50% THF-EtOAc (200mL) and washed with
brine saturated pH 4.0 buffer (2x40mL), dried (MgSO₄) and
25 chromatographed over a Biotage 25M column (eluted with 0%
to 15% MeOH in CH₂Cl₂) to supply 217 mg (54%) of the
desired product N-BOC-P2[(4R)-(7-methoxyquinoline-4-oxo)-
S-proline]-P1(1R,2S VinylAcCa)-CONHSO₂Cyclopropane. ¹H
NMR (Methanol-d₄) δ 1.01-1.10 (m, 2H), 1.11-1.18 (m, 1H),
30 1.20-1.27 (m, 1H), 1.39-1.48 (m, 1H), 1.44 (s, 9H), 1.87
(dd, $J=8$, 5 Hz, 1H), 2.01-2.38 (m, 2H), 2.57 (dd, $J=14$, 7
Hz, 1H), 2.91-2.96 (m, 1H), 3.83-3.92 (m, 2H), 3.94 (s,
3H), 4.36-4.39 (m, 1H), 5.11 (d, $J=10$ Hz, 1H), 5.29 (d,

$J=17$ Hz, 1H), 5.38 (m, 1H), 5.74-5.81 (m, 1H), 6.91 (d, $J=5.5$ Hz, 1H), 7.20 (dd, $J=9.2$, 2.4 Hz, 1H), 7.29 (m, 1H), 8.07 (d, $J=9.2$ Hz, 1H), 8.60 (d, $J=5.5$ Hz, 1H). LC-MS (retention time: 1.28, method I). MS m/z 601 ($M^+ + 1$)

5

Steps 11H and 11I: Preparation of BOCNH-P3(2(S)-tert-butoxycarbonylamino-8-nonenoyl)-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S-VinylAcca)-CONHSO₂Cyclopropane or (1R,2S) P1 Isomer of {1-[2-(1-Cyclopropanesulfonylamino-carbonyl-2-vinyl-cyclopropyl-carbamoyl)-4-(7-methoxyquinolin-4-yloxy)-pyrrolidine-1-carbonyl]-oct-7-enyl} carbamic acid tert-butyl ester.


15 **Steps 11H)** A total of 600 mg (1.0 mmol) of BOC-P2 [(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S Vinyl Acca-CONHSO₂Cyclopropane), (1R,2S) VinylAcca P1 isomer of 2-(1-Cyclopropanesulfonylamino-carbonyl-2-vinylcyclopropyl-carbamoyl)-4-(7-methoxyquinolin-4-yloxy)pyrrolidine-1-carboxylic acid tert-butyl ester, was
20 dissolved in 4N HCl/dioxane (4ml, 16 mmol) and was stirred for 3h at rt. The reaction mixture was concentrated to supply the crude HN-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S Vinyl Acca)
25 -CONHSO₂Cyclopropane, Bis HCl Salt: (1R,2S)-Vinyl Acca P1 isomer of 4-(7-Methoxyquinolin-4-yloxy)-pyrrolidine-2-carboxylic acid (1-cyclopropane-sulfonylamino-carbonyl-2-vinylcyclopropyl)amide dihydrochloride as a tan solid which was used immediately in Step 11I.

30

Step 11I) The HN-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S Vinyl Acca)-CONHSO₂Cyclopropane, Bis HCl Salt from step 11H, was suspended in 20 mL of

dichloromethane. To this mixture was added 352 mg (1.30 mmol) of 2(*S*)-*tert*-butoxycarbonylamino-8-nonenoic acid purchased from RSP Amino Acids, HOAT (82 mg, 0.60 mmol), DIPEA (0.74 ml, 5.0 mmol), and HATU (494 mg, 1.30 mmol) at rt. The reaction mixture was stirred 16h, and the majority of the CH₂Cl₂ removed in vacuo. The mixture was diluted with saturated pH 4.0 buffer (150 mL), and extracted into EtOAc (4x200 mL). The combined organic layers were dried (MgSO₄), concentrated and purified over a Biotage 40M column (eluted with 0% to 15% MeOH in CH₂Cl₂) to afford 574 mg (76%) of BOCNH-P3(2(*R/S*)-*tert*-butoxycarbonylamino-8-nonenoic acid)-P2[(4*R*)-(7-methoxyquinoline-4-oxo)-*S*-proline]-P1(1*R*,2*S*-VinylAcce)-CONHSO₂Cyclopropane or (1*R*,2*S*) P1 Isomer of {1-[2-(1-Cyclopropanesulfonylamino-2-vinyl-cyclopropylcarbamoyl)-4-(7-methoxyquinolin-4-yl)-pyrrolidine-1-carbonyl]-oct-7-enyl}carbamic acid *tert*-butyl ester. LC-MS *m/e* 754 (retention time: 1.64, method I).

85




Preparation of Compound 11, Example 11,
(1S,4R,6S,13S,18R)-7-cis-[4-Cyclopropanesulfonyl-
aminocarbonyl-18-(7-methoxy-2-phenyl-quinolin-4-yloxy)-
2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-en-
5 14-yl]carbamic acid tert-butyl ester and Compound 12,
Example 11, (1S,4R,6S,13R,18R)-7-cis-[4-
Cyclopropanesulfonyl-aminocar-bonyl-18-(7-methoxy-2-
phenylquinolin-4-yloxy)-2,15-dioxo-3,16-
diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-en-14-yl]carbamic
10 acid tert-butyl ester.

Step 11J) To a solution of BOCNH-P3(2(S)-tert-butoxycarbonylamino-8-nonenoyl)-P2[(4R)-(7-methoxyquinoline-4-oxo)-S-proline]-P1(1R,2S-VinylAcCa)-
15 CONHSO₂Cyclopropane or (1R,2S) P1 Isomer of {1-[2-(1-Cyclopropanesulfonylamino-carbonyl-2-vinylcyclopropyl-carbamoyl)-4-(7-methoxyquinolin-4-yloxy)-pyrrolidine-1-carbonyl]-oct-7-enyl}carbamic acid tert-butyl ester from
20 Step 11I (537 mg, 0.712 mmol) in 0.7L of Argon degassed 1,2-DCE, was added 60 mg of Grubbs catalyst, the mixture degassed under Ar and heated to reflux for 4 h. The reaction mixture was cooled to rt, the mixture degassed once more, an additional 60 mg portion of Grubbs catalyst added, and the mixture heated to reflux for 4 h. The
25 reaction mixture was cooled to rt, the mixture degassed once more, and a final 60 mg portion of Grubbs catalyst added, and the mixture heated to reflux for 2 h. The mixture was cooled to rt, stirred 9 h more and the resulting dark brown solution concentrated in vacuo and
30 purified using three Analtech 1000μ PTLC plates (20X40 cm, eluted sequentially with 0% to 3% MeOH in CH₂Cl₂ to afford 80mg of crude product. This material was further purified using four 500 μ E-Merck PTLC plates to afford

10 mg of Compound 11, Example 11, (1*S*,4*R*,6*S*,13*S*,18*R*)-7-
cis-[4-Cyclopropanesulfonyl-aminocarbonyl-18-(7-methoxy-
2-phenyl-quinolin-4-yloxy)-2,15-dioxo-3,16-
diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-en-14-yl]carbamic
5 acid *tert*-butyl ester and P3 Epimer Compound 12, Example
11, (1*S*,4*R*,6*S*,13*R*,18*R*)-7-cis-[4-
Cyclopropanesulfonylaminocarbonyl-18-(7-methoxy-2-phenyl-
quinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo-
[14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid *tert*-butyl
10 ester. Data for Compound 11: ¹H NMR (CD₃OD) δ 0.87-1.10
(m, 4H), 1.15-2.01 (m, 11H), 1.21 (s, 9H), 2.19-2.29 (m,
1H), 2.45-2.73 (m, 3H), 2.83-2.92 (m, 1H), 3.93 (s, 3H),
4.11-4.17 (m, 1H), 4.20-4.29 (m, 1H), 4.60-4.69 (m, 2H),
5.21-5.27 (m, 1H), 5.47 (m, 1H), 5.54-5.63 (m, 1H), 6.96
15 (d, *J*=5.5 Hz, 1H), 7.10 (dd, *J*=9.5, 2.5 Hz, 1H), 7.27 (d,
J=2.5 Hz, 1H), 8.17 (d, *J*=9.5 Hz, 1H), 8.61 (d, *J*=5.5 Hz,
1H). LC-MS *m/e* 726 (retention time: 1.57, method A). Data
for Compound 12: ¹H NMR (CD₃OD) δ 0.79-0.99 (m, 2H), 1.00-
1.65 (m, 9H), 1.15 (s, 9H), 1.80-2.05 (m, 4H), 2.32-2.57
20 (m, 2H), 2.59-2.83 (m, 2H), 2.84-2.92 (m, 1H), 3.92 (s,
3H), 4.03-4.16 (m, 1H), 4.27-4.31 (m, 1H), 4.53-4.78 (m,
2H), 5.42 (m, 3H), 6.91 (d, *J*=5 Hz, 1H), 7.11 (dd, *J*=9.2,
2.2 Hz, 1H), 7.26 (d, *J*=2.2 Hz, 1H), 8.11 (d, *J*=9.2 Hz,
1H), 8.60 (d, *J*=5 Hz, 1H). LC-MS *m/e* 726 (retention time:
25 1.49, method A).


30

35

Examples 12,13Preparation of Compounds 13 and 14

Compound 13, Example 12 - R is Boc
Compound 14, Example 13 - R is H

5




Compound 13, Example 12

Preparation of (1*S*,4*R*,6*R*,13*S*,17*R*)-[4-Cyclopropane-sulfonylaminocarbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]-octadec-13-yl]-carbamic acid tert-butyl ester, Compound 13, Example 12.

Method A. A solution of 30 mg (0.038 mmol) of
10 (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-[4-Cyclopropanesulfonyl-aminocarbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-7-en-13-yl]carbamic acid tert-butyl ester, Compound 7, Example 7, was dissolved in 15 mL of MeOH and then evacuated via
15 vacuum and degassed repeatedly under Ar. To this solution was added 15 mg of 20% Pd(OH)₂/C and the mixture evacuated and put under 1 atmosphere of H₂ gas three times and finally allowed to stir under 1 atmosphere of H₂ over 45 minutes. The reaction contents were then then
20 evacuated and put under Argon, filtered over celite and the resulting crude residue purified over preparative HPLC (column: Xterra C18 S5 30X75 mm, 35% to 80% Solvent B/A for 28 min gradient, hold time 3 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is
25 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). Compound 13, Example 12 eluted first (8.8 mg, 29.3%), followed by 15 mg of mixed fractions containing Compound 13 and an unidentified product: ¹H NMR (methanol-d₄) δ
0.84-2.08 (m, 26H), 2.51-2.69 (m, 1H), 2.74-2.78 (m, 1H),
30 2.91-3.01 (m, 1H), 3.96 (s, 3H), 4.04-4.09 (m, 1H), 4.17 (d, *J*=11 Hz, 1H), 4.72 (t, *J*=8 Hz, 1H), 4.97 (d, *J*=12 Hz, 1H), 5.64 m, 1H), 7.13 (dd, 9, 1.6 Hz, 1H), 7.32 (s, 1H), 7.40 (d, *J*=1.6 Hz, 1H), 7.53-7.62 (m, 3H), 8.02-8.05 (m,

2H), 8.15 (d, $J=9$ Hz, 1H). LC-MS m/e 790 (retention time: 2.55, method D, except gradient time increased from 2 to 3 min).




Compound 13, Example 12- R is BOC
Compound 14, Example 13- R is H

5

Step i of Method B) To a solution of 400 mg (0.584 mmol) of (1S,4R,6S,13S,17R)-7-*cis*-13-*tert*-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octa-dec-7-ene-4-carboxylic acid and 1.90 g (11.7 mmol) of potassium diazodicarboxylate in 20 mL of MeOH, was added a solution of 1.34 mL (22.4 mmol) of glacial AcOH in 19 mL of MeOH slowly dropwise via syringe pump over 5 h, followed by stirring for an additional 6 h. To this mixture was added an additional 380 mg of potassium diazodicarboxylate (prepared as in *Org. React.* **1991**, 40, p.91), followed by 400 μL of glacial AcOH in 4 mL of MeOH via syringe pump over 70 min. The mixture was stirred overnight, diluted with 200 mL of Et₂O, filtered and concentrated in vacuo. The residue was chromatographed over 20 g of SiO₂ (eluted with 5% MeOH/CH₂Cl₂) to afford 450 mg of crude which was dissolved in 12 mL of MeOH and injected six times (6 x 2 mL injections) onto a preparative HPLC column: Xterra C18 30X100 mm S5, 30% to 85% Solvent B/A for 25 min gradient,

hold time 5 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined extracts were dried (MgSO₄) and concentrated to afford 360 mg (90%) of (1*S*,4*R*,6*R*,13*S*,17*R*)-13-tert-Butoxycarbonylamino-17-(7-methoxy-2-phenyl-quinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadecane-4-carboxylic acid: ¹H NMR (methanol-d₄) δ 1.10 (s, 9 H), 1.34 (m, 12 H), 1.68 (m, 1 H), 1.84 (m, 1 H), 2.05 (s, 1 H), 2.58 (m, 1 H), 2.71 (m, 1 H) 3.94 (s, 3 H), 4.05 (dd, *J*=11.4, 2.9 Hz, 1 H), 4.16 (dd, *J*=11.8, 2.9 Hz, 1 H), 4.67 (t, *J*=8.1 Hz, 1 H), 4.90 (m, 1 H), 5.61 (s, 1 H), 7.10 (dd, *J*=9, 2 Hz, 1 H), 7.29 (s, 1 H), 7.38 (d, *J*=2 Hz, 1 H), 7.55 (m, 3 H), 8.03 (d, *J*=6.7 Hz, 2 H), 8.13 (d, *J*=9 Hz, 1 H). LC-MS *m/e* 687 (retention time: 2.84, method D, except gradient time increased from 2 to 4 min). LC-MS *m/e* 687 (retention time: 2.84, method D, except gradient time increased from 2 to 4 min).

Step ii/12j-Method B Preparation of Compound 13, Example 12, Preparation of (1*S*,4*R*,6*R*,13*S*,17*R*)-[4-Cyclopropanesulfonylamino-carbonyl-17-(7-methoxy-2-phenyl-quinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-13-yl]-carbamic acid tert-butyl ester, Compound 13, Example 12.



Compound 13, Example 12.


Method B-Step ii/Step 12j) Following the procedural and purification conditions of Step 1j, except
 5 preparative HPLC/Extractive workup was used instead of normal phase chromatography, 260 mg (0.38 mmol) of (1*S*,4*R*,6*R*,13*S*,17*R*)-13-*tert*-Butoxycarbonyl-amino-17-(7-methoxy-2-phenyl-quinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0.4,6]octa-decane-4-carboxylic acid
 10 was converted to 192 mg (64%) (1*S*,4*R*,6*R*,13*S*,17*R*)-[4-Cyclopropanesulfonylamino-carbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0.4,6]octadec-13-yl]carbamic acid *tert*-butyl ester, Compound 13, Example 12 identical
 15 spectroscopically to that prepared in Method A.
 Representative Preparative HPLC /Extractive Workup
 Conditions: Xterra C18 30X100 mm S5, 35% to 100% Solvent B/A for 25 min gradient, hold time 5 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is
 20 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N

aqueous HCl solution, and repeatedly extracted with EtOAc. The combined extracts were dried (MgSO₄) and concentrated to afford the aforementioned amount of Compound 13.

5

Method B-Stepiii/Step 13k: Preparation of Compound 14,
Example 13, (1S,4R,6R,13S,17R)-Cyclopropanesulfonic acid
[13-amino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-
dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]-octadecane-4-
carbonyl]amide dihydrochloride

10




Compound 14, Example 13

Method B-Step iii/Step 13k) A solution of 2.5 mL (10
 15 mmol) of 4N HCl/dioxane was added to 161 mg (0.204 mmol)
 of (1S,4R,6R,13S)-[4-Cyclopropanesulfonylamino-carbonyl-
 17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-
 diazatricyclo[13.3.0.0^{4,6}]-octadec-13-yl]carbamic acid
 tert-butyl ester, the mixture stirred 3h, concentrated
 20 and dried (40 °C; 20 torr; 24h) to afford 156 mg (~100%)
 of Compound 14, Example 13, (1S,4R,6R,13S,17R)-
 Cyclopropanesulfonic acid [13-amino-17-(7-methoxy-2-
 phenylquinolin-4-yloxy)-2,14-dioxo-3,15-


diazatricyclo[13.3.0.0^{4,6}]octadecane-4-carbonyl]amide dihydrochloride as a white solid: ¹H NMR (500 MHz, CD₃OD) δ 1.03-1.26 (m, 3 H), 1.26-1.75 (m, 14 H), 1.84 (s, 1 H), 2.02 (m, 1 H), 2.75¹ (m, 1 H), 2.85 (m, 1 H), 2.96 (m, 1 H), 4.05 (s, 3 H), 4.23 (m, 1 H), 4.31 (s, 1 H), 4.46 (d, *J* = 11.29 Hz, 1 H), 5.95 (s, 1 H), 7.44 (d, *J* = 8.85 Hz, 1 H), 7.57 (s, 1 H), 7.61 (s, 1 H), 7.75 (dd, *J* = 8.55, 7.32 Hz, 3 H), 8.13 (d, *J* = 6.71 Hz, 1 H), 8.38 (d, *J* = 9.16 Hz, 1 H). LC-MS *m/e* 690 (retention time: 2.10, method D, except gradient time increased from 2 to 4 min).

Examples 14, 15

Preparation of Compounds 15 and 16



Compound 15, Example 14 - R is Boc
Compound 16, Example 15 - R is H (Bis HCl Salt)




Compound 15, Example 14- R is BOC
Compound 16, Example 15- R is H

Step i of Method B) To a solution of 470 mg (0.67 mmol) of (1*S*,4*R*,6*S*,14*S*,18*R*)-8-*cis*-14-*tert*-butoxycarbonylamino-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.^{4,6}]nona-dec-7-ene-4-carboxylic acid and 1.90 g (11.7 mmol) of potassium diazodicarboxylate (prepared as in *Org. React.* **1991**, 40, p.91) in 20 mL of MeOH, was added a solution of 1.34 mL (22.4 mmol) of glacial AcOH in 20 mL of MeOH slowly dropwise via syringe pump over 6.5 h. This procedure was repeated twice, but using a 12 h time interval instead of 6.5 h during which the reaction was stopped at approximately 90% completion. The mixture was concentrated in vacuo and then chromatographed over 20 g of SiO₂ (eluted with 5% MeOH/CH₂Cl₂). The resulting residue was dissolved in 18 mL of MeOH and injected nine times (9 x 2 mL injections) onto a preparative HPLC column: Xterra C18 30X100 mm S5, 30% to 75% Solvent B/A for 30 min gradient, hold time 5 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAC. The

combined extracts were dried (MgSO₄) and concentrated to afford 278 mg (59%) of (1*S*,4*R*,6*R*,14*S*,18*R*)-14-*tert*-butoxycarbonylamino-18-(7-methoxy-2-phenyl-quinolin-4-yloxy)-2,15-dioxo-3,16-diazatricycl

5 o[14.3.0.0^{4,6}]nonadecane-4-carboxylic acid: ¹H NMR (500 MHz, methanol-d₄) δ 1.23 (s, 9 H), 1.12-1.49 (m, 11 H), 1.50-1.67 (m, 4 H), 1.71-1.86 (m, 2 H), 2.54-2.60 (m, 1 H), 2.65-2.72 (m, 1 H), 3.94 (s, 3 H), 4.05 (dd, *J*=11.29, 3.36 Hz, 1 H), 4.28-4.33 (m, 1 H), 4.65 (d, *J*=11.29 Hz, 1
10 H), 4.70 (t, *J*=8.09 Hz, 1 H), 5.55-5.61 (m, 1 H), 7.07-7.15 (m, 1 H), 7.28 (s, 1 H), 7.38 (d, *J*=1.83 Hz, 1 H), 7.50-7.59 (m, 3 H), 8.04 (d, *J*=7.02 Hz, 2 H), 8.15 (d, *J*=9.16 Hz, 1 H). LC-MS *m/e* 701 (retention time: 3.01, method D, except gradient time increased from 2 to 4
15 min).

Step ii/14j-Method B Preparation of Compound 15, Example 14, Preparation of (1*S*,4*R*,6*R*,14*S*,18*R*)-[4-Cyclopropanesulfonylamino-
20 carbonyl-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo-
[14.3.0.0^{4,6}]nonadec-14-yl]carbamic acid *tert*-butyl ester.



Compound 15, Example 14


Method B-Step ii/Step 14j) Following the analogous procedural and purification conditions of Steps 1j and 12j, respectively, 100 mg (0.143 mmol) of

5 (1*S*,4*R*,6*R*,14*S*,18*R*)-14-*tert*-Butoxycarbonylamino-18-(7-methoxy-2-phenyl-quinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadecane-4-carboxylic acid was converted to 108 mg (93%) (1*S*,4*R*,6*R*,14*S*)-[4-Cyclopropanesulfonylamino-18-(7-methoxy-2-phenyl-

10 quinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-14-yl]carbamic acid *tert*-butyl ester, Compound 15, Example 14: ¹H NMR (500 MHz, methanol-*d*₄) δ 1.20 (m, 9 H) 0.83-1.83 (m, 21 H), 2.48 (s, 1 H), 2.73 (s, 1 H), 2.95 (s, 1 H), 3.94 (s, 3 H), 4.04 (m, 1 H),

15 4.24 (m, 1 H), 4.67 (m, 2 H), 5.60 (s, 1 H), 7.12 (m, *J*=8.85 Hz, 1 H), 7.30 (s, 1 H), 7.38 (s, 1 H), 7.57 (m, *J*=6.10 Hz, 3 H), 8.03 (m, *J*=5.80 Hz, 2 H), 8.16 (d, *J*=8.85 Hz, 1 H). LC-MS *m/e* 804 (retention time: 1.73, method D.

Method B-Stepiii/Step 15k: Preparation of Compound 16,
 Example 15, (1*S*,4*R*,6*R*,14*S*,18*R*)-Cyclopropanesulfonic acid
 [14-amino-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-
 dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadecane-4-
 5 carbonyl]amide dihydrochloride




Compound 16, Example 15

Method B-Step iii/Step 15k) A solution of 2.5 mL (10 mmol) of 4N HCl/dioxane was added to 101 mg (0.126 mmol) of (1*S*,4*R*,6*R*,14*S*,18*R*)-[4-Cyclopropanesulfonyl-amino-carbonyl-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nona-dec-14-yl]carbamic acid tert-butyl ester (Compound 15 from Example 14), the mixture stirred 3h, concentrated and dried (40 °C; 20 torr; 24h) to afford 98 mg (~100%) of Compound 16, Example 15, (1*S*,4*R*,6*R*,14*S*,18*R*)-Cyclopropanesulfonic acid [14-amino-18-(7-methoxy-2-phenylquin-olin-4-yloxy)-2,15-dioxo-3,16-diazatri-cyclo[14.3.0.0^{4,6}]nonadecane-4-carbonyl]amide
 20 dihydrochloride as a white solid: ¹H NMR (500 MHz, CD₃OD) δ 1.05-1.77 (m, 19 H), 1.88 (m, 2 H), 2.60 (m, 1 H), 2.90 (m, 1 H), 2.96 (m, 1 H), 4.05 (s, 3 H), 4.21 (m, 1 H), 4.41 (m, 1 H), 4.46 (s, 1 H), 5.93 (s, 1 H), 7.44 (d,

$J=8.55$ Hz, 1 H), 7.58 (s, 1 H), 7.62 (s, 1 H), 7.74 (m, 3 H), 8.10 (m, $J=7.02$ Hz, 2 H), 8.33 (d, $J=9.16$ Hz, 1 H).
LC-MS m/e 704 (retention time: 2.24, method D, except gradient time increased from 2 to 4 min).

5

Step 16j: Preparation of Compound 17, Example 16
(1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-[4-Cyclobutanesulfonylamino-
carbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-
dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-7-en-13-
10 yl]carbamic acid *tert*-butyl ester




Compound 17, Example 16

Step 16j) Following the analogous procedural
15 conditions of Steps 1j except cyclobutylsulfonamide was
substituted for cyclopropylsulfonamide as well using the
following preparative HPLC/Extractive Workup
Purification: HPLC Column Xterra C18 30X100 mm S5, 35% to
85% Solvent B/A for 23 min gradient, hold time 4 min;
20 where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent
B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 30
ml/min. The pH of the combined fractions was adjusted to
neutral using 6N aqueous NaOH solution and the MeOH
removed in vacuo. The solution was adjusted to pH 4
25 using 3N aqueous HCl solution, and repeatedly extracted

with EtOAC. The combined extracts were dried (MgSO₄) and concentrated to give the resulting conversion: 52 mg (0.076 mmol) of (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-13-*tert*-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octa-dec-7-ene-4-carboxylic acid was converted to 12 mg (20%) of (1*S*,4*R*,6*S*,13*S*,17*R*)-7-*cis*-[4-Cyclobutanesulfonylamino-carbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-7-en-13-yl]carbamic acid *tert*-butyl ester: ¹H NMR (500 MHz, CD₃OD) δ 1.18 (s, 9 H), 1.23-1.56 (m, 11 H), 1.76 (m, 1 H), 1.98-2.13 (m, 3 H), 2.15-2.34 (m, 2 H), 2.48 (m, 1 H), 2.62 (m, 1 H), 2.74 (m, 1 H), 3.98 (m, 3 H), 4.29 (m, 3 H), 4.66 (m, 1 H), 5.41 (m, 1 H), 5.64 (m, 2 H), 7.12 (d, *J*=8.85 Hz, 1 H), 7.29 (s, 1 H), 7.42 (m, 1 H), 7.56 (m, 3 H), 8.10 (m, 3 H). LC-MS (Method D-retention time: 1.61), MS *m/z* 802 (M⁺+1).


Step 17j-Preparation of Compound 18, Example 17,
20 Preparation of (1*S*,4*R*,6*R*,13*S*,17*R*)-[4-Cyclobutanesulfonylamino-carbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octa-dec-13-yl]carbamic acid *tert*-butyl ester.



Compound 18, Example 17

Step 17j) Following the procedural and purification conditions of Step 1j, except preparative HPLC/Extractive workup was used instead of normal phase chromatography, 56 mg (0.082 mmol) of (1*S*,4*R*,6*R*,13*S*,17*R*)-13-*tert*-butoxycarbonylamino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadecane-4-carboxylic acid was converted to 12 mg (20%) of (1*S*,4*R*,6*R*,13*S*,17*R*)-[4-Cyclobutanesulfonylamino-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octadec-13-yl]carbamic acid *tert*-butyl ester, Compound 18, Example 17. Representative Preparative HPLC/Extractive Workup Conditions: Xterra C18 30X100 mm S5, 35% to 85% Solvent B/A for 25 min gradient, hold time 5 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAC. The combined extracts were dried (MgSO₄) and concentrated to afford the aforementioned amount of Compound 18: ¹H NMR (500 MHz, CD₃OD) δ 1.07 (s, 9 H), 1.22-1.62 (m, 15 H), 1.63-1.78 (m, 2 H), 2.04 (m, 1 H), 2.52 (m, 4 H), 2.87 (m, 1 H), 4.04 (s, 3 H), 4.14 (m, 2 H), 4.36 (m, 1 H), 4.78 (m, 1 H), 5.07 (m, 1 H), 5.82 (s, 1 H), 7.32 (d, *J*=9.16 Hz, 1 H), 7.51 (s, 1 H), 7.56 (d, *J*=8.55 Hz, 1 H), 7.72 (m, 3 H), 8.08 (m, 2 H), 8.32 (m, 1 H). LC-MS (Method D-retention time: 1.68), MS *m/z* 804 (M⁺+1).


Step 18j-Preparation of Compound 19, Example 18,
 Preparation of (1*S*,4*R*,6*R*,14*S*,18*R*)-[4-Cyclobutanesulf-
 onylaminocarbonyl-18-(7-methoxy-2-phenylquinolin-4-
 yloxy)-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nona-
 5 dec-14-yl]carbamic acid tert-butyl ester.



Compound 19, Example 18

Step 18j) Following the analogous procedural and
 10 purification conditions of Steps 1j and 17j,
 respectively, 50 mg (0.071 mmol) of (1*S*,4*R*,6*R*,14*S*, 18*R*)-
 14-*tert*-Butoxycarbonylamino-18-(7-methoxy-2-phenyl-
 quinolin-4-yloxy)-2,15-dioxo-3,16-
 diazatricyclo[14.3.0.0^{4,6}]nonadecane-4-carboxylic acid was
 15 converted to 22 mg (38%) of (1*S*,4*R*,6*R*,14*S*,18*R*)-[4-
 Cyclobutanesulfonylaminocarbonyl-18-(7-methoxy-2-phenyl-
 quinolin-4-yloxy)-2,15-dioxo-3,16-diazatri-
 cyclo[14.3.0.0^{4,6}]nonadec-14-yl]carbamic acid *tert*-butyl
 ester, Compound 19, Example 18: ¹H NMR (500 MHz, Solvent
 20 methanol-*d*₄) δ 1.21 (s, 9 H), 1.14-1.88 (m, 20 H), 2.35
 (m, 2 H), 2.53 (m, 3 H), 2.83 (dd, *J*=13.4, 6.4 Hz, 1 H),
 4.01 (s, 3 H), 4.12 (d, *J*=11.3 Hz, 1 H), 4.27 (m, 1 H),
 4.36 (m, 1 H), 4.70 (m, 1 H), 5.73 (m, 1 H), 7.21 (d,

$J=9.2$ Hz, 1 H), 7.41 (m, 1 H), 7.46 (m, 1 H), 7.64 (m, 3 H), 8.08 (m, 2 H), 8.25 (d, $J=9.2$ Hz, 1 H). LC-MS m/e 818 (retention time: 1.75, method D).



5

Compound 10, Example 10


Compound 20, Example 19

Step 191: Preparation of Compound 20, Example 19

Preparation of (1S,4R,6S,14S,18R)-7-cis-Cyclopropane-sulfonic acid [14-(3-tert-butylureido)-18-(7-methoxy-2-phenyl-quinolin-4-yloxy)-2,15-dioxo-3,16-diazatri-
 10 cyclo[14.3.0.0^{4,6}]nona-dec-7-ene-4-carbonyl]amide

Step 191) To a suspension of 59 mg (0.0762 mmol) of compound 10 and 53.4 μ L (0.381 mmol) of Et_3N in 2 mL of CH_2Cl_2 , was added 24 μ L (0.208 mmol) of tert-butyl isocyanate (Aldrich). The mixture was stirred 18h, concentrated in vacuo and dissolved up in 2 mL of MeOH. This solution was injected onto a preparative HPLC using the following conditions: Column Xterra C18 30X100 mm S5, 20 37% to 85% Solvent B/A for 30 min gradient, hold time 5 min; where Solvent A is 10% MeOH/90% H_2O with 0.1%TFA, Solvent B is 90%MeOH/10% H_2O with 0.1%TFA and flow rate is 35 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and 25 the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly


extracted with EtOAC. The combined extracts were dried (MgSO₄) and concentrated to afford 30 mg (49%) of Compound 20, Example 19: ¹H NMR (500 MHz, methanol-d₄) δ 1.03 (s, 9 H) 1.11 (m, 3 H) 1.30 (m, 5 H) 1.49 (m, 5 H) 1.65 (m, 1 H) 1.78 (m, 2 H) 1.91 (m, 1 H) 2.41 (m, 1 H) 2.69 (dd, J=14.04, 4.58 Hz, 2 H) 2.85 (m, 1 H) 2.94 (m, 1 H) 4.07 (s, 3 H) 4.13 (dd, J=12.2, 3.4 Hz, 1 H) 4.25 (d, J=10.4 Hz, 1 H) 4.75 (m, 1 H) 4.98 (d, J=12.2 Hz, 1 H) 5.11 (m, 1 H) 5.74 (m, 1 H) 5.91 (s, 1 H) 7.40 (dd, J=9.3, 2.4 Hz, 1 H) 7.55 (m, 1 H) 7.67 (s, 1 H) 7.77 (m, 3 H) 8.10 (d, J=7.6 Hz, 1 H) 8.42 (d, J=9.3 Hz, 1 H). LC-MS m/e 801 (retention time: 2.88, method D, except gradient time increased from 2 to 4 min).



Step 20I: Preparation of Compound 21, Example 20
Preparation of (1S,4R,6S,14S,18R)-7-cis-[4-
Cyclopropanesulfonylaminocarbonyl-18-(7-methoxy-2-
 20 phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo-
[14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid 2-
fluoroethyl ester

Step 20I) To a suspension of 59 mg (0.0762 mmol) of
 25 compound 10 and 53.4 μL (0.381 mmol) of Et₃N in 2 mL of
 CH₂Cl₂, was added 20 μL (0.208 mmol) of 2-fluoroethyl

chloroformate (Aldrich). The mixture was stirred 18h, concentrated in vacuo and dissolved up in 2 mL of MeOH. This solution was injected onto a preparative HPLC using the following conditions: Column Xterra C18 30X100 mm S5, 30% to 80% Solvent B/A for 30 min gradient, hold time 2 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 35 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined extracts were dried (MgSO₄) and concentrated to afford 38 mg (64%) of Compound 21, Example 20: ¹H NMR (500 MHz, methanol-d₄) δ 1.05 (m, 3 H) 1.25-1.89 (m, 8 H) 1.62 (dd, J=9.5, 5.49 Hz, 1 H) 1.74 (dd, J=8.1, 5.34 Hz, 1 H) 1.84 (m, 2 H) 2.39 (m, 1 H) 2.66 (m, 2 H) 2.83 (m, 1 H) 2.91 (m, 1 H) 3.87 (m, 2 H) 4.06 (s, 3 H) 4.09 (m, 1 H) 4.22 (m, 1 H) 4.30 (m, 1 H) 4.40 (m, 1 H) 4.71 (m, 1 H) 5.08 (m, 1 H) 5.70 (m, 1 H) 5.91 (s, 1 H) 7.41 (m, 1 H) 7.53 (s, 1 H) 7.65 (s, 1 H) 7.73 (m, 3 H) 8.07 (d, J=7.6 Hz, 2 H) 8.39 (d, J=9.2 Hz, 1 H). LC-MS m/e 792 (retention time: 2.66, method D, except gradient time increased from 2 to 4 min).




Compound 16, Example 15

Compound 22, Example 21


Step 211: Preparation of Compound 22, Example 21

Preparation of (1S,4R,6R,14S,18R)-[4-Cyclopropane-sulfonylaminocarbonyl-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-14-yl]carbamic acid 2,2-dimethylpropyl ester

Step 211) To a suspension of 45 mg (0.058 mmol) of compound 16 and 42 μ L (0.29 mmol) of Et₃N in 2 mL of CH₂Cl₂, was added 26 μ L (0.174 mmol) of neopentyl chloroformate (Aldrich). The mixture was stirred 18h, concentrated in vacuo and dissolved up in 2 mL of MeOH. This solution was injected onto a preparative HPLC using the following conditions: Column Xterra C18 30X100 mm S5, 30% to 80% Solvent B/A for 29 min gradient, hold time 2 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 35 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined extracts were dried (MgSO₄) and concentrated to afford 35 mg (74%) of Compound 22, Example 21: ¹H NMR (500 MHz, methanol-d₄) δ 0.81 (s, 9 H), 1.10-1.25 (m, 4 H), 1.27-1.79 (m, 16 H), 1.86 (m, 1 H) 2.60 (m, 1 H) 2.90 (m, 1 H), 3.01 (m, 1 H), 3.18 (d, J=10.1 Hz, 1 H), 3.35 (m, 1 H), 4.07 (m, 3 H), 4.13 (m, 1 H), 4.26 (m, 1 H), 4.76 (m, 1 H), 4.90 (m, 1 H), 5.93 (m, 1 H), 7.43 (m, 1 H), 7.55 (m, 1 H), 7.68 (m, 1 H), 7.79 (m, 3 H), 8.12 (m, 2 H), 8.39 (d, J=9.2 Hz, 1 H). LC-MS m/e 818 (retention time: 3.21, method D, except gradient time increased from 2 to 4 min).

General Procedure For Preparation of chloroformates

This procedure was used for the preparation of non-commercially available chloroformates. To a solution of 5.96 g (67.6 mmol) of commercially available reagents (S)-3-hydroxytetrahydrofuran and pyridine (5.8 mL; 72 mmol) in THF (150 mL) cooled to 0 °C was added a 1.93 M solution of phosgene in toluene (48 mL, 92.6 mmol over 10 min under argon. The resulting solution was allowed to warm to rt over 2 h, the resulting solid filtered, and the mother liquor carefully concentrated in vacuo at room temperature until theoretical mass was obtained. The resulting residue was dissolved in 100 mL of THF to prepare a 0.68M stock solution of 3(S)-oxo-tetrahydrofuran chloroformate that could be stored in the freezer until use. In analogous fashion, other commercially available alcohols could be converted to 0.68M stock solutions of the corresponding chloroformates.



Step 221: Preparation of Compound 23, Example 22Preparation of (1S,4R,6R,14S,18R)-[4-Cyclopropanesulfonyl-aminocarbonyl-18-(7-methoxy-2-phenylquinolin-4-yloxy)-2,15-dioxo-3,16-

5 diazatricyclo[14.3.0.04,6]nonadec-14-yl]carbamic acid
tetrahydro[4H]pyran-4-yl ester

Step 221) To a suspension of 45 mg (0.058 mmol) of compound 16 and 42 μ L (0.29 mmol) of Et_3N in 2 mL of

10 CH_2Cl_2 , was added 217 μ L (0.174 mmol) of ~0.8M tetrahydro[4H]pyran-4-yl chloroformate (prepared from Aldrich quality tetrahydro[4H]pyran-4-ol using the same method as in the preparation of 3(S)-oxo-tetrahydrofuran chloroformate) in CH_2Cl_2 . The mixture was stirred 18h,


15 concentrated in vacuo and dissolved up in 2 mL of MeOH. This solution was injected onto a preparative HPLC using the following conditions: Column Xterra C18 30X100 mm S5, 30% to 80% Solvent B/A for 29 min gradient, hold time 2 min; where Solvent A is 10% MeOH/90% H_2O with 0.1%TFA,

20 Solvent B is 90%MeOH/10% H_2O with 0.1%TFA and flow rate is 35 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly

25 extracted with EtOAC. The combined extracts were dried (MgSO_4), concentrated, and rechromatographed over a Biotage 12M column (eluted with 0 to 8% MeOH/ CH_2Cl_2) to afford 14.2 mg (30%) of Compound 23, Example 22: ^1H NMR (500 MHz, CDCl_3) δ 1.24 (m, 25 H), 2.48 (m, 1 H) 2.64 (m, 1 H), 2.93 (m, 1 H), 3.44 (m, 2 H), 3.83 (m, 2 H), 3.93 (s, 3 H), 4.05 (m, 1 H), 4.39 (d, $J=11.3$ Hz, 1 H), 4.45 (t, $J=6.9$ Hz, 1 H), 4.65 (m, 2 H), 5.36 (m, 2 H,) 6.95 (s, 1 H), 7.02 (dd, $J=9.0, 2.3$ Hz, 1 H), 7.49 (m, 4 H),

30

7.93 (d, $J=8.8$ Hz, 1 H), 8.03 (d, $J=7.3$ Hz, 2 H), 10.26 (s, 1 H). LC-MS m/e 832 (retention time: 2.90, method D, except gradient time increased from 2 to 4 min).



5

Step 231: Preparation of Compound 24, Example 23

Preparation of (1*S*,4*R*,6*R*,14*S*,17*R*)-*N*-[4-Cyclopropane-
sulfonylaminocarbonyl-17-(7-methoxy-2-phenylquinolin-4-
yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.⁰_{4,6}]-octadec-
13-yl]-2-cyclopropylacetamide


10

Step 231) To a suspension of 50 mg (0.065 mmol) of compound 14, 14 mg (0.139 mmol) of cyclopropyl acetic acid and 97 μ L of DIPEA in 2 mL of CH_2Cl_2 , was added 53 mg (0.139 mmol) of HATU. The mixture was stirred 18h, concentrated in vacuo and dissolved up in 2 mL of MeOH. This solution was injected onto a preparative HPLC using the following conditions: Column Xterra C18 30X100 mm S5, 30% to 80% Solvent B/A for 30 min gradient, hold time 2 min; where Solvent A is 10% MeOH/90% H_2O with 0.1%TFA, Solvent B is 90%MeOH/10% H_2O with 0.1%TFA and flow rate is 35 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly

20

25

extracted with EtOAC. The combined extracts were dried (MgSO₄) and concentrated to afford 20 mg (40%) of Compound 24, Example 23: ¹H NMR (500 MHz, methanol-d₄) δ -0.01 (m, 2 H), 0.30 (m, 2 H), 0.66 (m, 1 H), 0.88 (m, 1 H), 1.08 (m, 3 H), 1.24-1.68 (m, 13 H), 1.75 (m, 1 H), 1.91 (m, 3 H), 2.63 (m, 1 H), 2.81 (s, 1 H), 2.98 (m, 1 H), 4.02 (s, 3 H), 4.15 (m, 1 H), 4.41 (m, 1 H), 4.73 (t, J=8.1 Hz, 1 H), 4.94 (d, J=11.6 Hz, 1 H), 5.86 (s, 1 H), 7.31 (dd, J=9.2, 2.2 Hz, 1 H), 7.50 (d, J=2.2 Hz, 1 H), 7.58 (s, 1 H), 7.69 (m, 3 H), 8.08 (m, 2 H), 8.27 (d, J=9.2 Hz, 1 H). LC-MS m/e 772 (retention time: 2.77, method D, except gradient time increased from 2 to 4 min).




Step 241: Preparation of Compound 25, Example 24

Preparation of (1S,4R,6R,14S,17R)-[4-Cyclopropanesulfonylaminocarbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]octa-dec-13-yl]carbamic acid cyclopentyl ester

Step 241) To a suspension of 50 mg (0.065 mmol) of compound 14, and 47 μL (0.32 mmol) of Et₃N in 2 mL of CH₂Cl₂, was added 300 μL (0.20 mmol) of ~0.67M cyclopentyl chloroformate (prepared from Aldrich quality

cyclopentanol using the same method as in the preparation of 3(S)-oxo-tetrahydrofuran chloroformate) in CH_2Cl_2 . The mixture was stirred 18h, concentrated in vacuo and dissolved up in 2 mL of MeOH. This solution was injected
5 onto a preparative HPLC using the following conditions: Column Xterra C18 30X100 mm S5, 30% to 80% Solvent B/A for 30 min gradient, hold time 2 min; where Solvent A is 10% MeOH/90% H_2O with 0.1%TFA, Solvent B is 90%MeOH/10% H_2O with 0.1%TFA and flow rate is 35 ml/min). The pH of
10 the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined extracts were dried (MgSO_4) and concentrated to
15 afford 26 mg (50%) of Compound 25, Example 24: ^1H NMR (500 MHz, methanol- d_4) δ 1.02-1.17 (m, 3 H), 1.20-1.67 (m, 22 H), 1.72 (m, 1 H), 1.88 (m, 1 H), 2.63 (m, 1 H), 2.86 (m, 1 H), 2.98 (m, 1 H), 4.03 (s, 3 H), 4.08 (m, 1 H), 4.16 (m, 1 H), 4.28 (m, 1 H), 4.78 (m, 1 H), 5.02 (d, $J=11.0$ Hz, 1 H), 5.89 (m, 1 H), 7.38 (m, 1 H), 7.51 (d, $J=2.1$ Hz, 1 H), 7.64 (s, 1 H), 7.73 (m, 3 H), 8.06 (d, $J=7.6$ Hz, 2 H), 8.34 (d, $J=9.5$ Hz, 1 H). LC-MS m/e 802 (retention time: 3.00, method D, except gradient time increased from 2 to 4 min).

25




Step 251: Preparation of Compound 26, Example 25

Preparation of (1*S*,4*R*,6*R*,14*S*,17*R*,3'*S*)-[4-Cyclopropane-sulfonyl-aminocarbonyl-17-(7-methoxy-2-phenylquinolin-4-yloxy)-2,14-dioxo-3,15-diazatricyclo[13.3.0.0^{4,6}]-octadec-13-yl]-carbamic acid tetrahydrofuran-3'-yl ester

Step 251) To a suspension of 50 mg (0.065 mmol) of compound 14, and 47 μ L (0.32 mmol) of Et₃N in 2 mL of CH₂Cl₂, was added 300 μ L (0.20 mmol) of ~0.67M 3(*S*)-oxo-tetrahydrofuran chloroformate (prepared as described above) in CH₂Cl₂. The mixture was stirred 18h, concentrated in vacuo and dissolved up in 2 mL of MeOH. This solution was injected onto a preparative HPLC using the following conditions: Column Xterra C18 30X100 mm S5, 30% to 80% Solvent B/A for 30 min gradient, hold time 2 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 35 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAC. The combined extracts were dried (MgSO₄) and concentrated and chromatographed over a Biotage 12M column (eluted with 0% to 8% MeOH/CH₂Cl₂) to afford 10 mg (25%) of Compound 26, Example 25: ¹H NMR (500 MHz, CDCl₃) δ 1.20 (m, 18 H), 1.73 (m, 2 H), 1.93 (m, 2 H), 2.59 (m, 1 H), 2.69 (m, 1 H), 2.95 (m, 1 H), 3.74 (m, 3 H), 3.93 (s, 3 H), 4.07 (m, 1 H), 4.31 (m, 1 H), 4.65 (t, *J*=7.48 Hz, 1 H), 4.71 (d, *J*=11.3 Hz, 1 H), 4.96 (s, 1 H), 5.37 (d, *J*=6.7 Hz, 1 H), 5.43 (s, 1 H), 7.02 (m, 3 H), 7.50 (m, 3 H), 8.07 (d, *J*=7.3 Hz, 2 H) 10.83 (s, 1 H). LC-MS *m/e* 804 (retention time: 2.69,

method D, except gradient time increased from 2 to 4 min).

The following steps (A)-(G) show the preparation of the intermediate, the product of step (G).




Step A) To a solution of 4.0 g (46.5 mmol) of 3-methyl-2-butanone (Aldrich) in 50 mL of MeOH was added dropwise a solution of 2.4 mL (46.5 mmol) of bromine over 40 min. The mixture was stirred 1.5h, diluted with 300 mL of pentane, washed with sat. aqueous NaHCO₃, dried (MgSO₄) and concentrated to afford 5.81 g of impure 1-Bromo-3-methyl-butan-2-one which was taken directly into the step B.

Step B) A neat solution of 5.58g (34 mmol) of 1-Bromo-3-methyl-butan-2-one and 4.50g (34 mmol) of ethyl thioamate (Aldrich) was heated at 70 °C over 18h and then cooled to room temperature. The mixture was partitioned between sat. aqueous NaHCO₃ and EtOAc, the EtOAc layer dried (MgSO₄), concentrated and chromatographed over SiO₂ (eluted with 2% to 40% EtOAc/hexanes) to afford 3.4 g (48% overall) of 4-isopropylthiazole-2-carboxylic acid ethyl ester as an oil: ¹H NMR (500 MHz, CDCl₃) δ 1.32 (d, J=7 Hz, 6 H), 1.42

(t, $J=7.2$ Hz, 3 H), 3.23 (m, 1 H), 4.46 (q, $J=7.2$ Hz, 2 H), 7.18 (s, 1 H).

Step C) To a solution of 3.12 g (15.7 mmol) of 4-isopropylthiazole-2-carboxylic acid ethyl ester in 32 mL of 75% THF/MeOH, was added 110 mg (31.3 mmol) of LiOH in 8 mL of H₂O. The mixture was stirred overnight, the solution adjusted to pH 5 using 1N aqueous HCl solution and concentrated in vacuo to afford 4-isopropylthiazole-2-carboxylic acid as a white solid (2.97g including salts) which was used directly in Step E: ¹H NMR (500 MHz, methanol-d₄) δ 1.29 (d, $J=6.7$ Hz, 6H), 3.20 (m, 1 H), 7.39 (m, 1 H).



15

Step D) To a solution of 21.8g (172 mmol) of *m*-anisidine in 800 mL of CH₂Cl₂ cooled to -50 °C, was added 172 mL (172 mmol) of 1M BCl₃ in heptane over 20 min to produce an amber colored mixture which was stirred 1 h. To this mixture was added 12.2 mL (172 mmol) of AcCl and 22.9g (172 mmol) of AlCl₃. The mixture was allowed to warm to room temperature overnight, poured into ice water, the solution adjusted to pH 9 using 10% aqueous NaOH and was extracted repeatedly using EtOAc. The combined organic extracts were washed with brine, dried (MgSO₄), concentrated and the resulting solid recrystallized from

the minimal amount of MeOH in CH₂Cl₂/hexanes (50%) to afford 13.4 g (47%) of slightly impure 2-amino-4-methoxybenzophenone as a crystalline solid. The solid could be taken directly into the next reaction or
5 purified to absolute purity over SiO₂ (eluted with 0% to 14% MeOH/CH₂Cl₂): ¹H NMR (500 MHz, CDCl₃) δ 2.50 (s, 3 H), 3.79 (s, 3 H), 6.05 (d, J=2.4 Hz, 1 H), 6.21 (dd, J=8.8, 2.4 Hz, 1 H), 6.38 (m, 2 H), 7.62 (d, J=8.8 Hz, 1 H).

10 **Step E)** To a suspension of 2.59g (15.7 mmol) of 2-amino-4-methoxybenzophenone (product of step D) and 2.68g (15.7 mmol) of 4-isopropylthiazole-2-carboxylic acid (product of step C) in 75 mL of pyridine cooled to -30 °C, was added 1.93 mL (23.5 mmol) of POCl₃ slowly
15 dropwise over 5 min. The mixture was stirred 3 h, warmed to room temperature and was stirred overnight. The reaction mixture was poured into ice water, and extracted several times with EtOAc. The combined EtOAc extracts were dried (MgSO₄), concentrated and chromatographed over
20 SiO₂ (eluted with 0% to 15% MeOH/EtOAc) to afford 2.57g (51%) of 4-Methylthiazole-2-carboxylic acid (2-acetyl-5-methoxyphenyl)amide as a yellow solid: ¹H NMR (500 MHz, CDCl₃) δ 1.41 (d, J=6.7 Hz, 6 H), 2.64 (s, 3 H), 3.24 (m, 1 H), 3.91 (s, 3 H), 6.67 (dd, J=9, 2.5 Hz, 1 H), 7.18 (s, 1 H), 7.86 (d, J=9 Hz, 1 H), 8.56 (d, 25 J=2.5 Hz, 1 H), 13.48 (s, 1 H).

Step F) To a solution of 2.5g (7.85 mmol) of 4-methylthiazole-2-carboxylic acid (2-acetyl-5-methoxyphenyl)amide (product of step E) in 50 mL of THF,
30 was added 19 mL (19 mmol) of 1M KOtBu in THF. The mixture was heated to 70 °C for 3h, cooled to rt and stirred overnight. The mixture was concentrated, cold

water added to form a suspension. The mixture was then acidified to pH 4, filtered and dried. The resulting solid was chromatographed over SiO₂ (eluted with 0% to 25% MeOH in CH₂Cl₂) to afford 1.31g (56%) of

5 2-(4-Isopropylthiazol-2-yl)-7-methoxyquinolin-4-ol as a beige solid: ¹H NMR (300 MHz, DMSO-D₆) δ 1.32 (d, J=6.6 Hz, 6 H), 3.14 (m, 1 H), 3.89 (s, 3 H), 7.06 (s, 1 H), 7.38 (s, 1 H), 7.51 (s, 1 H), 7.99 (d, J=9.2 Hz, 1 H), 11.77 (m, 1 H). LC-MS m/e 301 (retention time: 1.53,

10 method D).

Step G) A suspension of 1.3g (4.3 mmol) of 2-(4-Isopropylthiazol-2-yl)-7-methoxyquinolin-4-ol, product of step F, in 60 mL of POCl₃ was heated to reflux for 2h.


15 The solvent was removed in vacuo, the residue diluted with ice cold water and the mixture adjusted to pH 9 while cooling to 0 °C. This aqueous solution was extracted several times with EtOAc. The combined EtOAc extracts were washed once with brine, pH 4 buffer, dried

20 (MgSO₄), and concentrated to afford 0.89g (64%) of 4-Chloro-2-(4-isopropylthiazol-2-yl)-7-methoxyquinoline as a yellow solid: ¹H NMR (500 MHz, CDCl₃) δ 1.38 (d, J=7 Hz, 6 H), 3.19 (m, 1 H), 3.98 (s, 3 H), 7.06 (s, 1 H), 7.26 (m, 1 H), 7.47 (d, J=2 Hz, 1 H), 8.10 (d, J=9 Hz, 1 H),

25 8.31 (s, 1 H). LC-MS m/e 319 (retention time: 2.20, method D).

Preparation of Example 26, (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-butoxycarbonylamino-18-hydroxy-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid using Steps 26i-26iv.


5



Example 26

Step 26i: Preparation of 1-(2(*S*)-*tert*-
Butoxycarbonylamino-non-8-enoyl)-4(*R*)-hydroxy-
pyrrolidine-2(*S*)-carboxylic acid methyl ester

10




A solution of 2(*S*)-*tert*-butoxycarbonylamino-8-
nonenoic acid (purchased from RSP Amino Acids) (3.5 g,
12.9 mmol) in 200 mL of DCM was treated sequentially with
4(*R*)-hydroxypyrrolidine-2(*S*)-carboxylic acid methyl ester
hydrochloride (2.15 g, 11.8 mmol), *N*-methyl morpholine
(4.25 mL, 38.6 mmol), and HATU (5.37 g, 14.1 mmol). The
reaction mixture was stirred at rt under N₂ for 3 days,
and then concentrated in vacuo. The residue was
partitioned between ethyl acetate and pH 4 buffer
(biphthalate). The organic phase was washed with sat. aq.

15

20

NaHCO₃, dried (MgSO₄), and concentrated in vacuo to give the crude product. Flash chromatography (50% ethyl acetate/hexane to 100% ethyl acetate) gave 4.7 g (~100%) of 1-(2(*S*)-tert-Butoxycarbonylamino-non-8-enoyl)-4(*R*)-hydroxy-pyrrolidine-2(*S*)-carboxylic acid methyl ester as a colorless oil: ¹H NMR (500 MHz, CD₃OD) δ 1.33-1.50 (m, 8 H), 1.46 (s, 9 H), 1.57 (m, 1 H), 1.72 (m, 1 H) 2.08 (m, 2 H), 2.28 (m, 1 H), 3.72 (s, 3 H,) 3.75-3.87 (m, 2 H), 4.36 (m, 1 H), 4.51 (bs, 1 H), 4.57 (t, *J*=8.2 Hz, 1 H), 4.95 (d, *J*=10.4 Hz, 1 H), 5.01 (m, 1 H), 5.83 (m, 1 H). LC-MS (Method J, retention time: 3.01 min), MS *m/z* 399 (*M*⁺+1).

Step 26ii: Preparation of 1-([1-(2(*S*)-tert-Butoxycarbonylamino-non-8-enoyl)-4(*R*)-hydroxy-pyrrolidine-2(*S*) carbonyl]-(1*R*)-amino}-2(*S*)-vinyl-cyclopropanecarboxylic acid ethyl ester



20


1-(2(*S*)-tert-Butoxycarbonylamino-non-8-enoyl)-4(*R*)-hydroxy-pyrrolidine-2(*S*)-carboxylic acid methyl ester (4.7 g, 11.8 mmol) was dissolved in THF (80 mL), methanol (20 mL), and water (40 mL). Powdered lithium hydroxide (5.6 g, 233 mmol) was added. The light yellow slurry was stirred at rt under N₂ for 16 h, and then concentrated in vacuo. The residue was partitioned between ether and water. The ether phase was discarded, and the

aqueous phase was treated with 1N HCl until the pH was 4. This acidic solution was extracted with EtOAc (3x). The combined EtOAc extracts were dried (MgSO₄) and concentrated in vacuo to give 4.36 g (96%) of 1-(2(S)-tert-butoxycarbonylamino-8-nonenoyl)-4(R)-hydroxy-pyrrolidine-2(S)-carboxylic acid as a white solid. This acid was then dissolved in 150 mL of DMF and (1R,2S)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride (2.61 g, 13.6 mmol), N-methyl morpholine (2.5 mL, 22.6 mmol), HATU (5.2 g, 13.7 mmol) was added. The reaction mixture was stirred at rt under N₂ for 16 h, and then concentrated in vacuo. The residue was partitioned between ethyl acetate and pH 4 buffer (bipthalate). The organic phase was washed with sat. aq. NaHCO₃, dried (MgSO₄), and concentrated in vacuo to give the crude product. Flash chromatography (60%-80% ethyl acetate/hexane) gave 6.0 g (98%) of 1-([1-(2(S)-tert-Butoxycarbonylamino-non-8-enoyl)-4(R)-hydroxy-pyrrolidine-2(S)carbonyl]-(1R)-amino)-2(S)-vinyl-cyclopropanecarboxylic acid ethyl ester as a white solid:

¹H NMR (500 MHz, CD₃OD) δ 1.25 (t, J=7.2 Hz, 3 H), 1.33-1.80 (m, 10 H), 1.46 (s, 9 H), 2.09 (m, 3 H), 2.25 (m, 2 H), 3.76 (m, 2 H), 4.14 (m, 2 H), 4.27 (dd, J=8.5, 5.2 Hz, 1 H), 4.50 (m, 2 H), 4.94 (d, J=10.1 Hz, 1 H), 5.01 (dd, J=17.1, 1.8 Hz, 1 H), 5.11 (dd, J=10.4, 1.8 Hz, 1 H), 5.30 (d, J=15.6 Hz, 1 H), 5.80 (m, 2 H), 8.57 (s, 1 H). LC-MS (Method J, retention time: 3.21 min), MS m/z 522 (M⁺+1).

Step 26iii: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-butoxycarbonylamino-18-hydroxy-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester

5




A solution of 1-([1-(2(*S*)-*tert*-Butoxycarbonyl-amino-non-8-enoyl)-4(*R*)-hydroxy-pyrrolidine-2(*S*) carbonyl]-(1*R*)-amino)-2(*S*)-vinylcyclopropane-carboxylic acid ethyl ester (800 mg, 1.53 mmol) in 2 L of methylene chloride was flushed with N₂ for 0.5 h. Then tricyclohexylphosphine[1,3-bis(2,4,6-trimethyl-phenyl)-4,5-dihydroimidazol-2-ylidene][benzylidene]-ruthenium (IV) dichloride (Strem) (64 mg, 0.075 mmol) was added, and the mixture was flushed with N₂ for another 10 min. The light orange homogeneous solution was refluxed for 2 h to give a dark orange solution. The reaction mixture was cooled to rt and concentrated in vacuo to give an orange oil. Flash chromatography (ethyl acetate) gave 460 mg (61%) of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-butoxycarbonylamino-18-hydroxy-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid ethyl ester as a gray solid. ¹H NMR (500 MHz, CDCl₃) δ

1.19 (t, *J*=7.2 Hz, 3 H), 1.42 (s, 9 H), 1.22-1.8 (m, 8 H), 1.87 (m, 2 H), 2.03-2.22 (m, 4 H), 2.63 (m, 1 H), 3.65 (m, 1 H), 4.09 (m, 3 H), 4.45 (m, 1 H), 4.56 (s, 1 H), 4.82 (m, 1 H), 5.23 (m, 1 H), 5.51 (s, 1 H), 7.16 (s,

1 H). LC-MS (Method J, retention time: 2.97 min), MS m/z 494 ($M^+ + 1$).

Step 26iv: Preparation of Example 26, (1*S*,4*R*,6*S*,14*S*,18*R*)-
5 7-*cis*-14-*tert*-butoxycarbonylamino-18-hydroxy-2,15-dioxo-
3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic
acid




10

To a solution of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-
butoxycarbonylamino-18-hydroxy-2,15-dioxo-3,16-
diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid
ethyl ester (493 mg, 1.0 mmol) in THF (4 mL), methanol (1
15 mL), and water (2 mL), was added powdered lithium
hydroxide (480 mg, 20 mmol), and the light yellow slurry
stirred at rt under N₂ for 16 h. The mixture was then
concentrated in vacuo and the residue partitioned between
ether and water. The ether phase was discarded, and the
20 aqueous phase was treated with 1 N HCl until pH 4. This
acidic solution was extracted with EtOAc three times. The
combined EtOAc extracts were dried (MgSO₄) and
concentrated in vacuo to give 460 mg (98%) of Example 26,
(1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-butoxycarbonylamino-18-
25 hydroxy-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-
nonadec-7-ene-4-carboxylic acid as a gray solid. ¹H NMR
(500 MHz, CD₃OD) δ ppm 1.26 (t, $J=7.2$ Hz, 3 H), 1.35-1.52
(m, 15 H), 1.57-1.68 (m, 3 H), 1.79 (m, 1 H), 2.04 (m, 1

H), 2.16–2.41 (m, 3 H), 3.80 (dd, $J=10.7$, 4.3 Hz, 1 H), 3.88 (m, 1 H), 4.38 (dd, $J=8.9$, 3.1 Hz, 1 H), 4.55 (m, 2 H), 5.39 (t, $J=9.8$ Hz, 1 H), 5.58 (m, 1 H). LC-MS (Method J, retention time: 2.64 min), MS m/z 466 ($M^+ + 1$).

5




Step H) To a suspension of 58 mg (0.125 mmol) of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-butoxycarbonylamino-18-hydroxy-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid, Example 26, and 31 mg (0.125 mmol) of LaCl₃ in 1.2 mL of DMF cooled to -78 °C was added 0.62 mL (0.62 mmol) of 1M KOtBu in THF, followed by 4-Chloro-2-(4-isopropylthiazol-2-yl)-7-methoxyquinoline, product of step G. The mixture was stirred one hour and warmed to rt showing little conversion to the product. Approximately 0.75 equivalent of solid KOtBu was added (10 mg), and the mixture stirred overnite. After 14h, approximately one-half equivalent of 4-Chloro-2-(4-isopropylthiazol-2-yl)-7-methoxyquinoline (22mg) and LaCl₃ (16 mg) were added, and the mixture stirred overnite. The mixture was quenched with pH 4 buffer and extracted into EtOAc. The combined EtOAc extracts were dried (MgSO₄), concentrated in vacuo


and dissolved up in 2 mL of MeOH. This solution was injected onto a preparative HPLC using the following conditions: Column Xterra 30X100 mm S5, 30% to 100% Solvent B/A for 18 min gradient, hold time 5 min; where
5 Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N
10 aqueous HCl solution, and repeatedly extracted with EtOAc. The combined EtOAc extracts were dried (MgSO₄) and concentrated in vacuo to provide 22 mg (31%) of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-Butoxycarbonylamino-18-[2-(2-isopropylthiazol-4-yl)-7-methoxyquinolin-4-yloxy]-
15 2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nona-dec-7-ene-4-carboxylic acid: ¹H NMR (500 MHz, Methanol-d⁴) δ 1.08-1.67 (m, 11 H), 1.39 (s, 9 H), 1.41 (d, *J*=7.0 Hz, 6 H), 1.78 (m, 1 H), 1.93 (m, 1 H), 2.32 (q, *J*=9 Hz, 1 H), 2.74 (m, 1 H), 3.23 (m, 1 H), 3.97 (m, 3 H), 4.07 (d, *J*=9 Hz,
20 1 H), 4.20 (d, *J*=8.2 Hz, 1 H), 4.66 (t, *J*=8.2 Hz, 1 H), 4.76 (d, *J*=11.3 Hz, 1 H), 5.36 (m, 1 H), 5.59 (m, 1 H), 5.64 (s, 1 H), 6.91 (s, 1 H), 7.13 (d, *J*=8.9 Hz, 1 H), 7.42 (s, 1 H), 7.45 (s, 1 H), 7.70 (s, 1 H), 8.19 (d, *J*=8.9 Hz, 1 H), 8.68 (s, 1 H). LC-MS *m/e* 748 (retention
25 time: 203, method D).


Step I: Preparation of Compound 27, Example 27,
(1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-{4-Cyclopropanesulfonylamino-
carbonyl-18-[2-(2-isopropylthiazol-4-yl)-7-methoxy-
30 quinolin-4-yloxy]-2,15-dioxo-3,16-diazatricyclo-
[14.3.0.0^{4,6}]nonadec-7-en-14-yl}carbamic acid *tert*-butyl
ester)

Step I) To a solution of 18 mg (0.024 mmol) of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-*tert*-Butoxycarbonylamino-18-[2-(2-isopropylthiazol-4-yl)-7-methoxyquinolin-4-yloxy]-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nona-dec-7-ene-4-carboxylic acid (product of Step H) in 2 mL of THF, was added CDI (29 mg, 0.036 mmol), and the resulting solution refluxed for 60 min and allowed to cool down to rt. Cyclopropylsulfonamide (4.4 mg, 0.036 mmol) was added in one portion before the addition of a neat solution of DBU (5.1 μ L, 0.034 mmol). The reaction was stirred for 16 h, diluted to 2 mL of methanol and injected onto a preparative HPLC using the following conditions: Column Xterra 30X100 mm S5, 30% to 100% Solvent B/A for 30 min gradient, hold time 5 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined EtOAc extracts were dried (MgSO₄) and concentrated in vacuo to provide 6.4 mg (31%) Compound 27, Example 27, as a foam: ¹H NMR (500 MHz, Methanol-d₄) δ 0.96-1.56(m, 17 H), 1.18 (s, 9 H), 1.61 (dd, *J*=9.5, 5.5 Hz, 1 H), 1.72 (dd, *J*=8.1, 5.3 Hz, 1 H), 1.84 (m, 2 H), 2.38 (m, 1 H), 2.61 (m, 1 H), 2.70 (m, 1 H), 2.80 (m, 1 H), 2.90 (m, 1 H), 3.24 (m, 1 H), 3.96 (s, 3 H), 4.07 (m, 1 H), 4.15 (d, *J*=10.7 Hz, 1 H), 4.66 (m, 1 H), 4.86 (m, 1 H), 5.08 (m, 1 H), 5.69 (m, 2 H), 7.13 (d, *J*=9.2 Hz, 1 H), 7.43 (s, 1 H), 7.45 (s, 1 H), 7.70 (s, 1 H), 8.20 (d, *J*=9.2 Hz, 1 H). LC/MS *m/e* 851 (retention time: 1.98, method D).




Example 28 (which corresponds to compound 35n in U.S. 6,323,180 at column 83, line 45) was prepared following the sequences of Example 35 of US Patent 6,323,180, but using a modification where the ethyl ester of 1-amino-2-vinylcyclopropane carboxylic acid hydrochloride was coupled in place of the methyl ester analogue (i.e. compound 35i of U.S. 6,323,180, column 80, line 5).



**Product of Steps 29f-g**

Steps 29f-29g) 500 mg (0.75mmol) of Example 28 was dissolved in HCl/dioxane (4.0M; 4.5 mL) and was stirred for 2.5 h at rt. The reaction mixture was concentrated and the crude product which was directly used in the next step, 29g. To suspension of the crude product from step 29f, 2(*S*)-tert-butoxycarbonyl-amino-8-nonenoic acid purchased from RSP Amino Acids (0.305g, 1.13mmol), NMM (0.44ml, 4.0mmol) in DMF (3.5mL) was added HATU (0.43g, 1.13 mmol). After being stirred for 2 days, the reaction mixture was diluted with EtOAc (200 mL), washed with water (50 mL), pH 4.0 buffer (50 mL), saturated aqueous NaHCO₃ (50 mL), dried (MgSO₄), and purified by a Biotage 40 M column (eluted with 0% to 5% MeOH in CH₂Cl₂) to supply the product of step 29g, as a yellow oil (0.50g, 81%) of approximate 80% purity by ¹H NMR: ¹H NMR (500 MHz, Solvent methanol-d₄) δ 1.32 (s, 9 H), 1.07-1.67 (m, 12 H), 1.74 (m, 1 H), 1.88 (m, 1 H), 2.02 (m, 2 H), 2.26 (m, 4 H), 2.42 (m, 1 H), 2.78 (m, 1 H), 3.94 (s, 3 H), 4.15 (m, 3 H), 4.49 (m, 1 H), 4.66 (m, 1 H), 4.83 (m, 1 H), 4.89 (d, *J*=9.5 Hz, 1 H), 4.96 (d, *J*=17.1 Hz, 1 H), 5.10 (d, *J*=10.4 Hz, 1 H), 5.28 (d, *J*=17.1 Hz, 1 H), 5.47 (s, 1 H), 5.76 (m, 2 H), 7.09 (m, 1 H), 7.37 (s, 1 H), 7.55 (s, 1 H), 7.95 (m, 1 H). LC-MS *m/e* 819 (retention time: 3.30, method D, except gradient time increased from 2 to 4 min).

127




Product of Step 29h

Step 29h: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-
 18-[2-(2-Acetylaminothiazol-4-yl)-7-methoxyquinolin-4-
 5 yloxy]-14-*tert*-butoxycarbonylamino-2,15-dioxo-3,16-
 diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid
 ethyl ester

Step 29h) To a solution of the product of from Step
 10 29g (380 mg of 80% purity, assumed 0.464 mmol) in 0.7L of
 Argon degassed benzene, was added 60 mg of
 Tricyclohexylphosphine [1,3-bis(2,4,6-trimethyl-phenyl)-
 4,5-dihydroimidazol-2-ylidene][benzylidene]ru-
 thenium(IV)dichloride catalyst from Strem, the mixture
 15 degassed under Ar and heated to reflux for 3 h. The
 reaction mixture was cooled to rt, the mixture degassed
 once more, an additional 60 mg portion of Tricyclohexyl-
 phosphine [1,3-bis(2,4,6-trimethyl-phenyl)-4,5-
 dihydroimidazol-2-ylidene][benzylidene]ruthen-
 20 ium(IV)dichloride catalyst added, and the mixture heated
 to reflux for 3 h. The reaction mixture was cooled to rt,
 the mixture degassed once more, and a final 5 mg portion
 of the catalyst added, and the mixture heated to reflux
 for an additional 1 h. The resulting dark brown solution
 25 was cooled to rt, concentrated in vacuo and purified over

a Biotage 40M column (eluted sequentially with 0% to 10% MeOH in CH₂Cl₂) to afford 309 mg (84%) of the titled product, a portion of which was taken directly into the next reaction: LC-MS *m/e* 791 (retention time: 1.83, method D).




Product of Step 29i

Step 29i: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-18-[2-(2-Acetylaminothiazol-4-yl)-7-methoxyquinolin-4-yloxy]-14-*tert*-butoxycarbonylamino-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid

Step 29i) Following an analogous experimental and purification procedure to Step 1i, (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-18-[2-(2-Acetylaminothiazol-4-yl)-7-methoxy-quinolin-4-yloxy]-14-*tert*-butoxycarbonylamino-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic ethyl ester (180 mg, 0.228 mmol) in 14 mL of THF and 3.5 mL of MeOH was reacted with 92 mg (2.3 mmol) of LiOH in 7 mL of H₂O to afford (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-18-[2-(2-Acetylaminothiazol-4-yl)-7-methoxyquinolin-4-yloxy]-14-*tert*-butoxycar-bonylamino-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid (150 mg, 86%): ¹H NMR (500 MHz, CD₃OD, some minor broadening due to rotomers) δ 11.29 (s, 9 H), 1.41 (m, 7

H), 1.62 (m, 2 H), 1.84 (m, 1 H), 1.96 (m, 1 H), 2.18 (m, 1 H), 2.24 (s, 3 H), 2.50 (m, 1 H), 2.62 (m, 1 H), 2.68 (m, 1 H), 3.96 (s, 3 H), 4.11 (s, 1 H), 4.29 (s, 1 H), 4.65 (s, 2 H), 5.51 (s, 3 H), 7.05 (s, 1 H), 7.37 (s, 1 H), 7.55 (s, 1 H), 7.95 (s, 1 H), 8.12 (s, 1 H): LC-MS m/e 763 (retention time: 3.09, method D, except gradient time increased from 2 to 4 min).



Compound 28
Example 29

10

Step 29j: Preparation of Compound 28, Example 29, (1S,4R,6S,14S,18R)-7-cis-[18-[2-(2-Acetylaminothiazol-4-yl)-7-methoxyquinolin-4-yloxy]-4-cyclopropanesulfonylaminocarbonyl-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-en-14-yl}carbamic acid tert-butyl ester

15


Step 29j) Following an analogous experimental and purification procedure to Step 1j, 52 mg (0.068 mol) of the product from step 29i, (1S,4R,6S,14S,18R)-7-cis-18-[2-(2-Acetylaminothiazol-4-yl)-7-methoxyquinolin-4-yloxy]-14-tert-butoxycarbonylamino-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid, was reacted with 16 mg (0.095 mmol) of CDI, 12 mg (0.095 mmol) of cyclopropyl-sulfonamide and 14 μ L (0.095

20

25


mmol) of DBU, to afford 25 mg (42%) of Compound 28, Example 29, (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-{18-[2-(2-Acetylaminothiazol-4-yl)-7-methoxyquinolin-4-yloxy]-4-cyclopropanesulfon-ylaminocarbonyl-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-en-14-yl}carbamic acid *tert*-butyl ester. To purify the compound, the compound was dissolved in 4 mL of MeOH and injected twice onto a preparative HPLC column (2 X 2 mL injections), afterwhich, an extractive purification was undertaken to isolate the product from the HPLC fractions: Column Xterra 30X100 mm S5, 35% to 80% Solvent B/A for 38 min gradient, hold time 2 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined EtOAc extracts were dried (MgSO₄) and concentrated in vacuo to provide the product, Compound 28, Example 29: ¹H NMR (500 MHz, Solvent methanol-*d*⁴) δ 0.76-1.61 (m, 20 H), 1.64 (dd, *J*=9.6, 5.3 Hz, 1 H), 1.75 (dd, *J*=8.1, 5.3 Hz, 1 H), 1.87 (m, 2 H), 2.29 (s, 3 H), 2.42 (m, 1 H), 2.60 (m, 1 H), 2.68 (m, 1 H), 2.81 (m, 1 H), 2.93 (m, 1 H), 3.99 (s, 3 H), 4.09 (m, 1 H), 4.22 (m, 1 H), 4.68 (m, 1 H), 4.88 (m, 1 H), 5.12 (m, 1 H), 5.59 (s, 1 H), 5.71 (m, 1 H), 7.10 (d, *J*=8.6 Hz, 1 H), 7.42 (s, 1 H), 7.63 (s, 1 H), 8.05 (s, 1 H), 8.20 (d, *J*=8.6 Hz, 1 H): LC-MS *m/e* 866 (retention time: 3.08, method D, except gradient time increased from 2 to 4 min).

Preparation of Example 30, Compound 29,
(1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-[4-Cyclopropanesulfonyl-
aminocarbonyl-2,15-dioxo-18-(quinolin-4-yloxy)-3,16-
diazatricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid
5 *tert*-butyl ester



Example 30, Compound 29

Step 30A: Preparation of 4(*R*)-(Quinolin-4-yloxy)-
10 pyrrolidine-1,2(*S*)-dicarboxylic acid 1-*tert*-butyl ester




Step 30A) A suspension of Boc-*L*-4-hydroxyproline [*N*-
15 Boc (2*S*,4*R*)-hydroxyproline] (10 g, 43.3 mmol) (CHEM-IMPEX
International) in DMSO (150 mL) was cooled in an ice bath
for ~ 3 minutes, and then *t*-BuOK (12.1 g, 108.2 mmol) was
added. The mixture was stirred for a few minutes until a
solid mass formed. At this time the reaction mixture was
20 allowed to warm to rt over 1.5 h to give a clear

colorless solution. 1-Chloroquinoline (7.7 g, 47.2 mmol) was added in two portions 10 min apart. The reaction was stirred for 24 h at rt. The dark reaction mixture was partitioned between ether (500 mL) and water (700 mL).

5 The aqueous phase was acidified to pH 4 using 4 N HCl. The resulting white precipitate was filtered and then washed with water (200 mL) to give the titled product of Step 30A as an off-white solid (13 g, 84%): ^1H NMR (500 MHz, DMSO) δ 1.32, 1.35 (2s, 9 H), 2.36 (m, 1 H), 2.66 (m, 1 H), 3.73 (m, 2 H), 4.35 (m, 1 H), 5.34 (s, 1 H), 7.06 (d, $J=5.2$ Hz, 1 H), 7.56 (t, $J=8.2$ Hz, 1 H), 7.75 (t, $J=8.2$ Hz, 1 H), 7.95 (d, $J=8.2$ Hz, 1 H), 8.11 (t, $J=8.2$ Hz, 1 H), 8.72 (d, $J=5.2$ Hz, 1 H). LC-MS (Method J, retention time: 1.87 min), MS m/z (M^++1).

15

Step 30B: Preparation of (2S)-(1(R)-ethoxycarbonyl-2(S)-vinylcyclopropylcarbamoyl)-4(R)-(quinolin-4-yloxy)pyrrolidine-1-carboxylic acid tert-butyl ester



20

Step 30B) A stirred slurry of 4(R)-(Quinolin-4-yloxy)-pyrrolidine-1,2(S)-dicarboxylic acid 1-tert-butyl ester (5.0 g, 14 mmol) in a mixture of 100 mL of methylene chloride and 100 mL of DMF was treated sequentially with N-methyl morpholine (3 mL, 27.3 mmol), HATU (PE biosystems) (6.3 g, 16.5 mmol), and 1R,2S-1-

amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride (3.2 g, 16.7 mmol). The solution was stirred at rt under N₂ for 18 h, and then concentrated in vacuo to give 10 g of a brown oil. This oil was


5 partitioned between ethyl acetate and sat. aq. NaHCO₃. The organic phase was washed with brine, dried (MgSO₄), and concentrated in vacuo. Flash chromatography (10% MeOH in ethyl acetate) gave 7.0 g (100%) of the titled product of Step 30B as a yellow solid: ¹H NMR (300 MHz, CD₃OD) δ

10 1.25 (t, $J=7.3$ Hz, 3 H), 1.44, 1.39 (2 s, 10 H), 1.79 (m, 1 H), 2.20 (m, 1 H), 2.43 (m, 1 H), 2.71 (m, 1 H), 3.92 (m, 2 H), 4.15 (q, $J=7.3$ Hz, 2 H), 4.42 (m, 1 H), 5.11 (m, 1 H), 5.29-5.39 (m, 2 H), 5.75 (m, 1 H), 7.05 (d, $J=5.5$ Hz, 1 H), 7.57 (t, $J=8.1$ Hz, 1 H), 7.76 (t, $J=8.4$

15 Hz, 1 H), 7.95 (d, $J=8.4$ Hz, 1 H), 8.16 (d, $J=8.4$ Hz, 1 H), 8.69 (d, $J=5.5$ Hz, 1 H). LC-MS (Method J, retention time: 2.35 min), MS m/z 496 (M^++1).

Step 30C: Preparation of 1-{[4(R)-(Quinolin-4-yloxy)-

20 pyrrolidine-2(S)-carbonyl]-1(R)-amino}-2(S)-vinyl-
cyclopropanecarboxylic acid ethyl ester bis hydrochloride




Step 30C) A stirred slurry of (2S)-(1(R)-

25 ethoxycarbonyl-2(S)-vinylcyclopropylcarbamoyl)-4(R)-(quinolin-4-yloxy)pyrrolidine-1-carboxylic acid tert-butyl ester (7.0 g, 14.1 mmol) was treated with 150 mL (300 mmol) of 2 N HCl/ether (Aldrich) for 24 h. The

reaction mixture was concentrated in vacuo to give 7.0 g (94%) of the titled product from step 30C as a yellow solid: ^1H NMR (500 MHz, CD_3OD) δ 1.28 (t, $J=7.0$ Hz, 3 H), 1.52 (dd, $J=5.5$ Hz, 4.0 Hz, 1 H), 1.65 (dd, $J=7.6$ Hz, 6.7 Hz, 1 H), 1.83 (dd, $J=5.5$ Hz, 2.7 Hz, 1 H), 2.29 (q, $J=8.8$ Hz, 1 H), 2.65 (m, 1 H), 3.02 (m, 2 H), 4.00 (s, 2 H), 4.20 (q, $J=7.0$ Hz, 2 H), 4.74 (dd, $J=7.6$ Hz, 2.7 Hz, 1 H), 5.17 (d, $J=10.4$ Hz, 1 H), 5.35 (d, $J=17.1$ Hz, 1 H), 5.81 (m, 1 H), 5.95 (m, 1 H), 7.69 (d, $J=6.7$ Hz, 1 H), 7.98 (t, $J=8.2$ Hz, 1 H), 8.19 (m, 2 H), 8.66 (d, $J=8.2$ Hz, 1 H), 9.14 (d, $J=6.7$ Hz, 1 H). LC-MS (Method J, retention time: 1.34 min), MS m/z 396 ($\text{M}^+ + 1 - 2\text{HCl}$).

Step 30D: Preparation of 1-([1-(2(S)-tert-butoxycarbonylamino-non-8-enoyl)-4(R)-(quinolin-4-yloxy)pyrrolidine-2(S)-carbonyl]-1(R)-amino)-2(S)-vinylcyclopropanecarboxylic acid ethyl ester




20

Step 30D) A solution of 2(S)-tert-butoxycarbonyl-amino-8-nonenoic acid (1.3 g, 4.79 mmol, purchased from RSP Amino Acids) dissolved in 100 mL of dichloromethane was treated sequentially with 1-([4(R)-(Quinolin-4-yloxy)pyrrolidine-2(S)-carbonyl]-1(R)-amino)-2(S)-vinylcyclopropanecarboxylic acid ethyl ester bis hydrochloride (2.0 g, 4.28 mmol), N-methyl morpholine (1.5 mL, 13.6

mmol), and HATU (PE biosystems) (1.92 g, 5.05 mmol). The reaction mixture was stirred at rt under N₂ for 16 h, and then concentrated in vacuo. The residue was partitioned between ethyl acetate and pH 4 buffer (biphthalate). The organic phase was washed with sat. aq. NaHCO₃, dried (MgSO₄), and concentrated in vacuo to give 2.2 g of the crude product. Flash chromatography (10% methanol/ethyl acetate) gave 2.0 g (73%) of the titled product of Step 30D as a yellow solid: ¹H NMR (500 MHz, CD₃OD) δ 1.27 (s, 9 H), 1.21-1.45 (m, 10 H), 1.63 (dd, J=11.29, 6.71 Hz, 2 H) 1.75 (m, 1 H) 2.04 (m, 2 H) 2.28 (q, J=8.75 Hz, 1 H) 2.44 (m, 1 H) 2.76 (m, 1 H) 4.01-4.25 (m, 4 H) 4.55 (d, J=12.21 Hz, 1 H) 4.67 (t, J=8.55 Hz, 1 H) 4.93 (d, J=10.07 Hz, 1 H) 5.00 (d, J=15.26 Hz, 1 H) 5.12 (dd, J=10.38, 1.83 Hz, 1 H) 5.31 (dd, J=17.24, 1.68 Hz, 1 H) 5.51 (s, 1 H) 5.79 (m, 2 H) 7.08 (d, J=5.49 Hz, 1 H) 7.56 (t, J=7.32 Hz, 1 H) 7.77 (t, J=7.48 Hz, 1 H) 7.97 (d, J=8.54 Hz, 1 H) 8.27 (d, J=8.55 Hz, 1 H) 8.73 (d, J=5.19 Hz, 1 H). LC-MS (Method J, retention time: 3.06 min), MS m/z 649 (M⁺+1).


Step 30E: Preparation of (1S,4R,6S,14S,18R)-7-cis-[14-tert-Butoxycarbonylamino-2,15-dioxo-18-(quinolin-4-yloxy)-3,16-diaza-tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester



Step 30E) A solution of 1-[[1-(2(*S*)-tert-Butoxycarbonylamino-non-8-enoyl)-4(*R*)-(quinolin-4-yloxy)pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino}-2(*S*)-vinylcyclopropanecarboxylic acid ethyl ester (1.6 g, 2.46 mmol) in 2 L of methylene chloride was flushed with N₂ for 0.5 h. Then tricyclohexylphosphine[1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene]-[benzylidene]-ruthenium (IV) dichloride (Strem) (150 mg, 0.17 mmol) was added, and the mixture was flushed with N₂ for another 10 min. The light orange homogeneous solution was refluxed for 16 h. Another 50 mg of the Ru catalyst was added, and the reaction mixture was refluxed for another 8 h. Finally, another 50 mg of the Ru catalyst was added and stirring was continued for 3 days at rt. The reaction mixture was concentrated in vacuo to give 1.7 g of an orange oil. Flash chromatography (ethyl acetate to 5% methanol in ethyl acetate) gave 1.1 g (68%) of the titled product of Step 30E as a white solid: ¹H NMR (500 MHz, CD₃OD), δ 1.19 (s, 9 H), 1.26 (t, *J*=7.0 Hz, 3 H), 1.29-1.59 (m, 7 H), 1.63 (dd, *J*=8.5 Hz, 4.8 Hz, 1 H), 1.67 (dd, *J*=9.5 Hz, 4.8 Hz, 1 H), 1.78-1.85 (m, 1 H), 1.92-1.99 (m, 1 H), 2.50 (m, 2 H), 2.70 (m, 1 H), 2.85 (q, *J*=8.8 Hz, 1 H), 4.06 (d, *J*=11.9 Hz, 1 H), 4.10-4.18 (m, 2 H), 4.23 (d, *J*=10.7 Hz, 1 H), 4.68 (m, 1 H), 4.79

(d, $J=11.3$ Hz, 1 H), 5.37 (t, $J=10.1$ Hz, 1 H), 5.53 (s, 1 H), 5.64 (m, 1 H), 7.08 (d, $J=5.5$ Hz, 1 H), 7.52 (t, $J=7.6$ Hz, 1 H), 7.75 (t, $J=8.2$ Hz, 1 H), 7.95 (d, $J=8.2$ Hz, 1 H), 8.72 (d, $J=5.5$ Hz, 1 H). (LC-MS (Method
 5 J, retention time: 2.71 min), MS m/z 621 (M^++1)).


Step 30F, Preparation of (1S,4R,6S,14S,18R)-7-cis-[14-
 tert-Butoxycarbonylamino-2,15-dioxo-18-(quinolin-4-
 yloxy)-3,16-diaza-tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-
 10 carboxylic acid



Step 30F) A solution of (1S,4R,6S,14S,18R)-7-cis-
 15 [14-tert-Butoxycarbonylamino-2,15-dioxo-18-(quinolin-4-yloxy)-3,16-diaza-tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester (1.10 g, 1.77 mmol) was dissolved in the mixed solvent system: THF (8 mL), methanol (2 mL), and water (4 mL). Powdered lithium
 20 hydroxide (425 mg, 17.7 mmol) was added. The light yellow slurry was stirred at rt under N₂ for 16 h, and then concentrated in vacuo. The residue was partitioned between ether and water. The ether phase was discarded, and the aqueous phase was treated with 1 N HCl until the
 25 pH was 4. This acidic solution was extracted with ethyl acetate four times. The combined ethyl acetate extracts

were dried (MgSO_4) and concentrated in vacuo to give the product of step 30F (0.90 g, 86%) as a white solid: ^1H NMR (300 MHz, CD_3OD) δ 1.16 (s, 9 H), 1.24 (m, 1 H), 1.33-1.55 (m, 6 H), 1.61 (m, 2 H), 1.79 (m, 1 H), 1.92 (m, 1 H), 2.32 (q, $J=8.8$ Hz, 1 H), 2.58 (m, 2 H), 2.68 (m, 1 H), 4.05 (d, $J=10.6$ Hz, 1 H), 4.22 (m, 1 H), 4.64 (t, $J=8.8$ Hz, 1 H), 4.74 (m, 1 H), 5.36 (t, $J=10.6$ Hz, 1 H), 5.49 (s, 1 H), 5.61 (m, 1 H), 7.07 (d, $J=5.5$ Hz, 1 H), 7.49 (t, $J=7.7$ Hz, 1 H), 7.73 (t, $J=7.7$ Hz, 1 H), 7.92 (d, $J=8.8$ Hz, 1 H), 8.30 (d, $J=8.8$ Hz, 1 H), 8.70 (d, $J=5.5$ Hz, 1 H). LC-MS (Method J, retention time: 2.54 min), MS m/z 593 (M^++1).

Step 30G: Preparation of Compound 29 of Example 30,
(1S,4R,6S,14S,18R)-7-cis-[4-Cyclopropanesulfonylamino-
carbonyl-2,15-dioxo-18-(quinolin-4-yloxy)-3,16-diaza-
tricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]-carbamic acid
tert-butyl ester



Compound 29
Example 30


20

Step 30G) To a solution of (1S,4R,6S,14S,18R)-7-cis-14-tert-Butoxycarbonylamino-2,15-dioxo-18-(quinolin-4-yloxy)-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid (100 mg, 0.17 mmol) was dissolved in 3 mL

25

of THF was added CDI (38 mg, 0.24 mmol), and the mixture heated to reflux for one hour. The mixture was cooled to rt and treated sequentially with cyclopropylsulfonamide (29 mg, 0.24 mmol) and DBU (36 mg, 0.24 mmol). After stirring for 24 h at rt, THF was removed by rotary evaporation. The residue was partitioned between ethyl acetate and pH 4 buffer. The organic phase was dried (MgSO₄) and concentrated in vacuo to give the crude product. Flash chromatography (10% MeOH/methylene chloride) gave 50 mg (42%) of the titled product, Compound 29 of Example 30, (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-[4-Cyclopropanesulfonyl-aminocarbonyl-2,15-dioxo-18-(quinolin-4-yl)oxy]-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid *tert*-butyl ester as a white powder: ¹H NMR (500 MHz, CD₃OD) δ 1.01 (m, 1 H), 1.18 (s, 9 H), 1.07-1.60 (m, 10 H), 1.65 (dd, *J*=9.5 Hz, 5.2 Hz, 1 H), 1.76 (dd, *J*=8.2 Hz, 5.5 Hz, 1 H), 1.86 (m, 2 H), 2.40 (m, 1H), 2.58 (m, 1 H), 2.68 (m, 1 H), 2.76 (m, 1 H), 2.93 (m, 1 H), 4.07 (d, *J*=10.7 Hz, 1 H), 4.19 (m, 1 H), 4.68 (dd, *J*=7.3 Hz, 1.81 Hz, 1 H), 4.85 (m, 1 H), 5.14 (t, *J*=7.6 Hz, 1 H), 5.55 (s, 1 H), 5.70 (m, 1 H), 6.68 (d, *J*=5.8 Hz, 1 H), 7.09 (d, *J*=5.2 Hz, 1 H), 7.51 (t, *J*=7.0 Hz, 1 H), 7.76 (t, *J*=7.0 Hz, 1 H), 7.95 (d, *J*=8.2 Hz, 1 H), 8.33 (d, *J*=8.2 Hz, 1 H), 8.72 (d, *J*=5.5 Hz, 1 H). LC-MS (Method A, retention time: 2.58 min), MS *m/z* 696 (M⁺+1).


Example 31, Preparation of Compound 30,
(1S,4R,6S,14S,18R)-7-cis-[4-Methanesulfonylamino-
carbonyl-2,15-dioxo-18-(quinolin-4-yloxy)-3,16-diaza-
tricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]-carbamic acid
 5 tert-butyl ester



(1S,4R,6S,14S,18R)-7-cis-[14-tert-
 10 butoxycarbonylamino-2,15-dioxo-18-(quinolin-4-yloxy)-
 3,16-diaza tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic
 acid, the product of Step F in Example 30 (50 mg, 0.075
 mmol) was reacted in analogous fashion with
 methanesulfonamide (12 mg, 0.12 mmol) as described in the
 15 preceding procedure above to give 10 mg (20%) of
 (1S,4R,6S,14S,18R)-7-cis-[4-methane-
 sulfonylamino-2,15-dioxo-18-(quinolin-4-yloxy)-
 3,16-diazatrimicyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]-
 carbamic acid tert-butyl ester as a white solid: ¹H NMR
 20 (300 MHz, CDCl₃) δ 1.26 (s, 9 H), 1.22-1.94 (m, 11 H),
 2.27 (m, 1 H), 2.55 (m, 1 H), 2.67 (m, 2 H), 3.17 (s, 3
 H), 4.05 (m, 1 H), 4.27 (m, 1 H), 4.61 (t, J=8.1 Hz, 1
 H), 4.73 (d, J=11.0 Hz, 1 H), 4.96-5.02 (m, 2 H), 5.35
 (s, 1 H), 5.71 (q, J=8.8 Hz, 1 H), 6.73 (d, J=5.1 Hz, 1
 25 H), 6.93 (bs, 1 H), 7.43 (t, J=7.7 Hz, 1 H), 7.69 (t,
 J=7.7 Hz, 1 H), 8.06 (d, J=8.8 Hz, 1 H), 8.15 (d, J=7.7

Hz, 1 H), 8.77 (d, $J=5.5$ Hz, 1 H). LC-MS (Method A, retention time: 2.43 min), MS m/z 670 (M^++1).

Preparation of Example 32, 2(S)-tert-butoxycarbonylamino-3-pentenylsulfanylpropionic acid methyl ester (Example 32) for use in Example 33.




Step 1: To a solution of *N*-Boc-cysteine methyl ester (3.36 g, 0.014 mol) in methanol (166 mL) at RT was added triethylamine (10.8 mL) and 1-bromopent-4-ene (3.19 g, 21 mmol, 1.5 equivalents) and the resulting solution was stirred at room temperature overnight. The mixture was then concentrated in vacuo and the resulting residual mixture was purified using flash chromatography (hexane, ethyl acetate gradient) to provide 1.76 g (41%) of the desired thioether. ^1H NMR (500 MHz, CDCl_3) δ 1.43 (s, 9H), 1.64 (m, 2H), 2.11 (m, 2H), 2.51 (m, 2H), 2.95 (m, 2H), 3.75 (s, 3H), 4.51 (m, 1H), 4.95–5.03 (m, 2H), 5.34 (m, 1H), 5.80 (1H, m). LC-MS (Method H, except gradient time was 3 min., and flow rate was 4 ml/min, retention time: 2.29 min), MS m/z 304 (M^++1).

Step 2: The thioether product of step 1 (9.51 g, 31.4 mmol) was added to a mixture of 1M LiOH in water (200mL) and THF (200mL) and the resulting mixture was stirred at room temperature overnight. The reaction mixture was then acidified using 1N hydrochloric acid and the resulting mixture was extracted several times with


ethyl acetate. The extracts were combined, dried over magnesium sulfate, and concentrated in vacuo to provide the desired acid which was used as is in the next reaction.

5

- Preparation of Example 33, Compound 31,
(1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-[4-Cyclopropanesulfonyl-
aminocarbonyl-18-(7-methoxy-2-phenyl-quinolin-4-yloxy)-
2,15-dioxo-12-thia-3,1
 10 6-diazatricyclo[14.3.0.⁰_{4,6}]nonadec-7-en-14-yl]-carbamic
acid *tert*-butyl ester.



- 15 Step 33A, Preparation of 1-{[1-(2(*S*)-*tert*-
Butoxycarbonylamino-3-pent-4-enylsulfanylpropionyl)-4(*R*)-
(7-methoxy-2-phenylquinolin-4-yloxy)pyrrolidine-2(*S*)-
carbonyl]-1(*R*)-amino}-2(*S*)-vinylcyclopropanecarboxylic
acid ethyl ester.



Step 33A) Note: the product of Step 1f (1-([4-(7-Methoxy-2-phenylquinolin-4(R)-yloxy)pyrrolidine-2(S)-

5 carbonyl]-1(R)-amino}-2(S)-vinylcyclopropanecarboxylic acid ethyl ester was used as starting material for the preparation of Compound 31. The carboxylic acid (282 mg, 976 μ mol) of example 32 was added to a DMF solution containing HATU (445mg, 1.17 mmol, 1.2 eq.), NMM (375 μ L, 3.42 mmol, 3.5 eq.) and the TFA salt of the product of


10 step 1f (600mg, 976 μ mol, 1eq.) and the resulting solution was stirred at room temperature over night. The resulting mixture was quenched by the addition of pH4 buffer and the resulting mixture was extracted (3X) with

15 ethyl acetate. The extracts were combined, washed with brine, dried over magnesium sulfate and concentrated in vacuo. The crude product mixture was purified by flash chromatography (hexane, ethyl acetate) to provide the product of Step 33A (753 mg, 84%). ^1H NMR (500 MHz,

20 CDCl_3) δ 1.20-1.25 (m, 3H), 1.36 (s, 9H), 1.54-1.68 (m, 4H), 1.88 (m, 1H), 2.10-2.20 (m, 3H), 2.55-2.64 (m, 2H), 2.73-2.97 (m, 3H), 4.08-4.15 (m, 3H), 4.35 (m, 1H), 4.68 (m, 1H), 4.90-5.04 (m, 3H), 5.10 (m, 1H), 5.27-5.44 (m, 3H), 5.68-5.82 (m, 2H), 7.05-7.09 (m, 2H), 7.42-7.59 (m,

25 4H), 7.92-8.19 (m, 3H).


Step 33B, Preparation of 14-*tert*-Butoxycarbonylamino-18-
 (7-methoxy-2-phenyl-quinolin-4-yloxy)-2,15-dioxo-12-thia-
 3,16-diaza-tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic
 5 acid ethyl ester



Step 33B) The tripeptide ester product of step 33A
 10 (390 mg, 505 μ mol) was dissolved in degassed (nitrogen
 purged) dichloroethane (145mL), Tricyclohexylphos-
 phine[1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-
 2-ylidene]-[benzylidene]-ruthenium (IV) dichloride
 (Strem) (45 mg, 53 μ mol) was added, and the reaction
 15 mixture was heated to 45 C and maintained at this
 temperature for 5 hr. At this time, additional ruthenium
 catalyst (45 mg, 53 μ mol) was added to the reaction
 mixture and heating was maintained overnight. The
 reaction mixture was then concentrated in vacuo and
 20 purified by flash chromatography (hexanes, ethyl acetate)
 to provide 159.4 mg (42%) of the desired olefin
 metathesis cyclization product of Step 33B. ¹H NMR (500
 MHz, CDCl₃) δ 1.24 (t, *J* = 7Hz, 3H), 1.37 (m, 1H), 1.41
 (s, 9H), 1.50-1.75 (m, 3H), 1.85 (m, 1H), 1.96 (m, 1H),
 25 2.13 (m, 1H), 2.21-2.45 (m, 3H), 2.52 (m, 1H), 2.81 (m,

1H), 3.07-3.15 (m, 3H), 4.12-4.25 (m, 3H), 4.85 (m, 1H),
 5.09 (m, 1H), 5.37-5.42 (m, 2H), 5.5 (m, 1H), 5.6 (m,
 1H), 7.05 (s, 1H), 7.09 (m, 1H), 7.40-7.58 (m, 4H), 8.00-
 8.10 (m, 3H).). LC-MS (Method H, except gradient time
 5 was 2 min, flow rate was 5 ml/min, and retention time:
 1.55 min), MS m/z 745 ($M^+ + 1$).


Step 33C, Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-
 tert-Butoxycarbonylamino-18-(7-methoxy-2-phenyl-quinolin-
 10 4-yloxy)-2,15-dioxo-12-thia-3,16-diaza-
 tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid



15 **Step 33C)** The macrocycle tripeptide ester product of
 step 33B (149.8 mg, 201 μ mol) was dissolved in a mixture
 of THF (4.1 mL), MeOH (1.9 mL), H₂O (0.45 mL) containing LiOH
 (97 mg, 4.04 mmol) and the resulting mixture was at
 stirred at room temperature overnight. The reaction was
 20 quenched by the addition of aqueous HCl (0.1N) and the
 resulting mixture was extracted with ethyl acetate. The
 organic extracts were dried with magnesium sulfate and
 concentrated in vacuo to provide a crude product mixture
 that was purified using flash chromatography (ethyl
 25 acetate, then 10% MeOH/CH₂Cl₂ as eluent) to provide the

desired macrocycle carboxylic acid (109.3 mg, 76%). ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 1.1 (m, 1H), 1.16 (s, 9H), 1.36 (m, 1H), 1.41-1.55 (m, 3H), 1.60 (m, 1H), 1.95 (m, 1H), 2.27 (m, 1H), 2.32-2.43 (m, 2H), 2.45-2.60 (m, 3H), 2.71 (m, 1H), 2.90 (m, 1H), 3.92 (s, 3H), 4.05 (m, 1H), 4.25-4.55 (m, 2H), 5.42-5.70 (m, 3H), 7.10 (m, 1H), 7.19 (m, 1H), 7.32-7.62 (m, 3H), 7.75 (m, 1), 8.04 (m, 1H), 8.24-8.30 (m, 2H), 8.50 (s, 1H). LC-MS (Method H, except gradient time was 2 min., and flow rate was 5 ml/min, retention time: 1.48 min), MS m/z 717 (M^++1).

Step 33D, Preparation of Example 33, Compound 31,
(1S,4R,6S,14S,18R)-7-cis-[4-Cyclopropanesulfonyl-
aminocarbonyl-18-(7-methoxy-2-phenyl-quinolin-4-yloxy)-
2,15-dioxo-12-thia-3,16-diaza-
tricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]-carbamic acid
tert-butyl ester




20

Step 33D) The macrocycle acid product of step 33C (77.9 mg, 109 μmol , 1 eq.) was dissolved in THF (770 μL) and CDI (21.1 mg, 130 μmol , 1.2 eq.) was added. The mixture was then heated to reflux and maintained at that temperature for 2h. The reaction mixture was then cooled

25

to room temperature, cyclopropylsulfonamide (15.5 mg, 128 umol, 1.2eq) and DBU (18.7 uL, 125 umol, 1.2eq) were added, and the resulting mixture was stirred at room temperature overnight. LC/MS analysis indicated the product mixture consisted of a mixture of starting material and desired product. The resulting mixture was partitioned between methylene chloride and 1N aqueous HCl, followed by back extraction of the aqueous layer. The combined extracts were dried over magnesium sulfate and concentrated in vacuo. The mixture of starting material and desired product was then resubjected to the above described reaction conditions using identical amounts of reagents. After the duration of the reaction the reaction mixture was worked up as before and then further purified by chromatography (Biotage 12M column, 75% ethyl acetate/hexane) to provide the desired macrocycle sulfonamide, Example 33, Compound 31, (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-[4-Cyclopropanesulfonyl-aminocarbonyl-18-(7-methoxy-2-phenyl-quinolin-4-yloxy)-2,15-dioxo-12-thia-3,16-diaza-tricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]-carbamic acid tert-butyl ester (30.6 mg, 34%). ¹H NMR (500 MHz, methanol-d₄) δ 0.90-1.50 (m, 7H), 1.24 (2, 9H), 1.55 (m, 1H), 2.60-2.67 (m, 2H), 2.83 (m, 1H), 2.15 (m, 1H), 2.30-2.47 (m, 2H), 2.5-2.9 (m, 7H), 3.8 (s, 3H), 4.22 (m, 1H), 4.35-4.41 (m, 2H), 4.7 (m, 1H), 5.38 (m, 1H), 5.48-5.68 (m, 2H), 6.82 (m, 1H), 7.06 (m, 1H), 7.25 (s, 1H), 7.35 (s, 1H), 7.49-7.60 (m, 3H), 8.03-8.15 (m, 3H). LC-MS (Method H, except gradient time was 2 min., flow rate was 5 ml/min, and retention time: 1.48 min), MS *m/z* 820 (M⁺+1).

Example 34, Preparation of Compound 32,
 (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-[4-Cyclopropanesulfonyl-
 aminocarbonyl-18-(7-methoxy-2-phenyl-quinolin-4-yloxy)-
 2,12,12,15-tetraoxo-12,16-thia-3,16-
 5 diazatricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid
tert-butyl ester



10

Oxone Step) To a solution of Compound 31 (9.5 mg, 11.6 μ mol) in methanol (0.276 mL) was added pH 7 buffer (0.093 mL) and the solution was cooled in an ice bath. Oxone (21 mg, 34.8 μ mol, 3 eq.) was added and the
 15 reaction was equilibrated to room temperature and stirred for two hours. Water was then added to the reaction mixture and the resulting mixture was extracted with three times with ethyl acetate. The combined organic extracts were concentrated in vacuo and the crude product
 20 was purified by preparative TLC (Hexane: Ethyl acetate, 1:3 eluent) to provide the desired product Example 34, Compound 32, (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-[4-Cyclopropanesulfonyl-aminocarbonyl-18-(7-methoxy-2-phenyl-quinolin-4-yloxy)-2,12,12,15-tetraoxo-12,16-thia-
 25 3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]carbamic acid *tert*-butyl ester (9.8 mg, 32%). ¹H NMR (500 MHz,


methanol- d_4) δ 0.74-0.94 (m, 3H), 1.0-1.08 (m, 2H), 1.26 (s, 9H), 1.22-1.33 (m, 2H), 1.58 (m, 1H), 1.76-1.82 (m, 2H), 2.04 (m, 1H), 2.17 (m, 1H), 2.32-2.40 (m, 2H), 2.60 (m, 1H), 2.82-2.89 (m, 2H), 3.65-3.78 (m, 2H), 3.95 (s, 3H), 4.38-4.45 (m, 2H), 4.75-4.99 (m, 3H), 5.48-5.62 (m, 3H), 7.14 (m, 1H), 7.29 (s, 1H), 7.42 (m, 1H), 7.46-7.58 (m, 3H), 8.06-8.18 (m, 3H). LC-MS (Method H, except gradient time was 2 min, flow rate was 5 ml/min, and retention time: 1.29 min), MS m/z 852 ($M^+ + 1$).

10

Example 35, Preparation of Compound 33,

(1S,4R,6S,14S,18R)-7-cis-[4-Cyclopropanesulfonyl-aminocarbonyl-2,15-dioxo-18-(8-trifluoromethylquinol-4-yloxy)-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl]-carbamic acid tert-butyl ester

15



Steps 35A-B) Compound 33, Example 35, was prepared by analogy using the chemistry described in the preparation of Compound 27 using Example 26 as starting material: Data for Compound 33: ^1H NMR (500 MHz, CD_3OD) δ ppm 1.02 (m, 1 H), 1.09 (s, 9 H), 1.05-1.60 (m, 10 H), 1.64 (dd, $J=9.5, 5.5$ Hz, 1 H), 1.75 (dd, $J=8.2, 5.5$ Hz, 1 H), 1.84 (bs, 2 H), 2.43 (q, $J=8.5$ Hz, 1 H), 2.59 (m, 1 H), 2.71 (m, 1 H), 2.79 (m, 1 H), 2.94 (m, 1 H), 4.03 (m, 1 H), 4.13 (d, $J=10.7$ Hz, 1 H), 4.71 (dd, $J=9.6,$


20

25

150

7.2 Hz, 1 H), 4.88 (m, 1 H), 5.11 (t, $J=9.3$ Hz, 1 H), 5.57 (s, 1 H), 5.72 (q, $J=8.9$ Hz, 1 H), 7.19 (d, $J=5.5$ Hz, 1 H), 7.60 (t, $J=7.6$ Hz, 1 H), 8.13 (d, $J=7.0$ Hz, 1 H), 8.58 (d, $J=8.2$ Hz, 1 H), 8.84 (d, $J=5.2$ Hz, 1 H).

5 LC-MS (Method J, retention time: 3.09 min), MS m/z 764 (M^++1).




Compound 34
Example 36

Method A: Preparation of Compound 34, Example 36,


10 (1S,4R,6S,14S,18R)-7-cis-{18-[2-(2-Acetylaminothiazol-4-yl)-7-methoxyquinolin-4-yloxy]-4-cyclopropanesulfon-
ylaminocarbonyl-2,15-dioxo-3,16-diazatricyclo-
[14.3.0.0^{4,6}]-nonadec-7-en-14-yl}carbamic acid tert-butyl
ester was accomplished using Steps 36A-36N

15

151



Preparation of 4(R)-(7-Methoxy-2-methoxycarbonyl-
quinolin-4-yloxy)pyrrolidine-1,2(S)-dicarboxylic acid 1-
tert-butyl ester




10

This compound was prepared following the sequences of Example 35 of US Patent 6,323,180.

Step 36A: Preparation of 2(S)--(1(R)-ethoxycarbonyl-2(S)-vinylcyclopropylcarbamoyl)-4(R)-(7-methoxy-2-methoxycarbonylquinolin-4-yloxy)pyrrolidine-1-carboxylic acid tert-butyl ester


5



To a solution of (1R,2S)-1-amino-2-vinylcyclopropane carboxylic acid ethyl ester hydrochloride (1.24 g, 6.47
10 mmol), DIPEA (4.18 mL, 32.35 mmol), and 4(R)-(7-Methoxy-2-methoxycarbonylquinolin-4-yloxy)-pyrrolidine-1,2(S)-dicarboxylic acid 1-tert-butyl ester (2.95 g, 6.61 mmol) in 35 mL of acetonitrile at RT, was added TBTU (2.76 g, 8.59 mmol, purchased from Aldrich). The reaction mixture
15 was stirred at rt under N₂ for 14 h, and then concentrated in vacuo. The residue was partitioned between ethyl acetate and sat. aq. NaHCO₃ (2 X 75 mL), dried (MgSO₄), and the organics were concentrated in vacuo to give the crude product. Flash chromatography
20 over a Biotage 65M column (eluted with 40%-95% ethyl acetate/hexane) gave 3.45 g (91%): ¹H NMR (500 MHz, Methanol-d⁴, rotamers~1/2) δ 1.28 (m, 3 H), 1.42 (s, 4 H), 1.47 (s, 6 H), 1.80 (m, 1 H), 2.24 (q, J=8.6 Hz, 1 H), 2.48 (m, 1 H), 2.74 (m, 1 H), 3.94 (m, 2 H), 3.99 (s,
25 3 H), 4.07 (s, 3 H), 4.20 (m, 2 H), 4.45 (m, 1 H), 5.15 (m, 1 H), 5.33 (m, 1 H), 5.47 (s, 1 H), 5.81 (m, 1 H), 7.31 (m, 1 H), 7.52 (s, 1 H), 7.56 (d, J=1.5 Hz, 1 H), 8.10 (d, J=9 Hz, 1 H). LC-MS m/e 588 (retention time:

4.76, method D, except gradient time increased from 2 to 8 min and hold time increased from 1 to 2 min).

5 Step 36B: Preparation of 2(S)-(1(R)-ethoxycarbonyl-2(S)-vinylcyclopropylcarbamoyl)-4(R)-(7-methoxy-2-methoxycarbonylquinolin-4-yloxy)pyrrolidine dihydrochloride



2 HCl Salt

Product of Step 36B

10


A solution of 3.05 g (5.23 mmol) of 4(R)-(7-Methoxy-2-methoxycarbonylquinolin-4-yloxy)-pyrrolidine-1,2(S)-dicarboxylic acid 1-tert-butyl ester was stirred in 26 mL
15 (116.5 mmol) of 4N HCl in dioxanes but solidified after 10 min. To this mixture was added an additional 26 mL (116.5 mmol) of 4N HCl in dioxanes, the resulting suspension chopped up to allow free stirring and the mixture stirred 3.5 h. The reaction mixture was
20 concentrated and was azeotroped several time with dioxanes (4X100 mL) to afford the titled compound of Step 36B. The resulting solid was then used directly in the next reaction: ¹H NMR (500 MHz, Methanol-d⁴) δ 1.25 (t, J=7.0 Hz, 3 H) 1.49 (dd, J=9.0, 5.34 Hz, 1 H) 1.79 (dd, J=7.6, 5.5 Hz, 1 H), 2.27 (q, J=8.6 Hz, 1 H), 2.61 (m, 1 H), 3.00 (m, 1 H), 3.97 (s, 2 H), 4.08 (s, 3 H), 4.16 (m, 5 H), 4.70 (m, 1 H), 5.13 (d, J=10.4 Hz, 1 H), 5.32 (d, J=17.1 Hz, 1 H), 5.78 (m, 1 H), 6.01 (s, 1 H), 7.58 (m, 1

25

H), 7.70 (s, 1 H), 7.91 (s, 1 H), 8.55 (d, $J=9$ Hz, 1 H).
LC-MS (Method D, retention time: 1.12 min), MS m/z
484 (M^++1).

- 5 Step 36C: Preparation of 1-([1-(2(*S*)-*tert*-butoxycarbonyl-
amino-non-8-enoyl)-4-(7-methoxy-2-
*methoxycarbonylquinolin-4(*R*)-yloxy)pyrrolidine-2(*S*)-*
*carbonyl]-1(*R*)-amino]-2(*S*)-vinylcyclopropanecarboxylic*
acid ethyl ester

10



Product of Step 36C


- A solution of 2(*S*)-*tert*-butoxycarbonyl-amino-8-nonenoic acid (2.13 g, 7.84 mmol, purchased from RSP
15 Amino Acids) dissolved in 18 mL of *N,N*-dimethylformamide was treated sequentially with 2(*S*)-(1(*R*)-ethoxycarbonyl-2(*S*)-vinylcyclopropylcarbonyl)-4(*R*)-(7-methoxy-2-methoxycarbonylquinolin-4-yloxy)pyrrolidine dihydrochloride (the product of step 36C-assumed 5.23
20 mmol), *N*-methyl morpholine (4.76 mL, 47 mmol), and HATU (PE biosystems) (2.97 g, 7.84 mmol). The reaction mixture was stirred at rt for 16 h, and then concentrated in vacuo. The residue was partitioned between 80% ethyl acetate/hexanes (400 mL) and water (100 mL). The organic
25 phase was washed with pH 4 buffer (100 mL), sat. aq. NaHCO_3 (100 mL), dried (MgSO_4), and concentrated in vacuo.

The residue was chromatographed over a Biotage 65M column (eluted with 50% to 100% ethyl acetate/hexanes) gave 3.43 g (89%) of the titled product of Step 36C as a foam: ^1H NMR (500 MHz, Methanol- d^4) δ 1.17-1.48 (m, 10 H), 1.30 (s, 9 H), 1.55 (m, 1 H), 1.64 (m, 1 H), 1.76 (dd, $J=7.9$, 5.2 Hz, 1 H), 2.03 (q, $J=7.2$ Hz, 2 H), 2.28 (q, $J=8.8$ Hz, 1 H), 2.44 (m, 1 H), 2.75 (dd, $J=14.3$, 7.9 Hz, 1 H), 3.98 (s, 3 H), 4.07 (s, 3 H), 4.15 (m, 4 H), 4.51 (d, $J=11.9$ Hz, 1 H), 4.68 (m, 1 H), 4.93 (d, $J=10.4$ Hz, 1 H), 4.99 (dd, $J=17.1$, 1.5 Hz, 1 H), 5.12 (dd, $J=10.4$, 1.8 Hz, 1 H), 5.31 (dd, $J=17.1$, 1.5 Hz, 1 H), 5.57 (s, 1 H), 5.80 (m, 2 H), 7.26 (dd, $J=9.2$, 2.4 Hz, 1 H), 7.51 (m, 1 H), 7.58 (s, 1 H), 8.17 (d, $J=9.2$ Hz, 1 H). LC-MS (Method D, retention time: 1.87 min), MS m/z 737 ($\text{M}^+ + 1$).

15

Step 36D: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-tert-Butoxycarbonylamino-18-(7-methoxy-2-methoxycarbonylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester

20



Product of Step 36D


To a solution of 1-{[1-(2(*S*)-tert-butoxycarbonylamino-non-8-enoyl)-4-(7-methoxy-2-carbomethoxyquinolin-4(*R*)-yloxy)-pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino]-2(*S*)-vinylcyclopropanecarboxylic acid ethyl ester (1.65 g,

25

2.24 mmol) in 1.4L of Argon degassed benzene, was added 200 mg of Tricyclohexylphosphine [1,3-bis(2,4,6-trimethyl-phenyl)-4,5-dihydroimidazol-2-ylidene]-[benzylidene]ruthen-ium(IV)dichloride catalyst, the mixture was degassed under Ar and heated to reflux for 3 h. The reaction mixture was cooled to rt, the mixture degassed again, 240 mg portion of Tricyclohexylphosphine [1,3-bis(2,4,6-trimethyl-phenyl)-4,5-dihydroimidazol-2-ylidene][benzylidene]ruthenium(IV)dichloride catalyst was added, and the mixture was heated to reflux for 3 h. The entire reaction sequence was repeated again using the exact same conditions. The combined dark brown solutions (resulting from 2 X 1.65 g metatheses reactions of 1-([1-(2(*S*)-*tert*-butoxycarbonylamino-non-8-enoyl)-4-(7-methoxy-2-carbomethoxyquinolin-4(*R*)-yloxy)-pyrrolidine-2(*S*)-carbonyl]-1(*R*)-amino]-2(*S*)-vinylcyclopropanecarboxylic acid ethyl ester) were concentrated in vacuo, purified over a Biotage 65 M column (eluted with 50% to 100% EtOAc in hexanes) to supply 2.00 g (63% yield, >99% pure) of the titled compound of Step 36D as a foam which solidified to a glass: ¹H NMR (500 MHz, Methanol-d⁴) δ 1.16-1.59 (m, 7 H), 1.20 (s, 9 H), 1.26 (t, *J*=7.0 Hz, 3 H), 1.61 (dd, *J*=8.4, 5.04 Hz, 1 H), 1.67 (dd, *J*=9.6, 5.0 Hz, 1 H), 1.80 (m, 1 H), 1.95 (m, 1 H), 2.34 (q, *J*=8.8 Hz, 1 H), 2.57 (m, 2 H), 2.70 (m, 1 H), 3.97 (s, 3 H), 4.06 (s, 3 H), 4.01-4.25 (m, 4 H), 4.67 (t, *J*=8.2 Hz, 1 H), 4.74 (d, *J*=11.3 Hz, 1 H), 5.37 (m, 1 H), 5.60 (m, 2 H), 7.18 (d, *J*=9.2 Hz, 1 H), 7.47 (s, 1 H), 7.53 (s, 1 H), 8.18 (d, *J*=9.2 Hz, 1 H). LC-MS (Method D except gradient time increased from 2 to 4 min, retention time: 3.03 min), MS *m/z* 709 (*M*⁺+1).

Step 36E: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-Amino-18-(7-methoxy-2-methoxycarbonylquinolin-4-yloxy)-2,15-dioxo-3,16-diaza-tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester dihydrochloride

5




Product of Step 36E

A solution of 1.60 g (2.26 mmol) of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-tert-Butoxycarbonylamino-18-(7-methoxy-2-methoxycarbonylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester (product of step 36D) in 17 mL (68 mmol) of 4N HCl/dioxanes was stirred. After 20 min. the reaction mixture solidified and 10 mL of CH₂Cl₂ was added. The resulting solid suspension was then chopped up to aid stirring, which proceeded for an additional 1h. After this time period the reaction was still not complete, so an additional portion of 4 mL (16 mmol) of 4N HCl/dioxanes was added and the mixture was stirred for an additional 1 h. The reaction mixture was then concentrated in vacuo to provide the product of Step 36E as a solid which was taken directly into step 36F: ¹H NMR (500 MHz, Methanol-d₄) δ 1.27 (m, 3 H), 1.27-1.64 (m, 7 H), 1.94 (m, 2 H), 2.16 (m, 1 H), 2.28 (q, J=9.2Hz, 1 H), 2.31 (m, 1 H), 2.72 (m, 1 H), 2.84 (m, 1 H), 4.10 (m, 3 H), 4.16 (m, 3 H), 4.22 (s, 3 H), 4.44 (m, 2 H), 4.84 (m,

1 H), 5.65 (m, 1 H), 5.99 (s, 1 H), 7.58 (dd, $J=9.2$, 2.1 Hz, 1 H), 7.73 (d, $J=2.1$ Hz, 1 H), 7.95 (s, 1 H), 8.45 (d, $J=9.2$ Hz, 1 H). LC-MS (Method D except gradient time increased from 2 to 4 min, retention time: 2.12 min), MS
 5 m/z 609 (M^++1).

Step 36F: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-Cyclopentyloxycarbonylamino-18-(7-methoxy-2-methoxycarbonylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester




Product of Step 36F

15 To a slurry of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-Amino-18-(7-methoxy-2-methoxycarbonylquinolin-4-yloxy)-2,15-dioxo-3,16-diaza-tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester dihydrochloride (product from step 36E-
 assumed 2.26 mmol) in 15 mL of THF cooled to 0 °C, was
 20 added 2 mL of Et₃N, followed by 5.7 mL (4.52 mmol) of freshly prepared 0.8M cyclopentyl chloroformate (prepared as described above). The mixture was stirred for 30 min then partitioned between 125 mL of ethyl acetate and 40 mL of a saturated aqueous NaHCO₃ solution. The ethyl
 25 acetate layer was dried (MgSO₄), concentrated and the crude residue chromatographed over a biotage 65M column

(eluted with 0% to 100% CH₂Cl₂/EtOAc) to afford 1.40g (86%) of the product of Step 36F as a white foam: ¹H NMR (500 MHz, Methanol-d₄) δ 1.23 (t, J=7.2 Hz, 3 H), 1.32-1.69 (m, 17 H), 1.79 (m, 1 H), 1.93 (m, 1 H), 2.31 (q, J=9.2 Hz, 1 H), 2.52 (m, 2 H), 2.67 (m, 1 H), 3.94 (s, 3 H), 4.03 (m, 1 H), 4.04 (s, 3 H), 4.11 (m, 2 H), 4.23 (dd, J=10.4, 2.75 Hz, 1 H), 4.54 (m, 1 H), 4.66 (m, 2 H), 5.33 (t, J=9.6 Hz, 1 H), 5.58 (m, 2 H), 7.17 (dd, J=9.2, 2 Hz, 1 H), 7.45 (d, J=2 Hz, 1 H), 7.52 (s, 1 H), 8.15 (d, J=9.2 Hz, 1 H). LC-MS (Method D except gradient time increased from 2 to 4 min, retention time: 3.12 min), MS m/z 721 (M⁺+1).

Step 36G: Preparation of (1S,4R,6S,14S,18R)-7-cis-18-(2-Carboxy-7-methoxyquinolin-4-yloxy)-14-cyclopentyloxy-carbonylamino-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid ethyl ester sodium salt



Product of Step 36G


20

To a solution of 1.37g (1.90 mmol) of (1S,4R,6S,14S,18R)-7-cis-14-Cyclopentyloxy-carbonylamino-18-(7-methoxy-2-methoxy-carbonylquinolin-4-yloxy)-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester (product from step 36G) in 23 mL of 2:1:1 THF/MeOH/H₂O was added 2.1 mL (2.1 mmol) of

25

1M aqueous NaOH. The solution was stirred 60 min at room temperature, glacial AcOH added dropwise until the solution pH was neutral and then the mixture concentrated in vacuo. The solid residue was dissolved in 100 mL of 30% THF/CH₂Cl₂, dried (MgSO₄), and concentrated to afford approximately 1.90 mmol of the product of Step 36G which was taken directly into the next reaction (Step 36H): ¹H NMR (500 MHz, Methanol-d₄) δ 1.22 (t, J=7.0 Hz, 3 H), 1.25-1.71 (m, 15 H), 1.79 (m, 1 H), 1.87 (m, 2 H), 1.94 (m, 1 H), 2.30 (q, J=9.0 Hz, 1 H), 2.50 (m, 2 H), 2.72 (m, 1 H), 3.95 (s, 3 H), 4.10 (m, 3 H), 4.26 (m, 1 H), 4.64 (m, 2 H), 4.70 (d, J=11.3 Hz, 1 H), 5.34 (t, J=9.6 Hz, 1 H), 5.59 (m, 2 H), 7.16 (d, J=8.4 Hz, 1 H), 7.52 (s, 1 H), 7.56 (s, 1 H), 8.20 (d, J=8.4 Hz, 1 H). LC-MS (Method D except gradient time increased from 2 to 4 min, retention time: 3.07 min), MS m/z 707 (M⁺+1).

Step 36H: Preparation of (1S,4R,6S,14S,18R)-7-cis-14-Cyclopentyloxy-carbonylamino-18-[2-(2-diazoacetyl)-7-methoxyquinolin-4-yloxy]-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester




Product of Step 36H

To a solution of (1S,4R,6S,14S,18R)-7-cis-18-(2-Carboxy-7-methoxyquinolin-4-yloxy)-14-cyclopentyloxy-

carbonylamino-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid ethyl ester sodium salt (1.90 mmol, product from step 36G) in 12 mL of THF cooled to 0 °C, was added 0.37 mL (2.66 mmol) of Et₃N, followed by 0.35 mL (2.66 mmol) of isobutylchloroformate. The solution was stirred 80 min, and an ethereal solution of approximately 40 mL of diazomethane added [prepared by adding 2.18 g (14.89 mmol) of 1-methyl-3-nitro-1-nitrosoguanidine in small portions to a rapidly stirring solution 40 mL of Et₂O and 30 mL of 40% aqueous KOH solution and then decanting off the CH₂N₂/Et₂O layer into the mixed anhydride reaction mixture]. The mixture was then stirred 30 min at 0 °C, 2 h at room temperature, and then concentrated in vacuo. The residue was dissolved in 200 mL of EtOAc, washed with saturated aqueous NaHCO₃ (2 X 75 mL), brine (2 X 75 mL), dried (MgSO₄), and concentrated to afford approximately 1.20 g (1.64 mmol, 86% over two steps) of the product of Step 36H which was taken directly into the next reaction (Step 36I): ¹H NMR (500 MHz, Methanol-d₄) δ 1.23 (t, J=7 Hz, 3 H), 1.30-1.73 (m, 17 H), 1.79 (m, 1 H), 1.94 (m, 1 H), 2.31 (q, J=9 Hz, 1 H), 2.53 (m, 2 H), 2.65 (m, 1 H), 3.94 (s, 3 H), 4.07 (m, 3 H), 4.25 (m, 1 H), 4.63 (m, 3 H), 5.33 (t, J=9.6 Hz, 1 H), 5.52 (s, 1 H), 5.58 (m, 1 H), 7.15 (dd, J=9.2, 2 Hz, 1 H), 7.37 (s, 1 H), 7.45 (s, 1 H), 8.14 (d, J=9.2 Hz, 1 H). LC-MS (Method D except gradient time increased from 2 to 4 min, retention time: 3.31 min), MS m/z 731 (M⁺+1).

Step 36I: Preparation of (1S,4R,6S,14S,18R)-7-cis-18-[2-(2-Bromoacetyl)-7-methoxyquinolin-4-yloxy]-14-cyclopentyloxycarbonylamino-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester


162

**Product of Step 36I**

To a solution of 1.20 g (1.64 mmol) of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-Cyclopentyloxycarbonylamino-18-[2-(2-diazoacetyl)-7-methoxyquinolin-4-yloxy]-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester (product of step 36H) in 17 mL of THF cooled to 0 °C, was added 1.25 mL of 48% aqueous HBr. The mixture was stirred 70 min, diluted with 250 mL of EtOAc, washed with saturated aqueous NaHCO₃ (2 X 75 mL), brine (2 X 75 mL), dried (MgSO₄), and concentrated to afford approximately 1.18 g (99%) of the product of step 36I as a yellow foam. The material was taken directly into the next reaction (step 36J) without further purification: ¹H NMR (500 MHz, CDCl₃) 1.20-1.82 (m, 19 H), 1.88 (m, 2 H), 2.19 (m, 3 H), 2.38 (m, 1 H), 3.06 (m, 1 H), 3.97 (s, 3 H), 4.13 (m, 4 H), 4.57 (m, 1 H), 4.93 (m, 2 H), 5.01 (m, 2 H), 5.25 (m, 1 H), 5.38 (s, 1 H), 5.52 (m, 1 H), 7.22 (dd, *J*=9.2, 2 Hz, 1 H), 7.35 (s, 1 H), 7.40 (s, 1 H), 8.08 (d, *J*=9.2 Hz, 1 H). LRMS *m/z* (M+H)⁺ 785.3.

Step 36J: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-Cyclopentyloxycarbonylamino-18-[2-(2-isopropylamino-thiazol-4-yl)-7-methoxyquinolin-4-yloxy]-2,15-dioxo-

3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester




Product of Step 36J

5 A solution of 1.18 g (1.62 mmol) of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-18-[2-(2-Bromoacetyl)-7-methoxyquinolin-4-yloxy]-14-cyclopentylloxycarbonylamino-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester (product of step 36I) and 382 mg (3.23 mmol) of isopropyl thiourea in 10 mL of isopropanol was heated to 80 °C for 1.5 h, cooled to RT, and concentrated in vacuo. The residue was diluted with 250 mL of EtOAc, washed with saturated aqueous NaHCO₃ (2 X 75 mL), brine (2 X 75 mL), dried (MgSO₄), and concentrated in vacuo. The crude residue was chromatographed over a biotage 65M column (eluted with 0% to 15% CH₂Cl₂/MeOH) to afford the desired product contaminated with isopropyl thiourea (~1.2 grams). The residue was then dissolved in a total volume of 26 mL of 75% DMSO/MeOH and purified by preparative HPLC by injection thirteen times (13 X 2 mL), afterwhich, an extractive purification was undertaken to isolate 900 mg (1.12 mmol, 69.4% yield) of the product from step 36J

from the combined HPLC fractions: Column Xterra 30X100 mm S5, 50% to 100% Solvent B/A for 10 min gradient, hold time 2 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and
5 flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined EtOAc
10 extracts were dried (MgSO₄) and concentrated in vacuo to isolate the pure product: ¹H NMR (500 MHz, Methanol-d₄) δ 1.22 (t, J=7.0 Hz, 3 H), 1.25-1.73 (m, 16 H), 1.31 (d, J=6.41 Hz, 6 H), 1.80 (m, 1 H), 1.98 (m, 1 H), 2.31 (q, J=8.8 Hz, 1 H), 2.50 (m, 2 H), 2.69 (dd, J=13, 8 Hz, 1
15 H), 3.93 (s, 3 H), 4.07 (m, 5 H), 4.29 (m, 1 H), 4.67 (m, 3 H), 5.33 (t, J=9.5 Hz, 1 H), 5.52 (s, 1 H), 5.58 (m, 1 H), 7.05 (d, J=9 Hz, 1 H), 7.37 (d, J=2 Hz, 1 H), 7.48 (s, 1 H), 7.51 (s, 1 H), 8.10 (d, J=9 Hz, 1 H). LC-MS (Method D except gradient time increased from 2 to 4 min,
20 retention time: 3.33 min), MS m/z 803 (M⁺+1).

Method A Step 36K: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-
 cis-18-{2-[2-(*tert*-Butoxycarbonylisopropylamino)thiazol-
 4-yl]-7-methoxy-quinolin-4-yloxy}-14-cyclopentyl-
 oxycarbonylamino-2,15-dioxo-3,16-diazatricyclo-
 5 [14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester




Product of Method A Step 36K

To a solution of 106 mg (0.132 mmol) of
 (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-Cyclopentylloxycarbonylamino-
 10 18-[2-(2-isopropylaminothiazol-4-yl)-7-methoxyquinolin-4-
 yloxy]-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-
 7-ene-4-carboxylic acid ethyl ester (product of step
 36J), 0.043 mL (0.311 mmol) of Et₃N, and 3.5 mg (0.00019
 mmol) of 4-DMAP in 1 mL of acetonitrile, was added 54 mg
 15 (0.245 mmol) of (BOC)₂O, and the mixture stirred for 3 h
 at RT. The residue was then diluted with MeOH until a
 total volume of 4 mL was obtained and the mixture
 purified by preparative HPLC using two injections (2 X 2
 mL), after which, an extractive purification was
 20 undertaken to isolate 22 mg (21%) of recovered
 (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-Cyclopentylloxycarbonylamino-
 18-[2-(2-isopropylaminothiazol-4-yl)-7-methoxyquinolin-4-
 yloxy]-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-
 7-ene-4-carboxylic acid ethyl ester (product of step 36J)
 25 and 60 mg (64% yield based on recovered SM) of the
 desired product from step 36K via Method A from the
 respective combined HPLC fractions. Conditions: Column

Xterra 30X100 mm S5, 45% to 100% Solvent B/A for 18 min gradient, hold time 4 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 30 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined EtOAc extracts were dried (MgSO₄) and concentrated in vacuo to isolate the respective compounds: ¹H NMR (500 MHz, Methanol-d₄) δ 1.25 (t, J=7.2 Hz, 3 H), 1.28–1.74 (m, 23 H), 1.65 (s, 9 H), 1.82 (m, 1 H), 1.98 (m, 1 H), 2.33 (q, J=8.9 Hz, 1 H), 2.51 (m, 1 H), 2.60 (m, 1 H), 2.72 (m, 1 H), 3.96 (s, 3 H), 4.08 (m, 4 H), 4.28 (d, J=9.5 Hz, 1 H), 4.70 (m, 2 H), 5.36 (m, 1 H), 5.61 (m, 3 H), 7.12 (d, J=9.2 Hz, 1 H), 7.46 (s, 1 H), 7.60 (s, 1 H), 8.17 (d, J=9.2 Hz, 1 H), 8.22 (s, 1 H). LC (Method D except gradient time increased from 2 to 4 min, retention time: 3.69 min).


Method A Step 36L: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-18-{2-[2-(*tert*-Butoxycarbonylisopropylamino)thiazol-4-yl]-7-methoxyquinolin-4-yloxy}-14-cyclopentyl-oxycarbonylamino-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid



Product of Method A Step 36L

To a solution of 60 mg (0.066 mmol) of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-18-{2-[2-(*tert*-Butoxycarbonyl-isopropylamino)thiazol-4-yl]-7-methoxyquinolin-4-yloxy}-14-cyclopentyloxy-carbonylamino-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid ethyl ester (product of step 36K, Method A) in 6.9 mL of 2:1 THF/MeOH, was added a solution of 25 mg (1.04 mmol) of LiOH in 2.3 mL of H₂O. The mixture was stirred for two days, the pH of the solution adjusted to neutral by the dropwise addition of 3N aqueous HCl solution, and the MeOH concentrated in vacuo. The solution was adjusted to pH 4 by the dropwise addition of 3N aqueous HCl solution and was extracted into EtOAc (3 X 25 mL). The combined organic layers were dried (MgSO₄), concentrated in vacuo. The residue was then diluted with 4 mL of MeOH and the mixture purified by preparative HPLC using two injections (2 X 2 mL), afterwhich, an extractive purification was undertaken to isolate 40 mg (69% yield) of the Method A product from step 36L from the respective combined HPLC fractions. HPLC/Extractive Workup Procedure: Column Xterra 30X100 mm S5, 50% to 90% Solvent B/A for 25 min gradient, hold time 3 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined EtOAc extracts were dried (MgSO₄) and concentrated in vacuo to isolate the product of step 36L: LC-MS (Method E except gradient time increased from 2 to 4 min, retention time: 3.77 min), MS *m/z* 875 (*M*⁺+1). A total of 25 mg was taken directly into Step 36M).

Method A Steps 36M-36N: Preparation of Compound 34,
 Example 36, (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-{4-Cyclopropane-
 ulfonylaminocarbonyl-18-[2-(2-isopropylaminothiazol-4-
 yl)-7-methoxyquinolin-4-yloxy]-2,15-dioxo-3,16-diaza-
 5 ricyclo-[14.3.0.0^{4,6}]nonadec-7-en-14-yl}carbamic acid
 cyclopentyl ester, from the Method A Product of Step 36L



Compound 34

10 To a solution of 25 mg (28.5 μ mol) of the
 macrocycle acid product of step 36L, (1*S*,4*R*,6*S*,14*S*,18*R*)-
 7-*cis*-18-{2-[2-(*tert*-butoxycarbonylisopropyl-
 amino)thiazol-4-yl]-7-methoxyquinolin-4-yloxy}-14-
 cyclopentyloxycarbonylamino-2,15-dioxo-3,16-diazatri-
 15 cyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid, was
 dissolved in THF (1 mL) and CDI (9 mg, 57 μ mol, 2 eq.)
 was added. The mixture was then heated to 60 °C for 45
 min. The reaction mixture was then cooled to room
 temperature, cyclopropylsulfonamide (7 mg, 57 μ mol, 2eq)
 20 and DBU (9 mg, 57 μ mol, 2eq) were added, and the
 resulting mixture was stirred at room temperature
 overnight. The resulting mixture was partitioned between
 EtOAc and water, followed by back extraction of the
 aqueous layer. The combined extracts were dried over
 25 magnesium sulfate and concentrated in vacuo. This crude
 material of step 36M was reacted with 3 mL of 67%


TFA/CH₂Cl₂ for 3 h, concentrated in vacuo, and the residue dissolved in 4 mL of MeOH. This solution was purified by preparative HPLC using two injections (2 X 2 mL), after which, an extractive purification was undertaken to isolate 11 mg of recovered carboxylic acid precursor, followed by 14 mg (56% yield, but 79% based on recovered carboxylic acid) of the product from step 36N (Method A), Compound 34, Example 36, from the respective combined HPLC fractions. HPLC/Extractive Workup

10 Procedure: Column Xterra 30X75 mm S5, 40% to 100% Solvent B/A for 30 min gradient, hold time 2 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined fractions was adjusted to neutral

15 using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined EtOAc extracts were dried (MgSO₄) and concentrated in vacuo to isolate compound 34: For


20 data, see additional Method B Preparation of Compound 34, Example 36 (Method B, Step 34M).

Method B (Steps 36L-36M): Preparation of Compound 34, Example 36, (1S,4R,6S,14S,18R)-7-cis-{4-Cyclopropanesulfonylaminocarbonyl-18-[2-(2-isopropylaminothiazol-4-yl)-7-methoxy-quinolin-4-yloxy]-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]nonadec-7-en-14-yl}carbamic acid cyclopentyl ester, from the Method A Product of Step 36J



Method A Prep. of Compound 34, Example 36

Method B Step 36L: Preparation of (1*S*,4*R*,6*S*,14*S*,18*R*)-7-
 cis-14-Cyclopentyloxycarbonylamino-18-[2-(2-isopropyl-
 5 aminothiazol-4-yl)-7-methoxy-quinolin-4-yloxy]-2,15-
 dioxo-3,16-diaza-tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-
 carboxylic acid from the Method A Product Of Step 36J



Product of Method B Step 36L


10

To a solution of 695 mg (0.866 mmol) of
 (1*S*,4*R*,6*S*,14*S*,18*R*)-7-cis-(1*S*,4*R*,6*S*,14*S*,18*R*)-7-cis-14-
 Cyclopentyloxycarbonylamino-18-[2-(2-isopropylaminothi-
 azol-4-yl)-7-methoxyquinolin-4-yloxy]-2,15-dioxo-3,16-
 15 diaza-tricyclo[14.3.0.0^{4,6}]nonadec-7-ene-4-carboxylic acid
 ethyl ester (Method A product of step 36J) in a solution

containing THF (57 mL), MeOH (14 mL), was added a solution of 311 mg (13 mmol) of LiOH in H₂O (20 mL). The mixture was stirred for 14 h, the pH of the solution adjusted to neutral by the dropwise addition of 3N aqueous HCl solution, and the MeOH concentrated in vacuo. The solution was adjusted to pH 4 by the dropwise addition of 3N aqueous HCl solution and was extracted into EtOAc (5 X 100 mL). The combined organic layers were dried (MgSO₄), concentrated in vacuo to afford 657 mg (98%) of the Method B product of Step 36L as a yellow solid: ¹H NMR (500 MHz, Methanol-d₄) δ 1.18-1.71 (m, 32 H), 1.79 (m, 1 H), 1.92 (m, 1 H), 2.30 (q, J=9 Hz, 1 H), 2.55 (m, 1 H), 2.64 (m, 1 H), 2.77 (m, 1 H), 4.02 (s, 3 H), 4.04 (m, 1 H), 4.15 (m, 2 H), 4.40 (s, 1 H), 4.71 (m, 1 H), 4.80 (m, 1 H), 5.36 (m, 1 H), 5.61 (m, 1 H), 5.81 (s, 1 H), 7.30 (d, J=9 Hz, 1 H), 7.73 (s, 2 H), 8.16 (s, 1 H), 8.29 (d, J=9 Hz, 1 H). LC-MS (Method D except gradient time increased from 2 to 4 min, retention time: 3.21 min), MS m/z 775 (M⁺+1).

20

Method B Step 36M, Preparation of Compound 34, Example 36, (1S,4R,6S,14S,18R)-7-cis-{4-Cyclopropanesulfonylaminocarbonyl-18-[2-(2-isopropylaminothiazol-4-yl)-7-methoxyquinolin-4-yloxy]-2,15-dioxo-3,16-diazatricyclo-[14.3.0.0^{4,6}]nonadec-7-en-14-yl}carbamic acid cyclopentyl ester, from the Method B product of Step 36L



Compound 34, Product of Method B Step 36M

To a solution of 66 mg (85.2 μmol) of the Method B acid product of step 36L, (1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-(1*S*,4*R*,6*S*,14*S*,18*R*)-7-*cis*-14-Cyclopentyloxycarbonylamino-18-[2-(2-isopropyl-aminothiazol-4-yl)-7-methoxy-quinolin-4-yloxy]-2,15-dioxo-3,16-diazatricyclo[14.3.0.0^{4,6}]-nonadec-7-ene-4-carboxylic acid, was dissolved in THF (1 mL) and CDI (21 mg, 130 μmol , 1.5 eq) was added. The mixture was then heated to 60 °C for 45 min. The reaction mixture was then cooled to room temperature,

10 cyclopropylsulfonamide (21 mg, 170 μmol , 2 eq) and DBU (19 mg, 130 μmol , 1.5 eq) were added, and the resulting mixture was stirred at room temperature overnight. The resulting mixture was partitioned between EtOAc and water, followed by back extraction of the aqueous layer.

15 The combined extracts were dried over magnesium sulfate and concentrated in vacuo. This residue was dissolved in 6 mL of MeOH. This solution was purified by preparative HPLC using two injections (3 X 2 mL), after which, an extractive purification was undertaken to isolate 51 mg

20 (68% yield) of the product from step 36M (Method B), Compound 34, Example 36, from the respective combined HPLC fractions. In addition, 20 mg more of compound 34 of ~80% purity (~20% yield) was isolated from earlier eluting fractions where the impurity was the carboxylic

25 acid precursor. HPLC/Isolation Procedure: Column Xterra 30X75 mm S5, 40% to 90% Solvent B/A for 32 min gradient, hold time 4 min; where Solvent A is 10% MeOH/90% H₂O with 0.1%TFA, Solvent B is 90%MeOH/10% H₂O with 0.1%TFA and flow rate is 40 ml/min). The pH of the combined

30 fractions was adjusted to neutral using 6N aqueous NaOH solution and the MeOH removed in vacuo. The solution was adjusted to pH 4 using 3N aqueous HCl solution, and repeatedly extracted with EtOAc. The combined EtOAc


extracts were dried (MgSO_4) and concentrated in vacuo to isolate the desired product of step 36N: ^1H NMR (500 MHz, Methanol- d_4) δ 0.83-1.66 (m, 23 H), 1.32 (d, $J=6$ Hz, 6 H), 1.72 (dd, $J=8.$, 5.34 Hz, 1 H), 1.81 (m, 2 H), 2.34 (m, 1 H), 2.62 (m, 2 H), 2.76 (m, 1 H), 2.89 (s, 1 H), 4.07 (m, 2 H), 4.18 (d, $J=9.77$ Hz, 1 H), 4.49 (s, 1 H), 4.68 (t, $J=8.09$ Hz, 1 H), 4.77 (m, 1 H), 5.09 (s, 1 H), 5.67 (m, 2 H), 7.19 (d, $J=9.2$ Hz, 1 H), 7.56 (s, 1 H), 7.60 (s, 1 H), 7.91 (s, 1 H), 8.21 (d, $J=9.2$ Hz, 1 H). LC-MS (Method D, retention time: 1.82 min), MS m/z 877 (M^++1).

Example 37















































































Preparation of Compounds Defined in Table 1 and Table 2 of Example 37





















































































Each of the Compounds shown in Table 1 and Table 2 of Example 37 are tripeptide acyl sulfonamides. These compounds can each be prepared from the corresponding carboxylic acid in a sulfonamide coupling process which converts the carboxylic acid to the said acyl sulfonamide. Examples of this process are found in the present application in the synthesis of Compounds 1 through 34. The carboxylic acids which are used as starting materials for the formation of the acyl sulfonamides shown in Table 1 can be prepared (for one example, see steps 9a-9i of Example 9 for the preparation of the carboxylic acid of step 9i) by the methods described in the present application and also from the teachings of International Application PCT/CA00/00353 (Publication No, WO 00/59929).

Table 1 of Example 37



Cmp	R ₁	R ₂	R ₃	R ₅	R ₄	Q
35						
36						
37						
38						
39						
40						
41						
42						















































































43						
44						
45						
46						
47						
48						
49						
50						
51						
52						
53						
54						
55						















































































56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						

70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						

83						
84						
85						
86						
87						
88						
89						
90						
91						
92						
93						
94						
95						
96						















































































97			$-\text{O}-\text{Me}$	$-\text{H}$		
98			$-\text{O}-\text{Me}$	$-\text{H}$		
99			$-\text{O}-\text{Me}$	$-\text{H}$		
100			$-\text{O}-\text{Me}$	$-\text{H}$		
101			$-\text{O}-\text{Me}$	$-\text{H}$		
102			$-\text{O}-\text{Me}$	$-\text{H}$		
103			$-\text{O}-\text{Me}$	$-\text{H}$		
104			$-\text{O}-\text{Me}$	$-\text{H}$		
105			$-\text{O}-\text{Me}$	$-\text{H}$		
106			$-\text{O}-\text{Me}$	$-\text{H}$		
107			$-\text{O}-\text{Me}$	$-\text{H}$		
108			$-\text{O}-\text{Me}$	$-\text{H}$		
109			$-\text{O}-\text{Me}$	$-\text{H}$		
110			$-\text{O}-\text{Me}$	$-\text{H}$		

111						
112						
113						
114						
115						
116						
117						
118						
119						
120						
121						
122						
123						















































































124						
125						
126						
127						
128						
129						
130						
131						
132						
133						
134						
135						
136						

137						
138						
139						
140						
141						
142						
143						
144						
145						
146						
147						
148						
149						

150						
151						
152						
153						
154						
155						
156						
157						
158						
159						
160						
161						
162						


163						
164						
165						
166						
167						
168						
169						
170						
171						
172						
173						
174						
175						

176						
177						
178						
179						
180						
181						
182						
183						
184						
185						
186						
187						
188						

189						
190						
191						
192						
193						
194						
195						
196						
197						
198						
199						
200						
201						





















































































202						
203						
204						
205						

Table 2 of Example 37

















































































5















































































Cmp	R ₁	R ₂	R ₃	R ₅	R ₄	Z
206						
207						
208						

























































209						
210						
211						
212						
213						
214						
215						
216						
217						
218						
219						
220						
221						
222						















































































223						
224						
225						
226						
227						
228						
229						
230						
231						
232						
233						
234						
235						





















































































236						
237						
238						
239						
240						
241						
242						
243						
244						
245						
246						
247						
248						

249						
250						
251						
252						
253						
254						
255						
256						
257						
258						
259						
260						
261						
262						















































































263						
264						
265						
266						
267						
268						
269						
270						
271						
272						
273						
274						
275						





















































































276			$-\text{O}-\text{Me}$	$-\text{H}$		
277			$-\text{O}-\text{Me}$	$-\text{H}$		
278			$-\text{O}-\text{Me}$	$-\text{H}$		
279			$-\text{O}-\text{Me}$	$-\text{H}$		
280			$-\text{O}-\text{Me}$	$-\text{H}$		
281			$-\text{O}-\text{Me}$	$-\text{H}$		
282			$-\text{O}-\text{Me}$	$-\text{H}$		
283			$-\text{O}-\text{Me}$	$-\text{H}$		
284			$-\text{O}-\text{Me}$	$-\text{H}$		
285			$-\text{O}-\text{Me}$	$-\text{H}$		
286			$-\text{O}-\text{Me}$	$-\text{H}$		
287			$-\text{O}-\text{Me}$	$-\text{H}$		
288			$-\text{O}-\text{Me}$	$-\text{H}$		
289			$-\text{O}-\text{Me}$	$-\text{H}$		















































































290						
291						
292						
293						
294						
295						
296						
297						
298						
299						
300						
301						
302						





















































































303						
304						
305						
306						
307						
308						
309						
310						
311						
312						
313						
314						
315						
316						















































































317			$-\text{O}-\text{Me}$	$-\text{H}$		
318			$-\text{O}-\text{Me}$	$-\text{H}$		
319			$-\text{O}-\text{Me}$	$-\text{H}$		
320			$-\text{O}-\text{Me}$	$-\text{H}$		
321			$-\text{O}-\text{Me}$	$-\text{H}$		
322			$-\text{O}-\text{Me}$	$-\text{H}$		
323			$-\text{O}-\text{Me}$	$-\text{H}$		
324			$-\text{O}-\text{Me}$	$-\text{H}$		
325			$-\text{O}-\text{Me}$	$-\text{H}$		
326			$-\text{O}-\text{Me}$	$-\text{H}$		
327			$-\text{O}-\text{Me}$	$-\text{H}$		
328			$-\text{O}-\text{Me}$	$-\text{H}$		
329			$-\text{O}-\text{Me}$	$-\text{H}$		















































































330						
331						
332						
333						
334						
335						
336						
337						
338						
339						
340						
341						
342						

























































343						
344						
345						
346						
347						
348						
349						
350						
351						
352						
353						
354						
355						
356						

357						
358						
359						
360						
361						
362						
363						
364						
365						
366						
367						
368						
369						

370						
371						
372						
373						
374						
375						
376						
377						
378						
379						
380						
381						
382						
383						

384						
385						
386						
387						
388						
389						
390						
391						
392						
393						
394						
395						
396						

397						
398						
399						
400						
401						
402						
403						
404						
405						
406						
407						
408						
409						


410			$-\text{O}-\text{Me}$	$-\text{H}$		
411			$-\text{O}-\text{Me}$	$-\text{H}$		
412			$-\text{O}-\text{Me}$	$-\text{H}$		
413			$-\text{O}-\text{Me}$	$-\text{H}$		
414			$-\text{O}-\text{Me}$	$-\text{H}$		
415			$-\text{O}-\text{Me}$	$-\text{H}$		
416			$-\text{O}-\text{Me}$	$-\text{H}$		
417			$-\text{O}-\text{Me}$	$-\text{H}$		
418			$-\text{O}-\text{Me}$	$-\text{H}$		
419			$-\text{O}-\text{Me}$	$-\text{H}$		
420			$-\text{O}-\text{Me}$	$-\text{H}$		
421			$-\text{O}-\text{Me}$	$-\text{H}$		
422			$-\text{O}-\text{Me}$	$-\text{H}$		
423			$-\text{O}-\text{Me}$	$-\text{H}$		

424						
425						
426						
427						
428						
429						
430						
431						
432						
433						
434						
435						
436						


437						
438						
439						
440						
441						
442						
443						
444						
445						
446						
447						

Preparation of Compound 441

A nonlimiting example of the above coupling process
 5 is shown for the preparation of compound 441 (where $R_1 =$



The starting material carboxylic acid
 Compound 441A can be prepared as described in
 International Application PCT/CA00/00353 (Publication No.,
 5 WO 00/59929) or Example 9 herein.



In the sulfonamide coupling step a tripeptide
 10 carboxylic acid, for example 441A (where $R_1 = \text{---}\triangle\text{---}$, $R_2 = \text{---}\begin{array}{c} \text{N} \\ \diagup \quad \diagdown \\ \text{O} \quad \text{N} \end{array}\text{---Me}$, $R_3 = \text{OMe}$, $R_5 = \text{H}$, $R_4 = \text{---NH---C(=O)---O---}\triangle\text{---}$, $Q = \text{---}\text{CH}_2\text{---CH}_2\text{---CH}_2\text{---CH}_2\text{---CH}_2\text{---CH}_2\text{---CH=CH}_2\text{---}\text{---}$)
 is dissolved in a suitable solvent, as for example THF or
 dichloroethane, under an inert atmosphere, as example
 argon or nitrogen. A suitable coupling reagent is then
 15 added. A non-limiting example of a coupling reagent is
 carbonyldiimidazole. The resulting solution is then
 stirred at room temperature for approximately 30 min and
 then heated at reflux for a period of 30 min. The
 solution is then cooled to room temperature and a
 20 sulfonamide is then added. In the present example
 cyclopropylsulfonamide is added in one portion, followed
 by the addition of a base. An example of a base used in

this transformation is DBU. The resulting reaction is stirred for 24 h, after which it is subjected to a work-up procedure. This work-up procedure can involve first diluting the reaction mixture with an organic solvent
5 such as ethyl acetate then washing the resulting mixture with an aqueous solution as for example an aqueous solution buffered to a pH of four. The organic phase is then separated and washed with brine. The organic extract is then dried (MgSO_4) and concentrated under
10 vacuum to provide a crude product. The crude product can be purified using conventional practices, such as but not limited to, flash chromatography, preparative thin layer chromatography (TLC) or reverse phase chromatography to provide the desire acylsulfonamide compound 441.

15

Example 38

Biological Studies

20 Recombinant HCV NS3/4A protease complex FRET peptide assay

The purpose of this in vitro assay was to measure the inhibition of HCV NS3 protease complexes, derived
25 from the BMS, H77C or J4L6S strains, as described below, by compounds of the present invention. This assay provides an indication of how effective compounds of the present invention would be in inhibiting HCV proteolytic activity.

30

Serum from an HCV-infected patient was obtained from Dr. T. Wright, San Francisco Hospital. An engineered full-length cDNA (compliment deoxyribonucleic acid) template of the HCV genome (BMS strain) was constructed

from DNA fragments obtained by reverse transcription-PCR (RT-PCR) of serum RNA (ribonucleic acid) and using primers selected on the basis of homology between other genotype 1a strains. From the determination of the entire genome sequence, a genotype 1a was assigned to the HCV isolate according to the classification of Simmonds et al. (See P Simmonds, KA Rose, S Graham, SW Chan, F McOmish, BC Dow, EA Follett, PL Yap and H Marsden, J. Clin. Microbiol., 31(6), 1493-1503 (1993)). The amino acid sequence of the nonstructural region, NS2-5B, was shown to be >97% identical to HCV genotype 1a (H77C) and 87% identical to genotype 1b (J4L6S). The infectious clones, H77C (1a genotype) and J4L6S (1b genotype) were obtained from R. Purcell (NIH) and the sequences are published in Genbank (AAB67036, see Yanagi, M., Purcell, R.H., Emerson, S.U. and Bukh, J. Proc. Natl. Acad. Sci. U.S.A. 94(16), 8738-8743 (1997); AF054247, see Yanagi, M., St Claire, M., Shapiro, M., Emerson, S.U., Purcell, R.H. and Bukh, J, Virology 244 (1), 161-172. (1998)).

The BMS, H77C and J4L6S strains were used for production of recombinant NS3/4A protease complexes. DNA encoding the recombinant HCV NS3/4A protease complex (amino acids 1027 to 1711) for these strains were manipulated as described by P. Gallinari et al. (see Gallinari P, Paolini C, Brennan D, Nardi C, Steinkuhler C, De Francesco R. Biochemistry. 38(17):5620-32, (1999)). Briefly, a three-lysine solubilizing tail was added at the 3'-end of the NS4A coding region. The cysteine in the P1 position of the NS4A-NS4B cleavage site (amino acid 1711) was changed to a glycine to avoid the proteolytic cleavage of the lysine tag. Furthermore, a

cysteine to serine mutation was introduced by PCR at amino acid position 1454 to prevent the autolytic cleavage in the NS3 helicase domain. The variant DNA fragment was cloned in the pET21b bacterial expression vector (Novagen) and the NS3/4A complex was expressed in *Escherichia coli* strain BL21 (DE3) (Invitrogen) following the protocol described by P. Gallinari et al. (see Gallinari P, Brennan D, Nardi C, Brunetti M, Tomei L, Steinkuhler C, De Francesco R., J Virol. 72(8):6758-69 (1998)) with modifications. Briefly, NS3/4A expression was induced with 0.5mM Isopropyl β -D-1-thiogalactopyranoside (IPTG) for 22hr at 20°C. A typical fermentation (10L) yielded approximately 80g of wet cell paste. The cells were resuspended in lysis buffer (10mL/g) consisting of 25mM N-(2-Hydroxyethyl)Piperazine-N'-(2-Ethane Sulfonic acid) (HEPES), pH7.5, 20% glycerol, 500mM Sodium Chloride (NaCl), 0.5% Triton-X100, 1ug/ml lysozyme, 5mM Magnesium Chloride (MgCl₂), 1ug/ml DnaseI, 5mM β -Mercaptoethanol (β ME), Protease inhibitor - Ethylenediamine Tetraacetic acid (EDTA) free (Roche), homogenized and incubated for 20 mins at 4°C. The homogenate was sonicated and clarified by ultracentrifugation at 235000g for 1hr at 4°C. Imidazole was added to the supernatant to a final concentration of 15mM and the pH adjusted to 8.0. The crude protein extract was loaded on a Nickel-Nitrilotriacetic acid (Ni-NTA) column pre-equilibrated with buffer B (25mM HEPES, pH8.0, 20% glycerol, 500mM NaCl, 0.5% Triton-X100, 15mM imidazole, 5mM β ME). The sample was loaded at a flow rate of 1mL/min. The column was washed with 15 column volumes of buffer C (same as buffer B except with 0.2% Triton-X100). The protein was eluted with 5 column

volumes of buffer D (same as buffer C except with 200mM Imidazole).

NS3/4A protease complex-containing fractions were
5 pooled and loaded on a desalting column Superdex-S200
pre-equilibrated with buffer D (25mM HEPES, pH7.5, 20%
glycerol, 300mM NaCl, 0.2% Triton-X100, 10mM β ME).
Sample was loaded at a flow rate of 1mL/min. NS3/4A
protease complex-containing fractions were pooled and
10 concentrated to approximately 0.5mg/ml. The purity of
the NS3/4A protease complexes, derived from the BMS, H77C
and J4L6S strains, were judged to be greater than 90% by
SDS-PAGE and mass spectrometry analyses.

15 The enzyme was stored at -80°C, thawed on ice and
diluted prior to use in assay buffer. The substrate used
for the NS3/4A protease assay was RET S1 (Resonance
Energy Transfer Depsipeptide Substrate; AnaSpec, Inc. cat
22991) (FRET peptide), described by Taliani et al. in
20 Anal. Biochem. 240(2):60-67 (1996). The sequence of this
peptide is loosely based on the NS4A/NS4B natural
cleavage site except there is an ester linkage rather
than an amide bond at the cleavage site. The peptide
substrate was incubated with one of the three recombinant
25 NS3/4A complexes, in the absence or presence of a
compound of the present invention, and the formation of
fluorescent reaction product was followed in real time
using a Cytofluor Series 4000.

30 The reagents were as follow: HEPES and Glycerol
(Ultrapure) were obtained from GIBCO-BRL. Dimethyl
Sulfoxide (DMSO) and bovine serum albumin (BSA) were
obtained from Sigma. β -Mercaptoethanol was obtained from

Bio Rad: Assay buffer: 50mM HEPES, pH7.5; 0.15M NaCl; 0.1% Triton; 15% Glycerol; 10mM β ME. Substrate: 2 μ M final concentration (from a 2mM stock solution in DMSO stored at -20°C). HCV NS3/4A type 1a (1b), 2-3 nM final concentration (from a 5 μ M stock solution in 25mM HEPES, pH7.5, 20% glycerol, 300mM NaCl, 0.2% Triton-X100, 10mM β ME). For compounds with potencies approaching the assay limit, the assay was made more sensitive by adding 50 μ g/ml BSA to the assay buffer and reducing the end protease concentration to 300 pM.

The assay was performed in a 96-well polystyrene black plate from Falcon. Each well contained 25 μ l NS3/4A protease complex in assay buffer, 50 μ l of a compound of the present invention in 10% DMSO/assay buffer and 25 μ l substrate in assay buffer. A control (no compound) was also prepared on the same assay plate. The enzyme complex was mixed with compound or control solution for 1 min before initiating the enzymatic reaction by the addition of substrate. The assay plate was read immediately using the Cytofluor Series 4000 (Perspective Biosystems). The instrument was set to read an emission of 340nm and excitation of 490nm at 25°C. Reactions were generally followed for approximately 15 minutes.

The percent inhibition was calculated with the following equation:

$$100 - [(\delta F_{inh}/\delta F_{con}) \times 100]$$

where δF is the change in fluorescence over the linear range of the curve. A non-linear curve fit was applied to

the inhibition-concentration data, and the 50% effective concentration (IC₅₀) was calculated by the use of Excel Xl-fit software using the equation, $y=A+(B-A)/(1+((C/x)^D))$.

5

All of the compounds tested were found to have IC₅₀s of 9 μM or less. Further, compounds of the present invention, which were tested against more than one type of NS3/4A complex, were found to have similar inhibitory properties though the compounds uniformly demonstrated greater potency against the 1b strains as compared to the 1a strains.

Specificity Assays

15

The specificity assays were performed to demonstrate the selectivity of the compounds of the present invention in inhibiting HCV NS3/4A protease as compared to other serine or cysteine proteases.

20

The specificities of compounds of the present invention were determined against a variety of serine proteases: human neutrophil elastase (HNE), porcine pancreatic elastase (PPE) and human pancreatic chymotrypsin and one cysteine protease: human liver cathepsin B. In all cases a 96-well plate format protocol using colorimetric p-nitroaniline (pNA) substrate specific for each enzyme was used as described previously (U.S. Patent 6,323,180) with some modifications to the serine protease assays. All enzymes were purchased from Sigma while the substrates were from Bachem.

30

Each assay included a 2hr enzyme-inhibitor pre-incubation at RT followed by addition of substrate and hydrolysis to ~30% conversion as measured on a Spectramax Pro microplate reader. Compound concentrations varied
5 from 100 to 0.4 μ M depending on their potency.

The final conditions for each assay were as follows:

50mM Tris(hydroxymethyl)aminomethane hydrochloride
10 (Tris-HCl) pH8, 0.5M Sodium Sulfate (Na_2SO_4), 50mM NaCl, 0.1mM EDTA, 3% DMSO, 0.01% Tween-20 with:
133 μ M succ-AAA-pNA and 20nM HNE or 8nM PPE; 100 μ M succ-AAPF-pNA and 250pM Chymotrypsin.

15 100mM NaHPO_4 (Sodium Hydrogen Phosphate) pH 6, 0.1mM EDTA, 3% DMSO, 1mM TCEP (Tris(2-carboxyethyl)phosphine hydrochloride), 0.01% Tween-20, 30 μ M Z-FR-pNA and 5nM Cathepsin B (enzyme stock
20 activated in buffer containing 20mM TCEP before use).

The percentage of inhibition was calculated using the formula:

25
$$[1 - ((\text{UV}_{\text{inh}} - \text{UV}_{\text{blank}}) / (\text{UV}_{\text{ctl}} - \text{UV}_{\text{blank}}))] \times 100$$

A non-linear curve fit was applied to the inhibition-concentration data, and the 50% effective concentration (IC_{50}) was calculated by the use of Excel
30 Xl-fit software.

HCV Replicon Cell-based Assay

An HCV replicon whole cell system was established as described by Lohmann V, Korner F, Koch J, Herian U, Theilmann L, Bartenschlager R., Science 285(5424):110-3 (1999). This system enabled us to evaluate the effects of our HCV Protease compounds on HCV RNA replication. Briefly, using the HCV strain 1B sequence described in the Lohmann paper (Assession number:AJ238799), an HCV cDNA was generated encoding the 5' internal ribosome entry site (IRES), the neomycin resistance gene, the EMCV (encephalomyocarditis viurs)-IRES and the HCV nonstructural proteins, NS3-NS5B, and 3' non-translated region (NTR). In vitro transcripts of the cDNA were transfected into the human hepatoma cell line, Huh7. Selection for cells constitutively expressing the HCV replicon was achieved in the presence of the selectable marker, neomycin (G418). Resulting cell lines were characterized for positive and negative strand RNA production and protein production over time.

Huh7 cells, constitutively expressing the HCV replicon, were grown in Dulbecco's Modified Eagle Media (DMEM) containing 10% Fetal calf serum (FCS) and 1mg/ml G418 (Gibco-BRL). Cells were seeded the night before (1.5×10^4 cells/well) in 96-well tissue-culture sterile plates. Compound and no compound controls were prepared in DMEM containing 4% FCS, 1:100 Penicillin / Streptomycin, 1:100 L-glutamine and 5% DMSO in the dilution plate (0.5% DMSO final concentration in the assay). Compound / DMSO mixes were added to the cells and incubated for 4 days at 37°C. After 4 days, plates were rinsed thoroughly with Phosphate-Buffered Saline (PBS) (3 times 150µl). The cells were lysed with 25µl of a lysis assay reagent containing the FRET peptide (RET S1, as

described for the in vitro enzyme assay). The lysis assay reagent was made from 5X cell Luciferase cell culture lysis reagent (Promega #E153A) diluted to 1X with distilled water, NaCl added to 150 mM final, the FRET peptide diluted to 10 μ M final from a 2 mM stock in 100% DMSO. The plate was then placed into the Cytofluor 4000 instrument which had been set to 340nm excitation / 490 emission, automatic mode for 21 cycles and the plate read in a kinetic mode. EC₅₀ determinations were carried out as described for the IC₅₀ determinations.

Two different secondary assays were used to confirm EC₅₀ determinations from the replicon FRET assay. These included a quantitative RNA assay and a transient luciferase cell reporter assay. For the quantitative RNA assay, compound / no compound controls were incubated with the cells as described for the replicon FRET assay. After 4 days, cells were lyzed using the Rneasy kit (Qiagen). Purified total RNA was normalized using RiboGreen (Jones LJ, Yue ST, Cheung CY, Singer VL, Anal. Chem., 265(2):368-74 (1998)) and relative quantitation of HCV RNA expression assessed using the Taqman procedure (Kolykhalov AA, Mihalik K, Feinstone SM, Rice CM, Journal of Virology 74, 2046-2051 (2000)) and the Platinum Quantitative RT-PCR Thermoscript One-Step kit (Invitrogen cat # 11731-015). Briefly, RNA made to a volume of 5 μ l (\leq 1ng) was added to a 20 μ l Ready-Mix containing the following: 1.25X Thermoscript reaction mix (containing Magnesium Sulfate and 2-deoxynucleoside 5'-triphosphates (dNTPs)), 3mM dNTPs, 200nM forward primer (sequence: 5'-gggagagccatagtggtctgc-3'), 600nM reverse primer (5'-cccaaattctccaggcattga-3'), 100nM probe (5'-6-FAM-cggaattgccaggacgaccgg-BHQ-1-3') (FAM: Fluorescein-

aminoethyl amidite; BHQ: Black Hole Quencher), 1 μ M Rox reference dye (Invitrogen cat # 12223-012) and Thermoscript Plus Platinum Taq polymerase mixture. All primers were designed with ABI Prism 7700 software and
5 obtained from Biosearch Technologies, Novato, CA. Samples containing known concentrations of HCV RNA transcript were run as standards. Using the following cycling protocol (50°C, 30 min; 95°C, 5 min; 40 cycles of 95°C, 15 sec, 60°C, 1 min), HCV RNA expression was
10 quantitated as described in the Perkin Elmer manual using the ABI Prism 7700 Sequence Detector.

The luciferase reporter assay was also used to confirm compound potency in the replicon. Utilization of
15 a replicon luciferase reporter assay was first described by Krieger et al (Krieger N, Lohmann V, and Bartenschlager R, J. Virol. 75(10):4614-4624 (2001)). The replicon construct described for our FRET assay was modified by replacing the resistance gene neomycin with
20 the Blastidicin-resistance gene fused to the N-terminus of the humanized form of Renilla luciferase (restriction sites AscI / PmeI used for the subcloning). The adaptive mutation at position 1179 (serine to isoleucine) was also introduced (Blight KJ, Kolykhalov, AA, Rice, CM, Science
25 290(5498):1972-1974). The luciferase reporter assay was set up by seeding huh7 cells the night before at a density of 2 x 10⁶ cells per T75 flask. Cells were washed the next day with 7.5ml Opti-MEM. Following the Invitrogen protocol, 40 μ l DMRIE-C was vortexed with 5 ml
30 Opti-MEM before adding 5 μ g HCV reporter replicon RNA. The mix was added to the washed huh7 cells and left for 4 hours at 37°C. In the mean time, serial compound

dilutions and no compound controls were prepared in DMEM containing 10% FCS and 5% DMSO in the dilution plate (0.5% DMSO final concentration in the assay). Compound / DMSO mixes were added to each well of a 24-well plate.

5 After 4 hours, the transfection mix was aspirated, and cells washed with 5ml of Opti-MEM before trypsinization. Trypsinized cells were resuspended in 10% DMEM and seeded at 2×10^4 cells/well in the 24-well plates containing compound or no compound controls. Plates were incubated

10 for 4 days. After 4 days, media was removed and cells washed with PBS. 100 μ l 1x Renilla Luciferase Lysis Buffer (Promega) was immediately added to each well and the plates either frozen at -80°C for later analysis, or assayed after 15 mins of lysis. Lysate (40 μ l) from each

15 well was transferred to a 96-well black plate (clear bottom) followed by 200 μ l 1x Renilla Luciferase assay substrate. Plates were read immediately on a Packard TopCount NXT using a luminescence program.

20 The percentage inhibition was calculated using the formula below:

$$\% \text{ control} = \frac{\text{average luciferase signal in experimental wells (+ compound)}}{\text{average luciferase signal in DMSO control wells (- compound)}}$$

25

The values were graphed and analyzed using XLFit to obtain the EC₅₀ value.

Biological Examples

30

The following Table lists compounds of the present invention which were assayed in vitro according to the method previously described. These IC₅₀ ranges used in

this table are: A is $<50\mu\text{M}$; B is $<5\mu\text{M}$; C is $<0.5\mu\text{M}$; D is $<0.05\mu\text{M}$. All of the compounds of the present invention that were tested were found to have IC_{50}s of $9\mu\text{M}$ or less against the BMS HCV NS3/4A complex. Preferred compounds

5 had IC_{50}s of $0.021\mu\text{M}$, as was found for Compound 6, or less. Further, compounds of the present invention, which were tested against more than one type of NS3/4A complex, were found to have similar inhibitory properties; somewhat better inhibitory properties were exhibited

10 against strain J4L6S. Representative compounds of the present invention were also assessed in the HCV replicon cell assay the results of which are provided in the Table. These EC_{50} ranges used in this table are: A is $<50\mu\text{M}$; B is $<5\mu\text{M}$; C is $<0.5\mu\text{M}$; D is $<0.05\mu\text{M}$. Compounds

15 of the present invention were found to have EC_{50}s of about $13.3\mu\text{M}$ or less. Preferred compounds have EC_{50}s of about $0.124\mu\text{M}$, as found for Compound 6, or less. More preferred compounds of the present invention were found to have EC_{50}s of $0.01\mu\text{M}$ or less.

20

In the specificity assays, the same compound was found to have the following activity: HLE = $30\mu\text{M}$; PPE $> 100\mu\text{M}$; Chymotrypsin $> 100\mu\text{M}$; Cathepsin B $> 100\mu\text{M}$. These results indicate this family of compounds are highly


25 specific for the NS3 protease and many of these members inhibit HCV replicon replication.

Compound	IC50	EC50	Compound	IC50	EC50
1	B	A	2	B	
3	B	A	4	A	
5	D	D	6	D	C
7	D	D	8	D	C
9	D	D	10	D	D
11	D	D	12	C	C
13	D	D	14	D	D
15	D	D	16	D	D
17	D	C	18	D	D
19	D	D	20	D	D
21	D	D	22	D	D
23	D	D	24	D	D
25	D	D	26	D	D
27	D	D	28	D	D
29	D	D	30	D	B
31	D	D	32	D	D
33	D	D	34	D	D

CLAIMS

What is claimed is:

1. A compound of formula I, including pharmaceutically acceptable salts thereof,



wherein:

(a) R_1 is C_{1-6} alkyl or unsubstituted C_{3-7} cycloalkyl;

(b) R_2 is H, halo, C_{1-6} alkyl, C_{1-6} alkoxy, C_{3-6} cycloalkoxy, C_6 or 10 aryl or heterocycle; wherein heterocycle is a five-, six-, or seven-membered saturated or unsaturated heterocycle containing from one to four heteroatoms selected from nitrogen, oxygen and sulfur; said aryl or heterocycle being substituted with R_{24} ;

wherein R_{24} is H, halo, C_{1-6} alkyl, $-N(R_a)(R_b)$,

5 $-N(R_a)C(O)(OR_d)$, $-N(R_a)C(O)(R_b)$, $-NHC(O)(R_a)(R_c)$ or $-N(R_a)SO_2R_b$; wherein R_a and R_b are independently H, C_{1-6} alkyl or C_{3-6} cycloalkyl; R_c is H, C_{1-6} alkyl, C_{3-6} cycloalkyl or C_{1-6} alkoxy; R_d is C_{1-6} alkyl or C_{3-6} cycloalkyl;

(c) R_3 is H, halo, CF_3 , C_{1-6} alkoxy or C_{3-6} cycloalkoxy;


10 (d) R_4 is NH_2 , or $-NH-R_{31}$; wherein R_{31} is $-C(O)-R_{32}$, $C(O)-NHR_{32}$ or $C(O)-OR_{32}$, wherein R_{32} is C_{1-6} alkyl optionally substituted with halo, $-(CH_2)_p-C_{3-7}$ cycloalkyl or a tetrahydrofuran ring linked through the C3 or C4 position of the ring;
15 wherein p is 0-6;

(e) R_5 is H; and

20 (f) Q is a three to nine atom saturated or unsaturated alkylene chain optionally containing one to three heteroatoms independently selected from O or $S(O)_m$; wherein m is 0, 1 or 2.


2. A compound of Claim 1 having the formula

222



3. A compound of Claim 2 wherein R_1 is cyclopropyl.
- 5 4. A compound of Claim 2 wherein R_2 is phenyl.
5. A compound of Claim 2 wherein R_3 is methoxy.
- 10 6. A compound of Claim 2 wherein R_4 is NH_2 or *tert*-butoxycarbonylNH-.
7. A compound of claim 2 wherein Q is $-(\text{CH}_2)_n\text{CH}=\text{C}-$ and n is 1-5.
- 15 8. A compound of claim 7 wherein n is 3 or 4.
9. A compound of claim 1 having the formula

223



wherein:

5 (a) R_4 is NH_2 or tert-butoxycarbonylNH-;



(b) Q is $-(\text{CH}_2)_n\text{CH}=\text{C}-$; and



(c) n is 1-5.

10

10. A compound of claim 9 wherein n is 3 or 4.

11. A compound of claim 2 wherein:

15 R_2 is  or  ;

R_4 is  or  ;

and R_1 is cyclopropyl.

20

12. A pharmaceutical composition, comprising a compound of claims 1-11, and a pharmaceutically acceptable carrier.
- 5 13. A method of inhibiting HCV NS3 protease which comprises administering to a mammal in need of such treatment a therapeutically effective amount of a compound of claims 1-11.
- 10 14. A method of treating an HCV infection, in a mammal in need thereof, comprising the administration to said mammal of a therapeutically effective amount of a compound of claims 1-11.

